

# 2020 APHRS/HRS expert consensus statement on the investigation of decedents with sudden unexplained death and patients with sudden cardiac arrest, and of their families.

**Title:** 2020 APHRS/HRS expert consensus statement on the investigation of decedents with sudden unexplained death and patients with sudden cardiac arrest, and of their families.

**Publication year:** 2021

**Journal:** Heart rhythm

**Authors:** Waikato Clinical School, Faculty of Medicine and Health Science, The University of Auckland, Hamilton, New Zealand.

**Society:** European Respiratory Society

**Origin country:** International (Europe)

**Target countries:** Europe

**DOI:** 10.1016/j.hrthm.2020.10.010

**PMID:** 33091602

**Source:** <https://pubmed.ncbi.nlm.nih.gov/33091602/>

Heart Rhythm Heart Rhythm 319 nihpa 101200317 Heart rhythm 1547-5271 1556-3871  
pmc-is-collection-domain yes pmc-collection-title NIHPA Author Manuscripts PMC8194370  
PMC8194370.1 8194370 8194370 NIHMS1704401 33091602 10.1016/j.hrthm.2020.10.010  
NIHMS1704401 NIHPA1704401 1 Article 2020 APHRS/HRS expert consensus statement on the  
investigation of decedents with sudden unexplained death and patients with sudden cardiac arrest,  
and of their families Stiles Martin K. MBChB, PhD, FHRS (APHRS Chair) 1 ^ Wilde Arthur A. M.  
MD, PhD (HRS Chair) 2 ^ ¥ II Abrams Dominic J. MBBS, MD, MBA 3 \* Ackerman Michael J. MD,  
PhD 4 † Albert Christine M. MD, MPH, FHRS 5 † Behr Elijah R. MA, MBBS, MD 6 ‡ ¥ II Chugh  
Sumeet S. MD, FHRS 5 † Cornel Martina C. MD, PhD 7 § Gardner Karen MPH, PhD 8 Δ Ingles  
Jodie GradDipGenCouns PhD, MPH, FHRS 9 # James Cynthia A. ScM, PhD, CGC 10 # Juang  
Jyh-Ming Jimmy MD, PhD 11 ¶ Kääb Stefan MD, PhD 12 ‡ ¥ Kaufman Elizabeth S. MD, FHRS 13 †  
Krahn Andrew D. MD, FHRS 14 † Lubitz Steven A. MD, MPH 15 † MacLeod Heather MS, CGC 16  
# Morillo Carlos A. MD, FHRS 17 \*\* Nademanee Koonlawee MD, FHRS, CCDS 18 ¶ Probst  
Vincent MD, PhD 19 ‡ ¥ II Saarel Elizabeth V. MD, FHRS, CEPS-P 20 \* Sacilotto Luciana MD, PhD  
21 \*\* Semsarian Christopher MBBS, MPH, PhD, FHRS 9 ¶ Sheppard Mary N. MB, BCH, BAO, MD,  
FRCPath 6 †† II Shimizu Wataru MD, PhD 22 ¶ Skinner Jonathan R. MBChB, MD, FHRS 23 \*  
Tfelt-Hansen Jacob MD, DMSc 24 ‡ ¥ II Wang Dao Wu MD, PhD 25 ¶ 1 Waikato Clinical School,  
Faculty of Medicine and Health Science, The University of Auckland, Hamilton, New Zealand 2  
Amsterdam University Medical Center, University of Amsterdam, Heart Center, Department of  
Clinical and Experimental Cardiology, Amsterdam, the Netherlands 3 Boston Children's Hospital,  
Boston, Massachusetts, USA 4 Mayo Clinic, Rochester, Minnesota, USA 5 Cedars-Sinai Medical  
Center, Los Angeles, California, USA 6 Cardiovascular Clinical Academic Group, Molecular and  
Clinical Sciences Institute, St George's, University of London, and St George's University Hospitals  
NHS Foundation Trust, London, United Kingdom 7 Amsterdam University Medical Center, Vrije  
Universiteit Amsterdam, Clinical Genetics, Amsterdam Public Health Research Institute,  
Amsterdam, the Netherlands 8 University of New South Wales, Canberra, Australia 9 Agnes  
Ginges Centre for Molecular Cardiology at Centenary Institute, The University of Sydney, Sydney,  
Australia 10 Johns Hopkins University, Baltimore, Maryland, USA 11 Cardiovascular Center and  
Division of Cardiology, Department of Internal Medicine, National Taiwan University Hospital and  
National Taiwan University College of Medicine, Taipei, Taiwan 12 Department of Medicine I,  
University Hospital, LMU Munich, Munich, Germany 13 MetroHealth Campus, Case Western  
Reserve University, Cleveland, Ohio, USA 14 The University of British Columbia, Vancouver, British  
Columbia, Canada 15 Massachusetts General Hospital, Boston, Massachusetts, USA 16 Data  
Coordinating Center for the Sudden Death in the Young Case Registry, Okemos, Michigan, USA 17

Libin Cardiovascular Institute, Calgary, Alberta, Canada 18 Chulalongkorn University, Faculty of Medicine, and Pacific Rim Electrophysiology Research Institute at Bumrungrad Hospital, Bangkok, Thailand 19 CHU de Nantes, Nantes, France 20 Cleveland Clinic Lerner College of Cardiology at Case Western Reserve University, Cleveland, Ohio, and St Luke's Medical Center, Boise, Idaho, USA 21 Heart Institute, University of São Paulo Medical School, São Paulo, Brazil 22 Department of Cardiovascular Medicine, Nippon Medical School, Tokyo, Japan 23 Cardiac Inherited Disease Group, Starship Hospital, Auckland, New Zealand 24 Department of Forensic Medicine, Faculty of Medical Sciences, Rigshospitalet, Copenhagen, Denmark 25 The First Affiliated Hospital of Nanjing Medical University, Nanjing, China. \* Representative of the Pediatric and Congenital Electrophysiology Society (PACES)

Representative of the Heart Rhythm Society (HRS)

Representative of the European Heart Rhythm Association (EHRA)

Representative of the European Society of Human Genetics (ESHG)

Representative of the Asia Pacific Heart Rhythm Society (APHRS)

Representative of the National Society of Genetic Counselors (NSGC)

Representative of the Latin American Heart Rhythm Society (LAHRS)

Representative of the Association for European Cardiovascular Pathology (AECVP)

Member of the European Cardiac Arrhythmia Genetics (ECGen) Focus Group of the European Heart Rhythm Association (EHRA)

Member of the European Reference Network for Rare and Low Prevalence Complex Diseases of the Heart: ERN GUARD-Heart

Martin K. Stiles, MBChB, PhD, FHRS, and Arthur A. M. Wilde, MD, PhD, are co-first authors.

19 10 2020 1 2021 18 1 383587 e1 e50 03 06 2021 11 06 2021 11 06 2021 02 08 2025  
<https://www.elsevier.com/about/our-business/policies/copyright/permissions> Permissions: Multiple copies, modification, alteration, enhancement, and/or distribution of this document are not permitted without the express permission of the Heart Rhythm Society. Instructions for obtaining permission are located at <https://www.elsevier.com/about/our-business/policies/copyright/permissions>.  
<https://creativecommons.org/licenses/by/4.0/> This article is published under the Creative Commons CC-BY license ( <http://creativecommons.org/licenses/by/4.0/> ). This international multidisciplinary document intends to provide clinicians with evidence-based practical patient-centered recommendations for evaluating patients with evidence-based practical patient-centered recommendations for evaluating patients and decedents with (aborted) sudden cardiac arrest and their families. The document includes a framework for the investigation of the family allowing steps to be taken, should an inherited condition be found, to minimize further events in affected relatives. Integral to the process is counseling of the patients and families, not only because of the emotionally charged subject, but because finding (or not finding) the cause of the arrest may influence management of family members. The formation of multidisciplinary teams is essential to provide a complete service to the patients and their families, and the varied expertise of the writing committee was formulated to reflect this need. The document sections were divided up and drafted by the writing committee members according to their expertise. The recommendations represent the consensus opinion of the entire writing committee, graded by Class of Recommendation and Level of Evidence. The recommendations were opened for public comment and reviewed by the relevant scientific and clinical document committees of the Asia Pacific Heart Rhythm Society (APHRS) and the Heart Rhythm Society (HRS); the document underwent external review and endorsement by the partner and collaborating societies. While the recommendations are for optimal

care, it is recognized that not all resources will be available to all clinicians. Nevertheless, this document articulates the evaluation that the clinician should aspire to provide for patients with sudden cardiac arrest, decedents with sudden unexplained death, and their families.

This expert consensus statement represents an international multidisciplinary effort led by the Asia Pacific Heart Rhythm Society (APHRS), in partnership with the Heart Rhythm Society (HRS) and in collaboration with the Association for European Cardiovascular Pathology (AECVP), the European Heart Rhythm Association (EHRA), the European Society of Human Genetics (ESHG), the Latin American Heart Rhythm Society (LAHRS), the National Society of Genetic Counselors (NSGC) (USA), the Pediatric and Congenital Electrophysiology Society (PACES), and the European Reference Network for Rare and Low Prevalence Complex Diseases of the Heart: ERN GUARD-Heart. The intent is to provide clinicians with practical patient-centered recommendations for evaluating patients with sudden cardiac arrest (SCA), decedents with sudden cardiac death (SCD), and their families, based on all available evidence. Although the recommendations are for optimal care, the writing committee recognizes that not all resources will be available to all clinicians. Nevertheless, this document articulates the evaluation that the clinician should aspire to provide.

The writing committee consisted of internationally recognized experts from 14 countries in the fields of cardiac electrophysiology, cardiology, pediatric cardiology, genetic counseling, community genetics and public health genomics, and cardiac pathology, representing APHRS, HRS, AECVP, EHRA, ESHG, LAHRS, NSGC, PACES, and ERN GUARD-Heart and selected according to each society's procedures. In addition, a patient representative was chosen to provide a consumer viewpoint. Each partner society nominated a chair, who did not have relevant relationships with industry and other entities (RWIs). In accordance with the APHRS policies, disclosure of any RWIs was required from the writing committee members ( Appendix 1 ) and from the peer reviewers ( Appendix 2 ); of the 28 committee members, 23 (82%) had no relevant RWIs. Recommendations were drafted by the writing committee members who did not have relevant RWIs.

After development of a preliminary outline, committee members were given writing assignments and a schedule of conference calls. Writing committee members conducted a comprehensive evidence search using MEDLINE/PubMed, Embase, and the Cochrane Library and summarized the evidence in standardized tables ( Appendix 3 ), with attention to the study type, size, inclusion criteria, and key findings. The writing committee reviewed evidence and established consensus to generate recommendations, which are presented in a modular knowledge chunk format, with each chunk including a table of recommendations, a brief synopsis, recommendation-specific supportive text, flow diagrams or tables as appropriate, and references. Recommendations were formulated according to the American College of Cardiology (ACC)/American Heart Association (AHA) Class of Recommendation (COR) and Level of Evidence (LOE) system 1 ( Table 1 ) and were subject to a period of public comment. The COR indicates the strength of a recommendation based on assessment of the estimated benefits and risks; LOE rates the quality of evidence that supports the recommendation based on type, quantity, and consistency of data from clinical trials and other sources. Case reports were not used to support recommendations. The threshold for consensus was considered as 80% or higher agreement. The 74 recommendations were balloted by the 28 writing committee members and approved by an average of 94%. A quorum of two-thirds of the writing committee was met for all votes.

After review by the entire writing committee, the recommendations were opened for public comment; the draft document was reviewed by the International Scientific Document Writing Committee of the APHRS and the Scientific and Clinical Documents Committee of the HRS and was revised prior to external review. The document underwent external peer review by reviewers appointed by the APHRS and HRS and each of the collaborating societies. After subsequent revisions and endorsement by the participating societies, the document was ready for publication.

This document provides a framework for the investigation of 1) patients with SCA, 2) decedents with sudden unexplained death (SUD), and 3) families of both SCA survivors and SUD victims, as

many conditions responsible for the cardiac arrest or unexplained death may be familial. Identifying a cause is important for preventing further events in the family, should an inherited condition be found. Integral to the process is the counseling of the patients and families, not only because of the emotionally charged subject, but because finding (or not finding) the cause of the arrest may influence the futures of the family members. The disciplines of cardiology, pediatrics, radiology, pathology, counseling, psychology, and genetics are all involved in this process. Therefore, the formation of multidisciplinary teams is essential to provide a complete service to the patients and their families.

While this document endeavors to provide clinicians with practical recommendations for evaluating patients with SCA, decedents with SUD, and their families, the best approach will vary with the situation and will be influenced by, for example, the subject's age and results of initial testing. Although some of the recommendations do specify an age cutoff, it is recognized that this age is somewhat arbitrary and may not be always appropriate for the disease being investigated for or the demographics of the patient's country. Nevertheless, where an age is specified in a recommendation, it has passed the consensus voting of the writing group. Referral to a center with a multidisciplinary team experienced in such evaluations is recommended because it can facilitate navigation of these complexities. A multidisciplinary team can also help organize interval follow-up evaluations for SCA survivors and their family members. Repeated interval follow-up can reveal important new clinical data and allows for integration of new knowledge into the continued evaluation and care of these patients. The writing committee members recognize that not all investigative modalities recommended will be available in all circumstances; however, this document is an attempt to outline an approach to which the clinician should aspire.

Table 2 lists pertinent guidelines and consensus statements that the writing committee considered for this document. The included documents contain relevant information for the diagnosis of patients with SCA and SCD.

The terms used in the consensus statement are defined in Table 3 .

"Sudden unexplained death" refers to an unexpected and sudden death in an individual older than 1 year. Sudden death occurring unexpectedly within the first year of life is termed "sudden unexplained death in infancy" (SUDI). Multiple definitions have been in use over the past decades, although most recent studies implement a definition that differs between witnessed and unwitnessed events; in witnessed cases, death has to occur within 1 hour of change in cardiovascular status, whereas unwitnessed cases have to be seen alive and functioning normally within 24 hours of being found dead. 12 , 13

SCD constitutes the majority of SUD. 14 – 16 Reported overall SCD incidence rates vary across studies and countries, in part due to large difference in SCD definitions and methods for estimation of SCD rates. Previous studies report overall SCD rates ranging from 15 to 159 SCD per 100,000 persons per annum, corresponding to 6–20% of all deaths. 17 – 24 However, both incidence and causes of SCD vary markedly with age. Lowest SCD incidence is observed in children and adolescents. 15 , 18 , 25 – 28 SCD incidence is low in children and the young under 35 years and increases dramatically up until the age of approximately 60–80 years. 15 , 19 , 20 , 29 , 30

In young persons aged 1–35 years, most SCDs are caused by potentially inherited heart diseases, including primary arrhythmogenic disorders (eg, congenital long QT syndrome and catecholaminergic polymorphic ventricular tachycardia [CPVT]), hypertrophic cardiomyopathy, arrhythmogenic cardiomyopathy, and dilated cardiomyopathy; 15 , 18 , 25 – 29 , 31 , 32 however, coronary artery disease, anomalous coronary arteries, aortic dissection, congenital heart disease, and myocarditis are also potential causes, potentially with a non-negligible genetic component ( Figure 1 ). From the age of 35 years, coronary artery disease becomes the most common cause of SCD, although potentially inherited heart diseases remain a common cause of SCD at least until the age of 50 years. 27 , 29 , 33 Individuals with SUD who subsequently have negative pathological and toxicological assessment may be assumed to have sudden arrhythmic death (syndrome), or

SAD(S), a term synonymous with “autopsy-negative SUD.”

At any age, males have higher SCD rates compared with females, even after adjustment for risk factors of coronary heart disease. 34 Ethnic background seems to have large effect. 35 , 3 6

#### Recommendations for improving outcomes from sudden death

SUD is a tragedy and, in the case of an underlying genetic predisposition, may be preventable. The main cause of SUD is SCD. SCD in the young often occurs in people who were thought to be well, may occur without warning symptoms, and is often the first presentation of an underlying genetic heart disease. Across all ages, estimates differ from 5% to 20% of all deaths, and ethnicity-specific data on SCD incidences worldwide are sparse. Cause of death changes according to age ( Figure 1 ). Exact estimates of the burden of SCD are crucial in order to adjudicate public health spending.

1. and 2. Inherited cardiac disorders are the main cause of SCD in the young. Sudden death is SCD in 60–90% depending on age, of which the majority is potentially from inherited cardiac disease. 2 , 15 , 40 Exact estimates of the burden of SCD are crucial in order to adjudicate public health spending. 25

3. Estimates of SCD among different ethnic backgrounds are sparse worldwide. 35 , 36 , 38 , 39

Out-of-hospital cardiac arrest (OHCA) is a leading cause of mortality globally 41 – 43 and is defined as the loss of functional cardiac mechanical activity in association with an absence of systemic circulation, occurring outside of a hospital setting. The exact burden of OHCA remains unknown, since a considerable number of cases are not attended by emergency medical services (EMS) and regional variations are prevalent in both reporting systems and survival. 17 , 44 , 45 Approximately 275,000 people in Europe have cardiac arrest treated by EMS per year, with only 29,000 (10.5%) surviving hospital discharge. 46 In England, 28,729 EMS-treated OHCA cases were reported in 2014 (53 cases per 100,000 of the resident population), with only 7.9% surviving to hospital discharge. 47 In the United States, reports from 35 communities suggested an incidence of 55 per 100,000 person-years 48 or approximately 155,000 individuals having an EMS-treated all-rhythm OHCA per year. 48 Globally, the weighted incidence estimates per 100,000 person-years of EMS-treated OHCA are 34.4 in Europe, 53.1 in North America, 59.4 in Asia, and 49.7 in Australia. For reported survival estimates, the percentage survival to discharge was 7.6% in Europe, 6.8% in North America, 3.0% in Asia, and 9.7% in Australia. 42

Significant geographical variation in the incidence of OHCA associated with poor outcomes has remained unchanged in the past 3 decades. 17 , 41 , 42 , 44 However, implementation of coordinated efforts targeted at improving the local chain of survival in some cities has improved regional survival to 20–40%. 49 , 50 This survival benefit can be partially attributed to varying definitions of OHCA, 42 but it is primarily due to a coordinated effort to optimize the effectiveness of the local chain of survival. 51 Identifying and improving weak links in the local chain of survival, paired with targeted approaches to improve the effectiveness, has resulted in positive outcomes achieved in several geographic regions. 49 , 52 – 54

OHCA causes are classified into cardiac and noncardiac causes. 47 , 55 , 56 Approximately 80% of individuals presenting with OHCA reached by EMS, and in whom resuscitation is considered possible, have a cardiac cause. 56

OHCA can affect seemingly fit and healthy athletes, young adults, or children. The incidence of SCD in athletes can range from 1 in 23,000 to 1 in 200,000 athletes per year, depending on a number of factors including populations studied. 57 , 58 In a retrospective analysis of the Rescu Epistery database of consecutive OHCA attended by EMS in a specific area of Ontario, Canada, the incidence of SCD during participation in competitive sports was reported to be 0.76 cases per 100,000 athlete-years. 59 The main causes of SCD were stratified by age. In those younger than 35

years, structural heart and primary arrhythmic causes were most common. In those aged between 35 and 45 years, coronary artery disease was the most frequent underlying pathology. 59 In a prospective study of children and young adults aged 1–35 years, 490 cases of SCD were identified from centers in Australia and New Zealand. 25 The cause of death was unexplained in 40% of these cases at autopsy, in whom a structurally normal heart was reported. 25 In this study, the annual incidence of SCD was calculated to be 1.3 cases per 100,000 people. When stratified according to age group, the highest incidence (3.2 cases per 100,000 people per year) was observed in those aged 31–35 years. Coronary artery disease was the most common cause ascribed. Younger age and SCD occurring at night were independently associated with unexplained SCD, probably due to congenital channelopathies. Less common causes were inherited cardiomyopathies (eg, dilated, hypertrophic, and arrhythmogenic right ventricular), myocarditis, and aortic dissection.

The Cardiac Arrest Registry to Enhance Survival (CARES), established by the Centers for Disease Control and Prevention (CDC), 60 evaluated OHCA events of presumed cardiac etiology that involve persons who received resuscitative effort. OHCA is defined in CARES as a cardiac arrest that occurred in the prehospital setting, had a presumed cardiac etiology, and involved a person who received resuscitative efforts, including cardiopulmonary resuscitation (CPR) or defibrillation. The registry includes 40,274 OHCA records, of which 31,689 OHCA events were presumed to be of cardiac etiology (eg, myocardial infarction or arrhythmia) that received resuscitation efforts in the prehospital setting (mean age 64.0 years [SD 18.2]; 61.1% male). The survival rate to hospital admission was 26.3%, and the overall survival rate from cardiac arrest to hospital discharge was 9.6% ( Figure 2 ). Approximately 36.7% of OHCA events were witnessed by a bystander. Only 33.3% of all patients received bystander CPR, and only 3.7% were treated by bystanders with an automated external defibrillator (AED) before the arrival of EMS providers.

The group most likely to survive an OHCA is persons who are witnessed to collapse by a bystander and found in a shockable rhythm (ie, arrhythmias leading to ventricular fibrillation or pulseless ventricular tachycardia). 60 Among this group, survival to discharge was 30.1% ( Figure 3 ). A subgroup analysis, performed among persons who experienced OHCA events unwitnessed by EMS, revealed that whites were significantly more likely to receive CPR than blacks, Hispanics, or members of other racial/ethnic populations (  $p < 0.001$  ). Overall survival to hospital discharge of patients whose events were not witnessed by EMS personnel was 8.5%. Of these, patients who received bystander CPR had a significantly higher rate of overall survival (11.2%) than those who did not (7.0%) (  $p < 0.001$  ).

Figure 4 shows bystander CPR and lay AED use by percentage of black residents in the area. Directing attention toward improving education, availability of AEDs, and treatment of cardiac arrest in predominantly black neighborhoods may save lives. 39

Bystander AED use in OHCA in pediatric populations is variable and uncommon, with important variations based on neighborhood characteristics leading to marked disparities in survival and outcomes. Griffis et al. 38 reported that AED use (likely due to availability) was more common in neighborhoods with a median household income of >\$50,000 per year (12.3%;  $p = 0.016$  ), <10% unemployment (12.1%;  $p = 0.002$  ), and >80% high school education (11.8%;  $p = 0.002$  ). Greater survival to hospital discharge and neurologically favorable survival were among arrests with bystander AED use, varying by neighborhood characteristics.

The majority of persons who experience an OHCA event, irrespective of etiology, do not receive bystander CPR or other timely interventions that are known to improve the likelihood of survival to hospital discharge (eg, defibrillation). 54 Because nearly half of cardiac arrest events are witnessed, efforts to increase survival rates should focus on timely and effective delivery of interventions by bystanders and EMS personnel ( Figure 5 ).

Education of public officials and community members regarding the importance of increasing rates of bystander CPR and promoting the use of early defibrillation by lay and professional rescuers is

critical to increasing survival rates. Reporting at local and national levels can enable local and national public health and EMS agencies to coordinate their efforts to target improving emergency response for OHCA events, regardless of etiology, which can lead to improvement in OHCA survival rates ( Figure 6 ).

#### Recommendations for improving outcomes in SCA survivors

OHCA remains a significant cause of mortality globally. Despite implementation of cardiac arrest protocols including CPR training and AEDs, only 33% of witnessed OHCA cases receive bystander CPR and less than 4% are defibrillated onsite. OHCA hospital discharge survival remains dismal at around 10% and has remained stagnant for the past 3 decades. Significant geographic variation in OHCA incidence and the role of social disparities merit further research. Public health campaigns promoting CPR training in at-risk communities and greater availability of AEDs are needed.

Coordinated efforts targeted at improving the local chain of survival have improved regional survival. 49 , 50 Targeted approaches to improve the effectiveness of CPR have resulted in positive outcomes. 52 – 54 The group most likely to survive an OHCA is persons who are witnessed to collapse by a bystander and found in a shockable rhythm, so widespread CPR training is recommended. Subgroup analysis has revealed that whites were significantly more likely to receive CPR than other racial/ethnic populations. 60 AED use was more common in neighborhoods with high median household income, <10% unemployment, and >80% high school education. 38 Therefore, maximum benefit will be gained from targeting CPR training to groups of high socioeconomic need and ethnic minorities.

The burden of OHCA and the response of bystanders appears to vary according to ethnicity and socioeconomic status. 35 , 38 , 39 , 60 Further investigation of these findings may result in targeted approaches to maximize outcome from investment when aimed at these communities.

Availability of AEDs has been shown to improve survival. 61 – 63 Therefore, as the majority of cardiac arrests are witnessed, AEDs at schools, stadiums, stations, etc, may be expected to increase survival. Venues where delivery of AEDs by emergency services is unlikely (eg, trains, ships, planes) are of particular importance. Appropriately maintained equipment and appropriate training of potential AED users are an essential component of this strategy.

The investigation of SCD and resuscitated SCA requires input from a variety of different disciplines. The coordination and the communication between them mandate the formation of a multidisciplinary team. Numerous consensus statements agree on the importance of a dedicated combined cardiac genetic service in this setting. 2 , 11 , 65 – 67

Certain key features can be identified in well-functioning multidisciplinary teams across specialties. Nancarrow et al. 68 propose 10 key attributes including positive leadership and management, communication strategies and structures, appropriate resources, appropriate skill mix, and a supportive team climate with a focus on education of each other. There should be open communication and shared decision-making.

The detection of inherited heart conditions by pathologists and by hospital clinicians requires heightened awareness of their existence and a simple referral pathway to a multidisciplinary service with cardiac genetic expertise. Clinical experience shows that the appointment of a coordinator, as well as an enthusiastic team leader, is essential to facilitate this process, and regular meetings increase relevance and improve attendance 69 ( Figures 7 and 8 ).

The investigation of SUD is led by (forensic) pathology and the investigation of resuscitated SCA by pediatric or adult cardiology, with cardiac heart rhythm specialists and genetic cardiologists often being central. Clinical and molecular genetic specialists and genetic counselors are needed because of the significant role of molecular genetics in achieving a diagnosis and cascade

screening, the consideration of multisystem genetic syndromes, and the high prevalence of genetic variants of uncertain significance. 70 , 71 The high levels of psychological morbidity among SCA survivors and family members of both SCA survivors and decedents mandates access to psychology expertise, 72 – 74 and input into the multidisciplinary team helps keep this in focus. 75 The presence of a specialist cardiac genetic nurse in the cardiology inpatient setting increases detection of inherited cardiac conditions following SCA. 76 Other clinical specialists can be helpful and be drafted in for certain cases—neurologists, pediatricians, metabolic specialists, and intensivists, for example.

The fact that the many disciplines may not be co-located highlights the importance of a coordinator to a multidisciplinary service. Co-location is not critical for effective collaboration, and non-co-location should not be an excuse for failed collaboration. A regional or institutional coordinator could be a nurse specialist, genetic counselor, or other allied professional and is vital to facilitate team meetings and communication between specialists and between centers, with primary care and across regions or between states and countries where necessary to facilitate family screening 69 , 77 ( Figure 7 ).

Links to other services as proposed in a recent scientific statement 66 and practiced by some centers already 69 include connections to molecular genetic expertise, researchers, primary health providers, between regions, and to a cardiac genetic clinical registry to facilitate family screening and follow-up across traditional boundaries ( Figure 8 ).

Clinical and genetic registries are generally voluntary and consent-based and have a research element. We do not consider that they are compulsory. However, in this setting they do have particular relevance because many cases remain unresolved after the initial investigation and families may find comfort in knowing that efforts to find a diagnosis continue. The multidisciplinary team also provides a mechanism to revisit family members if new findings appear in the wider family or if the pathogenicity of a genetic variant is redefined.

Recommendations for the role of a multidisciplinary team for investigation of SUD and SCA

The cardiac and genetic investigation of SUD and resuscitated SCA should be overseen by a multidisciplinary team with appropriate expertise in this area. Recommendations include adequate resourcing, a dedicated coordinator, strong leadership, and a mutually supportive team that meets regularly.

For regions where coordinated cardiac genetic services exist that include the investigation of SUD, detection of inherited heart conditions is higher than in regions where they are not. 78 Families prefer specialized clinics that combine co-located cardiac and genetic expertise and genetic counseling. 79 Many such dedicated clinics internationally have led to the detection of inherited heart conditions following SCD and resuscitated SCA. 80 – 82 It is therefore logical that multidisciplinary teams should have links to such clinics. Continued productive dialogue among pathology, coronial, police, and cardiac genetic services is recommended to improve the quality and relevance of forensic pathologists' reports. 86

Genetic testing in this context leads to a significant proportion of both pathogenic and unclassified variants, and precise evaluation of clinical phenotype is imperative for the correct assignment of such variants, so that a service that combines specialist cardiology and genetic expertise is essential. 83 , 84 Specialized clinics that combine co-located cardiac and genetic expertise and genetic counseling are preferred, 79 and such combined clinics have a high detection of inherited cardiac conditions following resuscitated SCA. 82 , 85

Genetic counseling is a process that aims to assist patients and their families to understand and adapt to the medical, psychosocial, and familial impact of inherited diseases. 87 , 88 Genetic counseling goes beyond the discussion of genetic testing and is important for all patients with a

genetic condition, at all stages of management. 89 Although genetic counseling may be performed by any number of health professionals, genetic counselors are specifically trained in this role and have grown to a large allied health workforce worldwide. 90 , 91 In some institutions, this role may be performed by a clinical/medical geneticist, genetic nurse, or other appropriately trained specialist.

In the setting of SCD or resuscitated SCA where a genetic cause is suspected, the inclusion of genetic counselors in the multidisciplinary team is widely advocated. The role of the cardiac genetic counselor includes taking a detailed family history, investigating and confirming details such as postmortem reports, providing education and awareness, assisting in coordinating family clinical screening, and providing psychosocial support. 74 , 92 – 95 Throughout the process of genetic testing, genetic counselors provide important pre- and post-test genetic counseling, assist with interpretation of the results, help communicate this information to relatives, and assist with cascade genetic testing 75 , 93 , 96 , 97 ( Table 4 ).

Where there are significant emotional difficulties (see Section 5 ), the process of effectively conveying genetic information can be challenging. 98 For families who have experienced a young SCD where a genetic cause is suspected, learning the potential inheritance risk to family members and need for clinical screening can add an additional stressor at a time of intense grief. Furthermore, with the increasing availability of postmortem genetic testing (see Sections 6.4 and 6.5 ), the need for complex genetic discussions with families is more commonplace. 99 Genetic counseling prior to and after genetic testing is important, particularly where genetic test results are not straightforward such as identification of variants of uncertain significance or in the event of a variant reclassification. 100 , 101

There is wide acknowledgment that genetic counseling as a process should go beyond just provision of information. 10 2 , 103 The psychosocial aspects of genetic counseling include psychological support, empathic listening, crisis intervention skills, knowledge of family dynamics, coping models, processes of grief, and adjustment to disease diagnoses, all of which align with the core competencies of genetic counseling accreditation. 10 4 , 105 Attending to the psychosocial needs, in addition to provision of education and information, has been demonstrated to positively impact patient outcomes, largely based around knowledge and recall, but healthy adjustment, empowerment, behavioral change, and satisfaction with decision-making also reduce anxiety and worry. 103 , 104 , 106 , 10 7

#### Recommendations for counseling families affected by SUD and SCA

Genetic counseling of patients and their families with genetic conditions is recommended, including those with SCD or resuscitated SCA where a genetic cause is suspected. Key aspects of the process include discussion of inheritance risks, education and awareness, pre- and post-test genetic counseling, interpretation of genetic results, taking a family history, coordination of clinical screening, and psychosocial support. Genetic counseling is focused on both information provision and psychosocial support and together has been shown to improve knowledge and recall; promote healthy adjustment, empowerment, and behavioral change; increase satisfaction with decision-making; and reduce anxiety and worry. While genetic counseling is a process often performed by a variety of health professionals, ideally a specifically trained genetic counselor or genetic nurse with appropriate skills in information provision and psychosocial support would perform this role.

Genetic counseling includes both information provision and psychosocial support. It is ideally performed by health professionals with specific training and experience; this includes genetic counselors, genetic nurses, or other qualified health professionals. 80 – 82 , 84 , 85 , 103 , 106 , 107 , 109

In the context of genetic testing, pre- and post-test genetic counseling must be performed. 75 , 95 , 96 , 99 In cases where there is uncertainty in the findings, such as a variant of uncertain significance or a variant reclassification, this is of particular importance. 100 , 101

SCD where a genetic cause is suspected has a profound psychological impact on the surviving members of the family. Grief is a normal emotional response to the loss of a loved one. Individuals will grieve differently, and while there is no single trajectory, many will experience disbelief, yearning, anger, sadness, and acceptance. 117 After a death, an individual will not return to normal, but rather create a revised meaningful life without the deceased. In a small proportion of bereaved individuals, the initial grief response does not resolve and may result in prolonged grief, or persistent complex bereavement disorder according to the Diagnostic and Statistical Manual of Mental Disorders, 5th Edition (DSM-5). 118 , 119 This occurs in approximately 7% of the general bereaved population, 120 and in 21% of first-degree relatives following SCD in the young. 121 Posttraumatic stress symptoms can also be experienced by family members. Posttraumatic stress occurs in response to a specific trigger, typically one that threatens one's own or a loved one's well-being. It is characterized by avoidance with hyperarousal and intrusive thoughts, including persistent and extreme fear and panic similar to that experienced by family members at the time of the event. 122 Posttraumatic stress has been shown in 44% of first-degree relatives following a young SCD. 121 Individuals with prolonged grief and/or posttraumatic stress symptoms can benefit from intervention with a clinical psychologist or other appropriately trained clinicians, and there is extensive evidence to support the efficacy of psychological treatments for these conditions in other settings. 120 Further, there is greater risk of other psychiatric comorbidities, 118 suicide, 123 and development of chronic medical conditions. 124

Factors associated with poor psychological outcomes have been investigated. One study showed that mothers of the deceased were more likely to report anxiety and depression symptoms. 72 In total, 53% of the mothers surveyed reported probable anxiety disorder on average 4 years after the death. In a larger study, after adjusting for factors including relationship to the decedent, those family members who witnessed the death or discovered the decedent's body had a 3-fold risk of posttraumatic stress symptoms (OR 3.3, 95% CI 1.2–8.7,  $p = 0.02$ ) and a 4-fold risk of prolonged grief (OR 4.0, 95% CI 1.3–12.5,  $p = 0.02$ ). 121 Given that half report symptoms indicating psychological difficulties, all first-degree relatives should be offered psychological evaluation and treatment. Although the evidence for psychological support is derived from studies investigating SCD where a genetic cause is suspected, it may logically apply to those individuals who have survived SCA and their families ( Figure 9 ).

There may be initial reluctance to seek psychological support given community stigma around mental health. Indeed, a recent study investigating families who had experienced a young SCD found that only 12% had sought psychological support, with most of those being self-referrals. 125 In discussing options for ongoing psychological support with patients and families, normalizing their response to a significant psychological stressor and describing common symptoms of prolonged grief and posttraumatic stress may reduce any perceived sense of stigma and increase interest in seeking support.

A recent needs analysis of parents who had experienced the SCD of their child (including adult children) found that while medical information and support were the most important need, psychological information and support were the most unmet need. 126 Nearly three-quarters reported wanting access to professional counseling or psychological services. Further, many indicated access to genetic testing or understanding the genetic cause to be an important need, highlighting the importance of maintaining realistic expectations regarding the diagnostic yield of postmortem genetic testing with families. 101 At present, this is likely not greater than 15%, and there is a high likelihood of uncertain genetic findings especially with increasing gene panel sizes. 25 , 70 A Swedish study of parents whose children died suddenly between 15 and 35 years of age likewise showed a critical lack of information and support in the acute grief stage. 127 This included a need for better communication of the postmortem examination process (how long it would take, when they would get results), time with a health professional to discuss the death, and information

about the cause of death. There was a lack of psychological support in the immediate aftermath, with many family members seeking their own care, including grief counselors and support groups. The need for support in the early aftermath has been shown to be important in other studies examining suddenly bereaved parents. 128 , 129

Community or peer-based bereavement support groups can also enhance social support. 120 , 130 Peer support programs come in many different forms but always involve people with similar backgrounds providing emotional, social, or practical support to each other. 131 Peer supporters draw on their shared experiences to provide empathic understanding, information, and advice to those they are helping. A key aim is to promote hope, recovery from illness or trauma, improved life skills, psychological well-being, and social integration. 132 A recent systematic review of peer support services for bereaved survivors of the sudden death of a loved one in multiple settings found evidence of reductions in grief and increased well-being and personal growth among participants, and improved personal growth and positive meaning in life among peer providers. 133 There is a current gap in care in addressing psychological support needs of families after the SCD of a young relative.

The psychological impact to the family following an SCD where a genetic cause is suspected can be significant. Although many family members will navigate their way through this traumatic experience, up to 44% may require additional psychological support from an appropriately trained health professional such as a clinical psychologist. Addressing community stigma around mental health needs to be considered and discussed with families. In addition, support services such as social workers, grief counselors, psychosocial teams, and peer support groups may be useful to many families. Whereas the evidence for psychological support is derived from studies investigating SCD where a genetic cause is suspected, it may logically apply in those families where there has been an SCA.

1. and 2. A clinical psychologist or appropriately trained health professional includes those equipped to assess and treat trauma; for example, those experienced in delivering cognitive behavioral therapies. While the evidence to date supports a need for psychological support in family members following a young SCD where a genetic cause is suspected, this may likewise be important for relatives of a patient who suffers an SCA. 72 , 121 , 134 There is a need to train personnel in psychological care for SUD and SCA, as this is an area where the need is not currently met.

Despite being “low-tech” and inexpensive, the history, as a tool for clinical phenotyping, is the essential and fundamental basis of approaching a patient with SCA because it can guide appropriate use and interpretation of other diagnostic modalities. The history should be focused toward both the decedent proband and also the wider family for evidence of other potentially affected members prior to investigations. Surviving family members should be investigated by a multidisciplinary team within a specialist program for cardiovascular genetic disorders with the all appropriate medical, genetic, and psychological personnel and ability for comprehensive investigations 2 , 10 ( Figure 10 ).

The proband age may help define potential etiologies; CPVT and long QT syndrome are typically diseases of the young, whereas coronary artery disease and cardiomyopathies become more common with age ( Figure 1 ). Although most deaths occur at rest or during sleep, 25 death during exertion may point to specific etiologies such as CPVT, long QT syndrome type 1, or arrhythmogenic cardiomyopathy. In addition to a detailed prior medical and medication history (including potential drugs of abuse), the decedent's health in the 24–48 hours preceding death including the presence of any viral prodrome or fever, as well as any prescribed medication, may be relevant. Myocarditis secondary to viral infection may be associated with viral and gastrointestinal symptoms, and both Brugada syndrome and long QT syndrome may be exacerbated by specific pharmacological agents through further inhibition of ion channel function. 135 Fever is a well-recognized trigger of ECG changes and arrhythmia in Brugada syndrome 136 and in some long QT syndrome subtypes, 137 and in young children may be misdiagnosed as febrile seizures.

Between 18% and 45% of sudden death cases may have experienced prior relevant symptoms, typically palpitations, chest pain, pre-syncope, or syncope, and may have undergone relevant investigations. 29 , 70 , 139 All medical records relevant to the sudden death etiology should be sought.

Relevant information from the family history should be collected by a health professional with specific experience in cardiovascular genetic disease (preferably a genetic counselor) and by an appropriately trained cardiologist. Symptoms and diagnoses in other family members as well as prior cardiovascular investigations should be sought. Noncardiac findings may be highly pertinent including unexplained epilepsy unresponsive to conventional therapy; skeletal muscle weakness; curled hair and subtle palmoplantar hyperkeratosis/keratoderma (arrhythmogenic cardiomyopathy) 140 ; attention deficit disorder and intellectual disability (CPVT) 141 ; and history of pneumothoraces, vascular disease, and gastrointestinal and uterine rupture (vascular Ehlers-Danlos syndrome). 142 Any other deaths or major cardiac events in the family should be recorded including those related to drowning in good swimmers, unexplained motor vehicle accidents, and sudden infant death or late fetal demise. If the SUD was observed, it is useful to collect witness accounts about the events occurring immediately prior to the collapse and during any resuscitation attempts.

#### Recommendations for investigation of sudden death: personal and family history

The personal medical and three-generation family history provides the initial information on which subsequent investigations will be based. Specific features within the wider family may suggest diagnoses and help direct subsequent investigation. The history should be recorded by cardiologists, specialist nurses, and geneticists or genetic counselors experienced in cardiovascular genetic diseases, ideally within the confines of a multidisciplinary program that can address the medical, genetic, and psychological needs of the family (see Section 3 ).

The personal and three-generation family history may provide critical information relevant to the etiology of SCD and provide a starting point for further investigations in both the decedent proband and surviving family members. 77 Multiple studies show a significant proportion of children and adults experience relevant cardiac symptoms prior to sudden death, 10 , 25 , 29 , 70 , 135 , 138 , 139 and some may have sought medical attention and undergone investigations. Available ECGs and cardiac imaging, together with autopsy findings, may allow a diagnosis to be made (or excluded) in the proband who, until family investigations have been performed, is the one definitively affected member of the family. Noncardiac features and symptoms may also provide important diagnostic information. 140 – 142

Further investigation is necessary where sudden death occurs in specific circumstances such as when a cardiac event may have triggered an apparently environmental death. Examples include road traffic accidents with no apparent cause and drowning in competent swimmers.

The presence and associated investigations for other noncardiac conditions should also be evaluated, specifically epilepsy. Failure to identify a neurological etiology or abnormality would suggest seizures may have had a cardiac etiology. Overlap syndromes exist between true neurological epilepsy and long QT syndrome type 2. 146 , 147

Individuals who have succumbed to SUD may have had pertinent investigations prior to their death that aid in the diagnosis of the cause of their SUD. Twelve-lead electrocardiogram (ECG) is the most useful pre-SUD investigation. Although long or short QT interval, spontaneous type 1 Brugada pattern, and early repolarization pattern are associated with fatal arrhythmias due to congenital long QT syndrome, short QT syndrome, Brugada syndrome, or early repolarization syndrome, 148 – 152 many patients with SUD without structural heart disease have a normal or near-normal ECG, 145 ,

153 particularly women. 154 Additional ECG findings suggestive of arrhythmic syncope include bifascicular block; intraventricular conduction abnormalities (QRS duration >0.12 s); Mobitz I second-degree atrioventricular block and first-degree atrioventricular block with markedly prolonged PR interval; sinus bradycardia (<40 bpm) or slow atrial fibrillation (<40 bpm); nonsustained ventricular tachycardia; pre-excited QRS complexes; negative T waves in right precordial leads or epsilon waves; and left ventricular hypertrophy, 5 any of which may indicate potential diagnoses of inherited arrhythmia syndromes such as progressive cardiac conduction defect, familial pre-excitation, arrhythmogenic cardiomyopathy, or hypertrophic cardiomyopathy. 4 , 151 In the general population, premature ventricular complexes (PVCs) are mostly benign; however, some frequent or complex PVCs significantly increase the risk of SCD. 155 – 157 If an ECG is recorded by the AED or EMS just before SCD, features such as J-wave or ST segment elevation (especially if augmented after a long pause) may help in the diagnosis of coronary spasm, early repolarization syndrome, or Brugada syndrome. 158 , 159 Interpretation of ECGs obtained immediately after resuscitation/defibrillation should be performed with great caution (see Section 7.4 ).

Syncope is a sentinel clinical symptom before SUD and may prompt investigations subsequently useful in making a retrospective diagnosis of the cause of SUD. In particular, the trigger for the syncopal event bears useful information. Ambulatory ECG monitoring during life may provide clues to the cause of SUD and should be sought.

If transthoracic echocardiography, cardiac computed tomography (CT), or cardiac magnetic resonance imaging (CMR) are performed during the patient's life, detailed review may indicate features of dilated cardiomyopathy, hypertrophic cardiomyopathy, or arrhythmogenic cardiomyopathy. 160 If blood or other tissue sample has been taken before SUD, this may be a source of DNA for genetic testing, should there not be a postmortem collection of tissue. 8 , 11 , 67 , 15 1 Neurological findings such as developmental delay or seizures thought to be suspicious for epilepsy during life may contribute to a diagnosis of a cardiac channelopathy, such as CPVT or long QT syndrome. 141 , 161 , 16 2 If a patient with SUD has a cardiovascular implantable electronic device (CIED) implanted, postmortem interrogation of the CIED is useful to determine the cause and timing of SCD. 163

Recommendations for investigation of sudden death: examination of premorbid investigations

During the investigation of SUD, pertinent investigations performed prior to death can aid in establishing the cause. Although ECG features such as QT interval, type 1 Brugada pattern, and early repolarization may be critical for diagnosis, many ECGs taken during life will be normal. Ambulatory ECG monitoring and cardiac imaging should be sought to provide clues to the diagnosis of SCD. Symptoms attributed to a neurological cause may be re-evaluated, in collaboration with neurologists. Any potential DNA sample before SUD should be stored if tissue is not gained at autopsy. ECG information from the AED, emergency services, or CIEDs may also be useful to determine the cause of SCD.

Investigations during life may provide clues to the cause of SUD and should be sought to aid in diagnosis. 173 These include resting, exercise and ambulatory ECG tracings, and cardiac imaging studies (echocardiography, CT, and CMR). 145 , 148 – 150 , 153 – 158 , 164

Samples for potential DNA testing taken during life may subsequently prove invaluable should they be the only source of DNA. Although it is a Class 1 recommendation that patients with SUD have an autopsy and material for DNA testing collected (see Section 6.3 ), it is recognized that sometimes this is not done, making blood or tissue samples taken during life the only remaining source for molecular autopsy. 84 , 165 – 167 Using such samples requires appropriate consent from family, unless ordered by the coroner. It should be recognized that success varies depending on the storage method, but attempts to gather useful DNA may be worthwhile even from suboptimal sources. Future extraction methods may improve the yield so continued storage is advisable.

Symptoms such as seizures thought to be suspicious for epilepsy during life may in fact be attributable to a cardiac channelopathy when further investigation is done. Other neurological findings such as developmental delay hold significance for diagnoses such as CPVT. Thus, meticulous recording of neurological events during life may lead to a diagnosis in SUD. 70 , 146 , 147 , 161 , 162 , 168 , 169

Recordings from emergency services continuous ECG monitoring or from interrogation of AEDs, when available, may provide clues to the etiology of SUD. However, it should be acknowledged that a finding of ventricular fibrillation is often due to this rhythm being a final common rhythm in arrhythmic death, regardless of the initial rhythm causing hemodynamic collapse. Nevertheless, at points such as reinitiation of arrhythmia and glimpses of normal rhythm in between arrhythmia may suggest a specific diagnosis. 158 , 170

The memory function of CIEDs may reveal the initiation pattern of cardiac arrhythmia and aid in the diagnosis of SCD. Therefore, if an SUD victim has a CIED implanted in life, interrogation of this device can provide useful clues in the diagnosis of SCD. 163 , 171 , 172

The critical components to the investigation of SUD include examining the circumstances of the death and the autopsy ( Figure 11 ). Identification of SUD relies on the reporting of EMS, police, hospitals, and witnesses. Investigation of a death is determined by the jurisdiction in which the death occurs. 14 , 174 , 175 Unexpected or unexplained deaths, when the individual was in apparent good health, should be carried out by a trained pathologist who has a thorough knowledge of cardiac pathology. 176 Autopsies vary not only by country but also by individual jurisdictions within countries. The autopsy should be comprehensive, examining all organs and conducted in a systematic and objective method with a focus on standardized reporting. 2 , 67 , 177 , 178 Cases should be referred to a cardiac pathologist when a cardiac cause is suspected. 2 , 176

Imaging includes X-rays and photography. Photography is useful in providing documentation of syndromic features and highlighting individual organ pathology. Postmortem CT and magnetic resonance imaging (MRI) have been shown to be useful 179 but are not universally available. Noncardiac causes should be looked for including infection, thromboembolism, tumors, intracerebral lesions, respiratory disease, and abdominal causes such as ruptured abdominal aneurysm. Body mass index should be recorded along with waist circumference.

Ancillary testing should be performed including microbiology/cultures for infectious disease, metabolic screening (particularly in younger children), toxicology, vitreous testing for biochemistry, genetic testing (see Sections 6.4 and 6.5 ), and other testing as indicated by the autopsy findings. Taking a sample for toxicology is recommended in all sudden unexpected deaths. 180

Samples for genetic testing should be saved at the time of autopsy from every sudden death case. 2 , 181 Ideally, two of the following three should be saved: a small piece of fresh frozen heart, a small piece of fresh frozen spleen/liver/thymus, and EDTA blood. If RNAlater (ThermoFisher Scientific, Waltham, MA, USA) or similar reagent to preserve DNA at room temperature is available, fresh tissue can be transported in this to the referral genetic center without need for freezing.

The heart should be examined thoroughly 67 and at least 7–10 samples taken for histology. Cardiovascular disease is the leading cause of sudden death in the young and is divided into two major groups: morphologically positive (eg, congenital heart disease, coronary artery disease, and cardiomyopathy) and morphologically normal hearts. Combined with negative toxicology, those with morphologically normal hearts have been labeled as having “autopsy-negative sudden unexplained death” or “sudden arrhythmic death (syndrome) or SAD(S).” 182 Samples should always be taken, even from a macroscopically normal heart, as histology may reveal inflammation and cardiomyopathies. Always consider sudden unexpected death in epilepsy (SUDEP) and sudden death in alcohol misuse (SUDAM) 183 where clinical history and circumstances are important. Pathologists and clinicians should not overinterpret findings in the heart at autopsy such as

nonsignificant coronary artery disease, etc. 184

Recommendations for investigation of sudden death: the postmortem examination and imaging

A comprehensive autopsy is an essential part of the investigation of SUD and should include collection and storage of tissue suitable for genetic analysis. When the autopsy suggests a possible genetic cause, or no cause and the heart is normal, referral to a multidisciplinary team for further investigation is indicated (see Section 3 ).

SUD should have an autopsy done by a trained pathologist. 174 – 176 Studies have shown that autopsies performed by pathologists who have a thorough knowledge of cardiac pathology have a superior diagnostic yield. 174 – 176 In cases where autopsy is not possible (eg, for religious reasons), a full body MRI or CT scan is recommended. 179

Samples should be taken for infection and toxicology. Histological sampling of all important organs especially the heart (from multiple sites) is essential even when macroscopically normal. 174 , 175

Blood or tissue suitable for DNA extraction and postmortem genetic testing should be obtained at all autopsies. 176 Following the initial investigation, DNA should be extracted and banked if genetic disease is suspected or if the cause remains unknown. 2 , 176 Ideally, lack of cost coverage should not be a reason not to comply with these recommendations.

Frozen myocardial tissue may be useful for subsequent RNA analysis or expression studies of aberrant proteins. 2 , 67

Cause of death should be discussed in a multidisciplinary meeting (see Section 3 ) and provided by a pathologist to the medical examiner/coroner. The findings and follow-up recommendations should be communicated to the family. 178 , 190 , 191

Finding of an abnormal or normal heart is important for family screening and directs much of the subsequent investigation (see Sections 6.4 and 6.5 ).

Autopsy phenotype should be established at a multidisciplinary meeting of pathologist, medical examiner, cardiologist, and clinical geneticist.

Genetic evaluation may be appropriate following SCD in two scenarios: most commonly, where the deceased individual is the proband with no prior medical history, or alternatively the deceased is part of a family where diagnosis is established but he/she has not yet undergone genetic evaluation. Initially, pathological examination should be performed by an experienced pathologist to ensure that all cardiac and extracardiac features relevant to the potential diagnosis are recognized (see Section 6.3 ).

In cases where the deceased is the proband and a postmortem diagnosis is established, identification of a pathological variant may facilitate genetic testing in the wider family evaluation. Genetic testing of DNA from the deceased proband may be performed directly after autopsy or deferred until first-degree family members have been clinically evaluated ( Figure 12 ). As part of familial evaluation, a three-generation pedigree (at a minimum) performed by a practitioner knowledgeable in the genetics of cardiovascular disease (eg, a genetic counselor or specialist nurse) is mandatory and should cover all potentially relevant cardiac and extracardiac features within the family (see Section 6.1 ). Genetic testing of deceased individuals may not be covered by health insurance in certain countries; in this instance, using a clinically affected family member as the testing proband with confirmatory testing in the deceased may be a more feasible strategy. Clinical and genetic testing in the proband and multiple family members will define segregation of the identified genetic variant(s), adding to the validity of the genetic findings. The yield of genetic testing in cases where a diagnosis of cardiomyopathy is established postmortem is significantly

greater than where structural changes are uncertain. 192

In cases where no other family members are clinically affected and the deceased proband is an apparently isolated case, genetic testing can be used as evaluation of the single definitely affected individual within the family. Families should be counseled about the expected benefits and potential outcomes of genetic investigations prior to testing. If identified variants are considered likely pathogenic or pathogenic, cascade testing across the family can be considered to identify at-risk individuals with no current clinical features.

In cases where the deceased is part of a family with a prior diagnosis of cardiovascular disease and known pathological variant, confirmatory genetic testing may be performed.

Recommendations for investigation of sudden death: genetic evaluation where the phenotype is known

A postmortem diagnosis following SCD significantly facilitates further clinical evaluation of family members and may provide an explanation for the family as to the underlying etiology. Genetic testing targeted toward the clinical diagnosis and phenotype is an important component of the overall evaluation of both the proband and family and provides additional support to the clinical diagnosis. Further investigations can be performed as clinically indicated. In cases where no other individuals within the family are clinically affected, identification of a definitively pathogenic variant in the proband facilitates cascade genetic testing in family members.

Genetic testing in the decedent proband is recommended to support the clinical diagnosis and facilitate cascade genetic testing within the family. 10 , 11 The yield of genetic testing is significantly higher when associated with a specific postmortem diagnosis. 84 , 192

If a cardiac phenotype has been identified in the deceased proband, genetic testing should be targeted toward that specific phenotype to maximize the chances of a clinically actionable result and minimize the risk of ambiguous secondary findings. Targeted panels for cardiomyopathy, channelopathies, familial thoracic aortic aneurysm, and familial hyperlipidemia are the preferred option. However, given the limited availability of proband DNA and potential financial implications, broader genetic testing including whole exome and genome with selective reporting based on the phenotype may be considered. Robust evidence for gene–disease association is currently probably only available in the Clinical Genome Resource (ClinGen) expert panels for long QT syndrome, Brugada syndrome, hypertrophic cardiomyopathy, arrhythmogenic cardiomyopathy. 195 , 196 , 202 Wider panels and whole-exome sequencing/whole-genome sequencing share the potential disadvantage of a high burden of variants of uncertain/unknown significance, but in experienced hands (ie, expert centers), this will not lead to unintended follow-up. 203

Clinical evaluation for family members of a proband with an evident or likely phenotype can be targeted based on that phenotype, although due to varied expressivity and overlap syndromes should be sufficiently broad to provide a comprehensive cardiovascular evaluation. If a pathogenic or likely pathogenic variant has been identified in the deceased proband in a gene consistent with the clinical diagnosis, genetic counseling and testing should be offered to family members. 25 , 80 – 82 , 84 , 85 , 143 , 189 , 193 Cascade testing for likely pathogenic variants should be done at the discretion of an experienced provider after reviewing the data. In cases with limited supporting data from the family, a more conservative approach may be appropriate.

Serial re-evaluation of variants should be performed based on new phenotype data in the family pedigree, or new data regarding both specific variants and whole genes, the pathogenicity of which may have been up- or downgraded based on contemporaneous evidence. 115 , 194 – 196 , 204 Responsibility for re-evaluation is unclear, but the re-evaluation is best carried out in a center of expertise.

Genetic variants identified in deceased probands and subsequently in family members should be correlated with clinical findings to determine segregation patterns within the family.

SCD may occur in an individual without any prior medical history or medical data. As indicated above, a three-generation pedigree including examination of all potentially relevant cardiac and extracardiac features within the family should be performed by a cardiologist experienced in genetic heart disease or clinical geneticist experienced in cardiology in an effort to optimize the identification of subtle clinical features before defining the cause of death as unknown. Specific triggers (eg, competitive athlete, emotional or physical stress, drug use, swimming, acoustic triggers, seizure) leading to the SCD event may help focus clinical and genetic investigation. Collection of blood and/or suitable tissue for molecular autopsy/postmortem genetic testing is recommended in all victims of SUD. 2, 10 Local protocols are recommended to ensure proper handling and extended storage of biosamples, as well as processes and consent allowing for contact of families for genetic testing in the future.

Family members of sudden death victims where the phenotype remains unknown should be instructed to request re-evaluation of the index event in case of future developments in the family. Clinical signs or symptoms in first-degree family members leading to a suspected phenotype of a sudden death victim may occur over many years and sometimes skip generations, so an unknown phenotype might prompt more extensive questioning of older family members.

In an SCD case where the phenotype remains unknown after expert evaluation, re-evaluation of family members to assess for new information that may impact diagnosis should be performed periodically, although the yield is low. 205 We suggest every 3 to 5 years, but shorter intervals should be considered if there is more than one SCD event in the family. Periodic re-evaluation should be stopped for individuals after age 45 years, unless the decedent died in this age range or new findings emerge.

Three scenarios may trigger arrhythmia syndrome–focused genetic evaluation of SCD even if the phenotype remains unknown: 1) documented arrhythmic death suggestive of an arrhythmia syndrome; 2) specific triggers associated with familial arrhythmia syndromes; 3) young age (<40 years) ( Figure 13 ).

Families should be counseled about the expected benefits and potential outcomes of genetic investigations prior to testing. When genotyping is performed, the identification of a genetic variant as causal remains challenging and requires reassessing the correct classification or potential reclassification periodically. With rapidly advancing genotyping technologies and the availability of large gene panels, the identification of genetic variants of uncertain/unknown significance becomes more frequent. Medical uncertainty in general elicits a variety of responses from patients. It is important to consider patients' responses to the ambiguous nature of genetic testing. Medical professionals ordering genetic testing should be prepared for the possibility of their patients' misinterpretation of such results. Pre-test counseling should include a discussion of the possibility of a variant of uncertain significance and what it would mean for the patient's care and its potential psychosocial impacts. When a variant of uncertain significance is found, post-test counseling should include additional education and a discussion of the variant's implications and medical management recommendations based on the results. If identified variants are considered likely pathogenic or pathogenic, cascade testing across the family should be offered to identify at-risk individuals with no current clinical features. Cascade testing should not be performed with variants of uncertain significance; however, careful investigation within a multidisciplinary team (including genetic counseling) may allow eventual reclassification of the variant so that it may then be used for cascade testing (see Sections 3 and 4 ).

Recommendations for investigation of sudden death: genetic evaluation where the phenotype is unknown

Collection and storage of blood and/or suitable tissue for postmortem genetic testing is recommended in all victims of SUD. In a large number of cases, the phenotype underlying SCD remains unknown despite comprehensive evaluation of the victim and their family. In SCD cases where the phenotype is determined as unknown after expert evaluation, re-evaluation should be performed periodically to assess for new information that may impact diagnosis. While hypothesis-free genetic testing is not indicated in cases of SCD where the phenotype remains unknown, arrhythmia syndrome–focused genetic evaluation of SCD is advised if 1) an arrhythmic death is documented suggestive of an inherited arrhythmia syndrome, 2) specific triggers associated with familial arrhythmia preceded the SCD, and/or 3) SCD occurred at young age.

Genetic diseases may express with reduced penetrance. Hence, a negative clinical screening does not exclude the (silent) presence of a genetic disorder. It is, therefore, reasonable to suggest repeated screening with a time interval between 3 to 5 years until at least age 45 years. Studies demonstrating the yield of clinical screening after the sudden death of a close relative usually report only the result of the first screening. 25 , 70 , 80 , 85 , 205 , 206 Responsibility for the repeated (infrequent) screening lies in the hands of the individual, but the local team should make sure that he/she is appropriately informed.

In an SCD case where the phenotype is unknown, arrhythmia syndrome–focused genetic testing of the proband should be considered if 1) documented arrhythmic death (such as torsades de pointes arrhythmias leading to ventricular fibrillation) is suggestive of an arrhythmia syndrome, and/or 2) SCD is preceded by specific triggers (eg, competitive athlete, emotional or physical stress, swimming, drug use, acoustic triggers, seizure) associated with familial arrhythmia syndromes. 10 , 25 , 70 , 80 , 85 , 205 , 206 , 212 Collection and storage of blood and/or suitable tissue for postmortem genetic testing is recommended in all victims of SUD irrespective of an identified phenotype at the time of death. Long-term storage of biosamples of SCD victims is recommended in expert centers to allow for genetic testing if indicated at present or in the future. 10 , 25 , 70 , 80 , 85 , 205 , 206 , 212

In an SCD case where the phenotype is unknown, arrhythmia syndrome–focused genetic testing of the proband can be considered if SCD occurred at young age. Testing for cardiomyopathy genes (such as LMNA ) has been studied and can increase the diagnostic rate, although it should be recognized that the yield is lower. 10 , 25 , 70 , 80 , 85 , 205 , 206 , 212

Hypothesis-free genetic testing is not indicated in cases of SCD where the phenotype remains unknown. Genetic testing using any range from large unfocused gene panels to whole-exome or whole-genome sequencing in the absence of a clinical phenotype or diagnosis may be considered in the context of a scientific effort but is not recommended for routine patient care and counseling. 213 , 214 The aim of discouraging hypothesis-free testing in clinical settings is to reduce the misinterpretation of genetic variants and their causality, specifically, variants of uncertain significance. A specific problem in this field is the nonuniformity in calling variants across different laboratories. 215

Figure 14 summarizes the recommendations from Section 6 .

When an individual has been resuscitated after SCA, the clinician must try to define the likely underlying cause, similar to that discussed in Section 6.1 focused on SCD. This should include information on age, sex, past medical history, recent symptoms, activity or emotional status at the time of SCA (eg, sleeping, exercising, or emotion), time of onset (eg, morning or night), and environment (eg, public or private location), exposure to medicinal or recreational drugs (particularly those that block potassium or sodium channels) or alcohol, and a detailed family history of three generations at least.

To focus the history taking, one must consider the differential diagnosis of SCA, which is similar to that discussed in Section 6.1 . If the SCA was observed, a description of the event by the observer

can add useful information. All records of the primary event, including initial rhythm recordings and details of the resuscitation, should be collected in addition to any prior ECGs and imaging studies. A family history of heart disease, syncope, or sudden death may point to a genetic cause. A study by Waddell-Smith et al. 77 showed that while inpatient cardiology teams identified a familial condition in only about 8% of cases, nurses trained in taking a family history detected a familial condition in 32% of cases. Thus, although most patients with SCA will be in a nonspecialty environment at first, ideally specially trained members of the cardiology and genetic counselor team with experience in genetic heart disorders should be utilized to elicit relevant details of the family history. Practical educational assistance can be found at <https://www.primarycaregenetics.org> .

#### Recommendations for investigation of SCA survivors: personal and family history

Observational studies have demonstrated that the cause of SCA can be determined in a substantial proportion of patients. Historical features, especially age, coronary risk factors, symptoms, activity at the time of SCA, exposure to drugs, and family history frequently provide important clues to the diagnosis and point the way to further investigation.

In a study of 37 patients with history of SCD, cardiomyopathy, or ventricular tachycardia, a family history obtained by specially trained personnel was far more likely to elicit a history of inherited cardiac disease than one obtained by an inpatient cardiac team. 77

A detailed history is a crucial component of diagnosing the cause of SCA. Studies utilizing a comprehensive, systematic approach including history, physical examination, ECGs (eg, 12-lead ECG, treadmill, 24-hour Holter, signal-averaged ECG, if needed), cardiac imaging (eg, coronary angiography, echocardiogram, CMR, CT), provocative testing, electrophysiological study, cardiac biopsy, and genetic testing have shown that a diagnosis can be established in a substantial proportion of survivors of SCA. 216 – 219

The next step in cardiac evaluation of SCA survivors is physical examination. The main purpose is to identify signs of syndromic and nonsyndromic diseases that can be associated with SCA. For example, obesity and/or the presence of xanthomata may indicate an increased likelihood of premature coronary atherosclerosis. 221 , 222 Syndromic features that may be relevant to genetic disorders include woolly hair and palmoplantar keratoderma (arrhythmogenic right ventricular cardiomyopathy), 140 joint contractures (Emery-Dreifuss muscular dystrophy), 223 muscle weakness and atrophy (lamin A/C and desmin cardiomyopathies, Triadin knockout syndrome), 224 micrognathia, syndactyly, clinodactyly (Andersen-Tawil and Timothy syndromes), 224 chest and limb deformities, and tall stature (Marfan syndrome). 225 , 226

Fever, hypothermia, dehydration, and signs of drug abuse may be detected on physical examination, although these signs may be confounded by neurological impairment after cardiac arrest. These factors can trigger life-threatening arrhythmia in genetic heart disease. For example, fever has been associated with malignant arrhythmias in 6% of patients with cardiac arrest and Brugada syndrome. 136

Cardiac murmurs can raise the suspicion of left ventricular outflow tract obstruction, mitral mid-systolic click (valve prolapse) with or without regurgitation, and Ebstein anomaly. Signs of pulmonary edema and hepatosplenomegaly can be detected in patients with severe systolic myocardial dysfunction. Importantly, examination findings may be affected by the SCA event and evolve during a hospital stay, requiring repeated physical examination to determine whether findings are related to the cause of SCA or the effect of SCA.

#### Recommendation for investigation of SCA survivors: examination

There are no data describing the usefulness of physical examination in resuscitated SCD. However, it is a first basic step in the diagnostic process that will focus subsequent complementary

investigations.

Protocols for clinics investigating SCA survivors support the use of physical examination as the first step of clinical overview. 10 , 216 , 227

In most emergency settings, a patient resuscitated from cardiac arrest in whom myocardial infarction is suspected will have undergone a coronary reperfusion strategy to treat acute occlusion. It is critical to obtain blood tests (cardiac enzymes, inflammatory markers, glucose, serum electrolytes, and white blood cell count) and pertinent toxicological analysis at presentation. The latter may include testing for drugs of abuse such as ethanol, opiates, and stimulants, as well as levels of prescribed medication that may prolong QT interval/QRS duration or cause respiratory depression. 228 While assisting in the acute management of a resuscitated patient, these results will also help to differentiate acute myocardial injury as a cause of cardiac arrest (eg, ischemia without clear evidence of coronary occlusion or myocarditis) and pick up other reversible causes such as drug overdose, electrolyte imbalance, or endocrine and metabolic disorders. Retention and storage of suitable blood samples on patient arrival in the emergency department will allow subsequent diagnostic evaluation including DNA extraction and analysis in a patient who dies prior to diagnosis or for later clinical and family review. In some cases, this may be the only opportunity to obtain genetic material for analysis.

In an OHCA, the use of AEDs is ever more widespread and increases survival. 229 Sensitivity for the diagnosis of cardiac rhythm at the time of arrest is about 99%. 230 Therefore, routine inspection of data from AED recordings may improve the quality of diagnosis (see Section 6.2 ). The underlying rhythm of cardiac arrest may provide information on the arrhythmogenic mechanism, assist in diagnosis, and eventually indicate any misdiagnosis of rhythm. 231 Any ECG tracings from emergency services, as well as recordings from interrogation of CIEDs or wearables can also contribute to diagnosis. 163

The 12-lead ECG in sinus rhythm or during arrhythmia recurrence is fundamental to the diagnostic investigation and should be repeated daily during recovery. 216 It may support diagnoses of primary electrical disorders, pre-excitation, and heart muscle diseases. However, abnormalities of cardiac conduction and repolarization may result from myocardial injury during the cardiac arrest and patients undergoing post-arrest hypothermia protocols may have transient ECG changes including QT prolongation and J point elevation that should be interpreted with caution. Information about electrolyte levels, drug prescription, and body temperature should be added to the ECG to prevent misinterpretation of such ECG abnormalities. A high precordial lead ECG is an inexpensive tool to increase detection of Brugada syndrome pattern. 184 , 232 – 235

In addition to standard ECG, a signal-averaged ECG may demonstrate late potentials. Two or more abnormalities in the absence of a prolonged QRS duration ( $\geq 110$  ms) on the standard ECG is a minor diagnostic criterion for arrhythmogenic cardiomyopathy and suggests ventricular depolarization abnormality. 4

Continuous heart rhythm monitoring is recommended during hospitalization due to the transient nature of some arrhythmias. Recording the onset (including pause-dependent or tachycardia-associated initiation) and late and short-coupled ventricular ectopics as triggers for torsade de pointes, polymorphic ventricular tachycardia, or ventricular fibrillation will elucidate cardiac arrest mechanism and likely diagnosis. 236 Evidence of dynamic ST elevation associated with chest pain may also indicate likelihood of coronary vasospasm. 216

Echocardiography is the screening tool of choice for structural heart disease, although early myocardial dysfunction may be present after cardiac arrest and, if present early, the test should be repeated later during the patient's convalescence. CMR allows detection of inflammatory diseases, such as myocarditis and sarcoidosis, through recognition of subepicardial edema. Identification of an inflammatory etiology is important, as it may be self-limiting or treatable. If sarcoidosis is

suspected, then positron emission tomography–CT scanning may be indicated. 237 The presence of subendocardial edema would suggest ischemic injury. 238 Late gadolinium enhancement indicates chronic fibrosis, permitting detection of a cardiomyopathic etiology, and can also contribute to the diagnosis of mitral valve prolapse associated with a risk of SCD. 239 , 240 Coronary imaging (at any age) will be important to exclude coronary artery disease not investigated at presentation as an emergency and ensure that an anomalous coronary circulation or coronary dissection is not missed. 241 This may be by cardiac catheterization or by CT coronary angiography. 242 Coronary angiography will only be required in select pediatric and young survivors.

Patients in whom the cause of their SCA remains undiagnosed may require periodic re-evaluation of the above investigations, as features may develop later that point to a cause of their SCA (similar to Section 6.5 , Recommendation 1).

Recommendations for investigation of SCA survivors: baseline investigations

Systematic clinical testing is paramount in SCA survivors. This includes blood testing, toxicology, ECG, signal-averaged ECG, high precordial lead ECG, continuous ECG monitoring, echocardiography, and coronary imaging. If the diagnosis remains elusive and cardiac arrest is deemed unexplained, then CMR is important to identify subtle forms of cardiomyopathy or acquired structural disease.

The usefulness of blood testing and toxicology is by consensus, and a diagnostic role is unquestionable. 2 , 10 , 228 Viral studies may be useful, but no systematic evidence is available as yet. 249 A potential role for biomarkers specific for one of the arrhythmia syndromes is anticipated. 250 , 251

Results of a forensic study indicate the value of postmortem CIED interrogation to define the cause and timing of death more accurately and to detect potential CIED-related safety issues. CIED interrogation in unexplained deaths clarified the manner of death in 60.8% of the cases including cardiac and nonarrhythmic death and device concerns. 163

Sensitivity for the diagnosis of cardiac rhythm via AED at the time of arrest is about 99%. 230 However, the AED seldom catches the initial rhythm of cardiac arrest and therefore may not contribute to the etiology of SCA.

Primary electrical disorders and specific cardiomyopathies may be detected by conventional ECG. 2 , 10 , 216 , 252 ECG findings in the immediate aftermath of a cardiac arrest, other than ST-segment elevation indicative of an acute coronary syndrome, may, however, have poor diagnostic accuracy. 253 , 254 These could be caused by abnormal repolarization following electrical cardioversion, 255 metabolic and electrolyte abnormalities, or even subarachnoid hemorrhage. 256 Therapeutic hypothermia may lead to misleading ECG changes such as prolongation in PR, QRS and QT intervals, and J point elevation. 257 , 258 Interpretation of ECGs obtained immediately after resuscitation/defibrillation should be performed with great caution.

Although there are no data describing directly the value of high precordial lead ECGs in SCD survivors, there is ample evidence of an increased yield of the type 1 Brugada ECG pattern. 189

Cardiac monitoring during short-term follow-up demonstrates an arrhythmogenic mechanism of SCD in some registries. 216 , 236 , 246

Signal-averaged ECG is part of the Task Force Criteria for arrhythmogenic right ventricular cardiomyopathy. 4 Signal-averaged ECG has been proposed as useful in other conditions (ie, Brugada syndrome), although systematic evaluation has not been performed.

Echocardiography is a valuable screening tool for detection of arrhythmogenic cardiomyopathy and other structural abnormalities useful in elucidating the cause of SCA. 85 Patients with functional abnormalities on initial echocardiogram should have this test repeated after recovery, to allow for the effects of the SCA itself and drugs used around the time of the arrest to wear off.

9. and 10. The utility of CMR has been evaluated in a series of studies involving survivors of unexplained arrest and has repeatedly been shown to provide significant incremental diagnostic value. A study of 137 individuals with unexplained aborted cardiac arrest found that CMR provided a diagnosis or identified an arrhythmic substrate in 76% of individuals, including an infarct pattern suggestive of occult myocardial infarction in 44%. Notably, the presence of late gadolinium enhancement, reflective of myocardial fibrosis, was associated with a 6.7 hazard ratio ( $p < 0.001$ ) of recurrent arrhythmic events on multivariate analysis. 248 The presence of subendocardial edema would suggest ischemic injury even when initial coronary imaging excludes significant obstruction. Coronary vasospasm and dissection might be misdiagnosed, and coronary re-evaluation may then be reconsidered. 238 The frequency of occult infarcts is not insignificant, 248 although the risk of SCD in patients with myocardial infarction and nonobstructive coronary arteries is low. 259 For Recommendation 10, primary electrical disease is not referring to an established diagnosis of long QT syndrome or CPVT, where MRI is unlikely to be of use.

Coronary artery disease is the leading cause of SCD in adults and might be treatable. Furthermore, coronary dissection and anomalies may also be relevant in this age group as well as in younger patients.

Once a cardiac arrest survivor has undergone initial thorough baseline evaluation, most overt acquired or genetic etiologies will have been diagnosed. However, concealed disorders may be uncovered by provocative maneuvers such as lying to standing ECGs, exercise ECG testing, epinephrine challenge, sodium channel blocker challenge, or ergonovine and acetylcholine testing. Some may even be employed in a resuscitated cardiac arrest survivor who is unlikely to survive due to neurological injury.

Exercise testing may uncover ventricular arrhythmia relevant to the cause of cardiac arrest. For example, evidence of monomorphic ventricular tachycardia arising from the right ventricle is part of Task Force Criteria for diagnosis of arrhythmogenic right ventricular cardiomyopathy (arrhythmogenic cardiomyopathy). 260 – 262 Exercise may also uncover concealed epsilon waves or even a type 1 Brugada ECG pattern. 263 , 264 The generation of bidirectional ventricular ectopy or tachycardia and/or polymorphic ventricular tachycardia in the absence of ischemia, structural disease, or digoxin toxicity is typical of CPVT 265 and has been evaluated in the Cardiac Arrest Survivors with Preserved Ejection Fraction Registry (CASPER) protocol. 216 Abnormal dynamics of repolarization in response to challenges may also inform the likelihood of underlying long QT syndrome. To this end, maximum QT prolongation (QT stretch) and T-wave morphology changes during lying to standing and then return to baseline heart rate (QT stunning) may be useful markers, although they are less specific in children. 266 – 268 These discriminate well in genotyped families and may add to diagnostic utility, but evaluation in unexplained cardiac arrest survivors has not been undertaken. 269 The recovery phase of exercise may also reveal readily measurable QT prolongation and T-wave abnormalities and has been validated in families with long QT syndrome for prediction of genotype 270 , 271 and in cardiac arrest survivors 216 such that a QTc >480 ms at 4 min of recovery forms part of the long QT syndrome risk score. 272

Epinephrine challenge with ECG monitoring has been advocated as an alternative to exercise testing for the diagnosis of long QT syndrome and CPVT, particularly where the patient is unable to exercise. In long QT syndrome, QT prolongation and secondary T-wave changes have been able to discriminate LQT1 and LQT2 patients from unaffected family members even though they have normal baseline QTc intervals. 273 , 274 The test has been assessed in unexplained cardiac arrest survivors and has suggested a low specificity for long QT syndrome. 275 The finding would be unlikely to provide a secure diagnosis in isolation and was proposed as useful in association with exercise testing and genetic testing. Indeed, in normal subjects, pharmacological sympathetic

stimulation does produce significant prolongation of QTc. 276 Epinephrine testing has also shown some diagnostic utility for CPVT in cardiac arrest survivors by inducing ventricular ectopic activity, bidirectional couplets, and ventricular tachycardia. 275 However, there is uncertainty as to the ideal cutoff for epinephrine-induced arrhythmia, and the diagnostic sensitivity compared to exercise testing in CPVT families is low. 277 Isoproterenol challenge for the diagnosis of arrhythmogenic right ventricular cardiomyopathy has also been advocated but has not been tested by other groups or in the unexplained cardiac arrest survivor without overt phenotype. 278

Sodium channel blocker challenge (ajmaline, procainamide, flecainide, and pilsicainide) has been used extensively for investigating the possibility of Brugada syndrome in cardiac arrest survivors, although these are mainly reported in series of patients with a strong suspicion of Brugada syndrome 279 – 281 rather than in unexplained cardiac arrest survivors. 282 The use of leads V1 and V2 in the second and third intercostal space or high right precordial ECG leads during provocation increases the diagnostic yield. 233 , 283 In the CASPER registry, there was a yield from procainamide challenge, 284 although this may underestimate the true burden, as different sodium channel blockers have different potencies for inducing the type 1 Brugada ECG pattern. For example, ajmaline is associated with an odds ratio of 8 for inducing the type 1 Brugada ECG pattern compared with procainamide 285 and a 4% yield in a small group of “healthy” controls. 286 Furthermore, while recent consensus guidelines would give a definite diagnosis to a cardiac arrest survivor with a type 2 or 3 pattern converting to a type 1, the implication of a drug-induced type 1 pattern without a baseline type 2 or 3 is not addressed. 152 Yet cardiac arrest survivors from CASPER without a type 2 or 3 pattern had positive procainamide challenges. 284 Other tests such as the full stomach test have been proposed but have not been taken up in general. 287

Different approaches for provocation of an underlying repolarization abnormality have been employed including drug challenge with quinidine and sotalol and mental stress tests. 288 – 291 These may offer utility in the future but have not been tested in the cardiac arrest survivor.

Coronary vasospasm, while a recognized cause of cardiac arrest, may not be picked up clinically at presentation. 292 Ergonovine or acetylcholine challenge has been proposed as a Class 1 indication by recent guidelines 293 and was employed selectively in the CASPER experience. 216 Hyperventilation has also been used as a diagnostic test. 294 Recent experience from the Paris Sudden Death Expertise Center investigators suggests that pharmacological challenge is useful for diagnosis and could be better employed. 295 Nonetheless, there are few centers with extensive experience in the use of the test, and there is a possibility of false-positive findings in the cardiac arrest survivor population.

Adenosine challenge has been used to unmask pre-excitation that may otherwise be missed. 296 , 297 In the absence of other causes, it will indicate the need for electrophysiological study to evaluate the risk of the accessory pathway (rapidity of antegrade conduction) followed by ablation therapy. However, electrophysiological study is not routinely included in the workup of unexplained cardiac arrest. 8 Indeed, previous consensus guidelines proposed a Class 3 indication when assessing a suspected primary electrical disorder. 8 It does not add additional diagnostic or prognostic value unless there is evidence to indicate otherwise. For example, pre-excited atrial fibrillation, bundle branch re-entrant ventricular tachycardia, and rapid supraventricular tachycardias that degenerate into ventricular fibrillation have all previously been described as culprits requiring an invasive approach to diagnosis and curative ablation therapy. 298 , 299 Electroanatomic voltage mapping of the right ventricle is a discretionary tool that may be considered to detect evidence of subclinical arrhythmogenic right ventricular cardiomyopathy. 300 More recently, extensive endocardial and epicardial mapping of unexplained cardiac arrest cases has been employed to identify cases with either Purkinje triggers and/or subtle depolarization abnormalities that may be suitable for ablation therapy. 301

Recommendations for investigation of SCA survivors: provocative testing

Concealed Brugada syndrome, long QT syndrome, CPVT, arrhythmogenic cardiomyopathy, pre-excitation, and coronary vasospasm may be uncovered by provocative maneuvers in the cardiac arrest survivor whose cause of cardiac arrest remains unknown after baseline clinical, ECG, and imaging investigations. Exercise ECG testing and sodium channel blocker challenge appear to offer most potential utility, whereas lying to standing ECGs; epinephrine, isoproterenol, and adenosine challenge; and hyperventilation, ergonovine, and acetylcholine testing may be considered in specific patients. Electrophysiological study and electroanatomic mapping may be useful to provide patient-specific insights into the mechanism of cardiac arrest and offer therapeutic options but should be avoided in the routine investigation of channelopathy. However, data in general are limited to case reports or case series, some with validation cohorts, and there are no randomized studies.

Exercise testing is a versatile, straightforward, and readily available test that may yield diagnoses due to the arrhythmic challenge or effects on depolarization and repolarization. As such, it should be a standard part of the investigative armamentarium. 216 , 220 , 263 – 265 , 270 – 272 , 302 , 303

Lying to standing ECGs may offer some insights into the likelihood of long QT syndrome, but in isolation a positive result may not fulfill diagnostic criteria and therefore may only complement the evaluation. 266 – 270

Similarly, epinephrine challenge has low specificity for long QT syndrome and may be insensitive for CPVT. In isolation, a positive result may not fulfill diagnostic criteria and therefore may only complement the evaluation. 273 – 275 , 277 , 304 There are risks to this test, both in inducing ventricular arrhythmia and in false diagnosis.

Sodium channel blocker testing has a clear role to play in evaluating the cardiac arrest survivor, although the implication of a positive result in patients with or without a high prior likelihood of Brugada syndrome is unclear, eg, male vs. female, cardiac arrest during sleep vs. exercise, type 2 or 3 ECG pattern vs. no Brugada pattern at baseline. This reflects the lack of a gold standard for the diagnosis of Brugada syndrome, and therefore the utility of sodium channel blocker challenge in patients without a clear prior likelihood of Brugada syndrome is less certain and merits a lower utility in the evaluation cascade. 233 , 273 – 275 , 277 , 286 Furthermore, sodium channel blocking agents differ in their sensitivity and specificity for inducing ECG changes and availability varies worldwide (see Section 7.4 text). 285

The utility of ergonovine or acetylcholine challenge or hyperventilation testing in all cardiac arrest survivors is unclear, as there are no studies of systematic testing in all cardiac arrest survivors. In particular, the specificity of the test is unknown, especially in the cardiac arrest survivor population. It can, however, lead to diagnoses when employed in a protocol and therefore should be considered as part of the armamentarium until more evidence is available. 216 , 293 – 295

Adenosine challenge has not been tested systematically in cardiac arrest survivors, but there are limited data suggesting that it will uncover concealed pre-excitation that may cause cardiac arrest in the setting of rapidly conducted pre-excited atrial tachyarrhythmias. 296 , 297

There are limited case series describing the use of electrophysiological study in diagnosis and treatment of pre-excited atrial arrhythmias and bundle branch re-entrant ventricular tachycardia. 298 , 299

Studies of endocardial mapping report a higher sensitivity for detection of arrhythmogenic cardiomyopathy, but this has not been explored in SCA survivors. 300 Sensitivity and specificity are unknown, particularly in patients where no other tests are abnormal.

Extensive endocardial and epicardial mapping of unexplained cardiac arrest cases has been employed to identify cases with either Purkinje triggers and/or subtle depolarization abnormalities that may be suitable for ablation therapy. 301

Although most SCA survivors will have an indication for an implantable defibrillator for secondary prevention of a cardiac arrest, genetic evaluation may influence final diagnosis, treatment recommendations, and family screening ( Figure 15 ). In some cases, genetic evaluation may enable therapy specific to the disease mechanism. Recent technological advances in genetic evaluations, establishment of reference databases of genetic variants, systematic annotation of causal genes, 305 and standardization of variant interpretation 306 have enabled efficient and comprehensive genetic assessment. However, the pace of genetic discovery and variant interpretation is evolving rapidly, creating a complex landscape surrounding genetic evaluation of SCA survivors.

The decision to pursue genetic evaluation is an individualized one in which the patient, with proper (clinical and genetic) counseling, must weigh the benefits, limitations, and personal and familial implications. The genetic basis of SCA and most individual conditions that predispose to SCA remain incompletely understood. As such, caution is advised when interpreting negative genetic test results or when examining genes with low likelihood of causing SCA. Prior clinical practice guidelines have specifically addressed the diagnostic considerations regarding specific genetic cardiovascular conditions. 4 , 10 , 307 The ClinGen represents a systematic effort to enumerate causal genes and variation related to human disease, including SCA-related conditions. 305 Previous reports have been developed for Brugada syndrome, 195 long QT syndrome, 196 and hypertrophic cardiomyopathy. 202

The yield of genetic testing varies substantially by condition. 188 Variant interpretation may differ by laboratory 215 despite recent efforts to standardize variant interpretation. 306 Given the rapid changes in available technology to evaluate the genome, complexities in variant interpretation, and nuances in the ethical and legal framework surrounding genetic testing in some settings, 66 , 115 , 215 , 308 – 310 it is recommended that SCA survivors undergoing genetic assessment have evaluations performed at centers with multidisciplinary experience in counseling, variant interpretation, and management of genetic heart disease (see Section 3 ). For SCA survivors who have undergone genetic testing, an offer of periodic re-evaluation of the genetic test results is advocated (similar to Section 6.4 , Recommendation 4).

#### Recommendations for genetic evaluation of SCA survivors

SCA can be caused by diverse etiologies, some of which may be predominantly or partially influenced by genetic predisposition. Whereas a thorough clinical evaluation leads to a diagnosis of the cause of SCA for most individuals, the cause of SCA may remain uncertain in others. 216 In some cases, genetic evaluation of SCA survivors can confirm a molecular etiology that predisposed to the SCA event, support the diagnosis of a specific phenotype, influence management, and facilitate screening in family members at risk via cascade genetic testing.

Genetic testing, after appropriate genetic counseling and informed consent, may facilitate identification of a molecular cause of SCA by identifying pathogenic variants in genes associated with specific phenotypes and fulfilling formal disease-based diagnostic criteria. 4 , 10 Examples of scenarios in which discovery of a genetic cause of SCA may influence management recommendations include administration of beta blockade for patients with long QT syndrome, 311 sodium channel inhibition in long QT syndrome type 3, 312 – 314 flecainide administration for patients with CPVT, 315 , 331 or exercise restriction recommendations in patients with arrhythmogenic cardiomyopathy. 316 , 317 Genetic evaluation may also influence family screening by facilitating cascade genetic testing and clinical surveillance in relatives at greatest risk for disease.

2. and 3. Genetic tests have variable yield and may result in discovery of variants of uncertain clinical significance, which can be frequent and challenging to interpret. Using genetic tests that comprise well-established genes related to a suspected or diagnosed genetic phenotype is most likely to result in discovery of disease-causing variants in an individual or family, while minimizing the probability of discovering a variant of uncertain clinical significance. In contrast, genetic tests with more comprehensive genomic coverage may lead to moderately increased diagnostic yield but at the expense of increased rates of discovery of variants of uncertain clinical significance. 84 , 308 Nevertheless, given a rapidly evolving understanding of the molecular causes of specific phenotypes and increased yield of broader genetic assessment, 82 , 84 , 246 , 323 , 324 tests that include broader coverage may be considered in select circumstances, such as when a heritable phenotype is being mapped within a family or assessment for de novo variation is sought through sequencing of multiple family members. The latter examples would only be pertinent when it becomes apparent that a familial trait is likely or there has been exome sequencing in a trio of confirmed parents and the index case.

Genetic evaluation appears to be highest for individuals with a phenotype consistent with a genetic cause 84 , 323 , 324 and is lower among SCA survivors without a clearly identifiable genetic phenotype. 82 , 84 , 246 , 323 – 326 Nevertheless, individuals with an idiopathic cause of SCA may eventually develop a diagnosis of a genetic etiology during long-term follow-up. 327 Genetic testing can identify variants during the concealed phase of a genetic disease such as arrhythmogenic cardiomyopathy, 328 or in individuals with conditions otherwise regarded as nongenetic such as in drug-induced long QT syndrome. 329 As such, a low but non-negligible yield for genetic testing appears to be present among individuals with idiopathic SCA.

In individuals with a well-established nongenetic cause of SCA, the routine use of genetic evaluation is not recommended owing to the potential for discovery of variants of uncertain significance and misdiagnosis.

Figure 16 summarizes the recommendations from Section 7 .

The sudden death of a young, apparently healthy individual raises many questions for family members. Apart from mourning and the question “could we have done something to avoid this,” a very relevant question is whether family members could be affected as well. Roles for health care providers include providing psychological support for the family, identifying a cause for the sudden death, and understanding the implications for family members (see Sections 4 and 5 ). For this chapter, it is especially relevant that families need support to organize clinical and genetic testing for family members and sometimes postmortem genetic testing of the deceased. 2 , 101 , 332

Currently, many cases and even familial forms of unexplained cardiac disease remain insufficiently investigated. 333 When clinical symptoms indicate that a cardiomyopathy or arrhythmia may have contributed to the death, further steps are needed to specify the diagnosis. If blood or tissue of the deceased is available, DNA testing can be done for a range of arrhythmia syndromes and cardiomyopathies, nowadays often using gene panels. Without a specific diagnosis in the deceased, clinical investigation of the first-degree relatives (parents, siblings, and children) can identify a person with similar symptoms or signs, although sometimes mild. This relative of the deceased can be the proband for DNA testing and thus provide the key to a diagnosis for the family.

Efforts are needed to increase the proportion of postmortem examinations (either forensic or medical autopsy) to clarify whether or not an underlying cause can be suspected or proven. The postmortem result should be communicated to the family as per local protocols. Without a postmortem diagnosis, efforts are needed to evaluate eventual clinical symptoms of the parents and other first-degree family members. 80 , 81 , 143 , 144 , 334 This is outlined in Section 6 . If DNA of the deceased person can be used for testing, or a relative who has similar symptoms can be tested, a monogenic form of cardiomyopathy or arrhythmia may be recognized that may also be present in family members. Typically, first-degree relatives are at 50% risk of carrying the same

pathogenic mutation, since many of these conditions follow an autosomal dominant pattern of inheritance.

Depending on whether or not autopsy has been performed, whether DNA is available, whether a relative is already diagnosed with a cardiogenetic condition, and whether symptoms were noticed during life, there are different situations possible: 1) sudden death, with clinical observations or DNA testing suggesting a specific diagnosis, and 2) sudden death, with no cause identified. The first situation will be discussed in Section 8.2 , the second in Section 8.3 ( Figure 17 ).

In cascade testing, after the postmortem diagnosis in an index patient by a DNA test, an invitation can be sent to parents, brothers and sisters, and children. 335 If any of the relatives is also diagnosed with the condition, a next circle of first-degree relatives is invited. If one of the first-degree relatives is not available (either deceased or not wanting to participate), second-degree relatives are invited, for instance, children of a deceased sibling. This systematic approach is very effective in autosomal dominant conditions, since first-degree relatives are at 50% a priori risk to carry the same pathogenic mutation and second-degree relatives at 25% risk. For minors, the age at which treatment starts determines the age before which DNA testing is advised. This may differ between countries and conditions. DNA testing of minors is not advised if the result would have no consequences in childhood.

Presymptomatic DNA testing makes it possible to organize preventive measures, such as regular cardiological follow-up, use of medication (eg, beta blockers), lifestyle advice (eg, avoid intensive sports), implantable cardioverter-defibrillators, or reproductive planning. 80 , 193 , 336 – 342 Presymptomatic DNA testing also puts an end to uncertainty and fear for those family members who test negative. 339 For them, follow-up investigations are no longer indicated.

In studies describing cascade screening, often the participation of relatives is limited. 343 – 348 This implies that many persons carrying pathogenic variants remain undiagnosed and are at continued risk of sudden death. Increasingly, geneticists and other stakeholders plea for an active approach to cascade testing for conditions where interventions are available. 349 – 351 Stakeholders agree on the importance of early diagnosis and informing the family. 2 , 340 , 352 Barriers to cascade screening include out-of-pocket expenses for the patient, limited resources for informing relatives, and privacy regulations. 339 , 351 , 353 , 354 To benefit from predictive, personalized, and preventive medicine, the roles and responsibilities of stakeholders in genetic testing as a preventive strategy need to be carefully aligned.

If clinical signs and symptoms suggest an inherited condition but no DNA test has been performed or no pathogenic variant has been identified, then history, examination, and clinical investigations of first-degree relatives are required to identify those at risk for SCD.

Recommendations for investigation of the family: cause identified—cascade testing, clinical and genetic investigations

A postmortem diagnosis in a victim of SCD implies the possibility of avoiding sudden death in relatives. This requires an active approach to inform parents, siblings, and children of the index case and offer clinical evaluation and potential genetic testing. A multidisciplinary service should support all elements of recognizing an inherited cardiac disorder in a victim (pathologist), identifying pathogenic mutations (geneticist), clinical evaluation and surveillance of carriers of the mutation (cardiologist), and supporting the patients and relatives (psychologist).

Since undiagnosed cardiogenetic conditions can be life-threatening and since interventions are available to reduce the risk of sudden death, first-degree relatives who are at 50% risk of carrying the pathogenic variant need to be informed about the possibilities of clinical investigations and genetic testing. 70 , 84 , 346 , 355 , 356 Second-degree relatives whose intervening first-degree relative refuses genetic testing (or is not available) should be offered testing due to a 25% risk.

Patients often can inform their relatives at risk, but medical professionals increasingly feel ethically responsible to support this process. 349 , 350 , 357 Supplying a letter for the patient to share with relatives is advised. Occasionally, directly contacting relatives may be possible in some jurisdictions. Studies have shown that there is substantial room for improvement of the uptake of family screening. Barriers include reluctance to consent to postmortem investigations, lack of information, and lack of funding for the services. While the services needed should involve many disciplines (pathologist, cardiologist, geneticist), the systematic approaching of family members fits the specialty of clinical genetics well. To achieve a high uptake, a systematic approach is needed.

The identification of at-risk individuals leads to the question of whether all require active treatment. Prospective clinical studies are needed to answer this question. To balance pros and cons for survival and for quality of life on treatment, long-term follow-up of persons following the suggested surveillance and interventions is needed. Precision prevention advice should build on this evidence.

In families with genetically affected individuals (with or without clinical findings), detailed prenatal counseling and guidance regarding inheritance patterns, variant penetrance, and risk should be offered, and other options including preimplantation genetic diagnosis should be explored. 336 – 338

Sudden death in the young is always a tragedy for those lost and the remaining family. When the cause of sudden death is not identified, either because there was no postmortem examination or because the autopsy was negative, significant anxiety in the family focuses on two major questions: why did the subject die suddenly and what risks apply to the remaining family members? An autopsy is typically requested, and in some jurisdictions mandated, but may not be completed because of cultural, family, or logistical limitations. When a postmortem examination establishes a cause of death, diagnosis transitions from SUD to death attributed to autopsy-related findings (see Section 8.2 ). When no underlying anatomic or toxicologic cause of death is identified with forensic autopsy, the description of the death goes from SUD to sudden arrhythmic death (syndrome)/SAD(S), since death is attributed to a presumed arrhythmia, or autopsy-negative SCD. In the case of nonspecific findings, follow-up of families should be similar to that with a negative autopsy. 184 , 359

When sudden death is classified as SAD or autopsy-negative SCD (ie, SUD), the differential diagnosis includes a breadth of inherited conditions that are predominantly ion channelopathies, with latent cardiomyopathy a consideration based on subtle autopsy findings. Causes include long QT syndrome, CPVT, short QT syndrome, Brugada syndrome, and arrhythmogenic cardiomyopathy. Careful review by a trained cardiac pathologist is recommended to ensure that assigned causes or absent causes are accurate (see Section 6.3 ). 2

Recent studies have shown that a genetic evaluation of the deceased subject's DNA associated with a clinical evaluation of first-degree relatives of the deceased subject retrospectively identified the cause of death in 20–40% of cases. 70 , 80 , 81 , 143 , 144 , 189 , 3 3 4 In this situation, the identification of the cause of sudden death provides an explanation to the family and facilitates further cascade screening. In a second step, it will enable prevention measures in the family to limit the risk of a second death.

Recommendations for investigation of the family: cause not identified—clinical and genetic investigations

Screening of first-degree relatives of the SCD victim is informed by findings from the forensic investigation. Though the yield of genetic testing is relatively low, results should be applied to all first-degree relatives in conjunction with clinical assessment. In the absence of genetic results, screening tests should include a medical history, standard and high precordial lead ECG, 24-hour ambulatory monitoring, echocardiography, and exercise test, with select use of pharmacological provocation and advanced imaging.

Although autopsy is recommended, family screening after sudden death in young patients is effective even when an autopsy is not conducted. Broad screening of first-degree relatives with systematic testing is warranted. 80 , 81 , 143 , 144 , 334 Combining molecular autopsy with clinical evaluation in surviving families increases diagnostic yield. 70 The value of surveillance testing after negative evaluation is uncertain, though commonly undertaken until age 45 years (range: 40–50 years), with decreasing frequency with age. 85 The context of SCD, the family history, and existing findings should inform the potential merits of ongoing surveillance, including frequency and duration. The age at which and from which onward surveillance is warranted depends on the (suspected) underlying condition.

When decedent genetic testing detects a pathogenic or likely pathogenic variant, the result enables identification of all family members at risk of SCD. 80 , 81 , 143 , 144 , 334 It is important to combine genetic and clinical evaluation, especially when the pathogenicity of the detected variant is uncertain, to evaluate the correlation between the genetic finding and clinical diagnosis for each family. 83 If DNA is not available from the decedent and no clinical phenotype is present in the family, genetic testing of family members should be strongly discouraged.

Several studies have demonstrated that family screening should include at least a medical history, standard and high precordial lead ECG (to improve the detection of the Brugada syndrome), echocardiography, exercise test, 80 , 81 , 143 , 144 , 334 and Holter monitoring on a case-by-case basis.

CMR and 24-hour ambulatory monitoring of family members can inform diagnosis. 4 , 5

5. and 6. A careful assessment of the circumstances of sudden death may point to a specific diagnosis. Sudden death in a young or middle-aged male occurring during a febrile illness or sleep suggests the diagnosis of Brugada syndrome, whereas sudden death occurring in a subject during physical activity suggests the diagnosis of long QT syndrome or CPVT. As Brugada syndrome may be masked or intermittent in some patients, sodium channel blocker challenge may unmask the type 1 pattern and increase the effectiveness of family screening. It should be recognized that there is a potential high rate of false positives, as data on the specificity and sensitivity of the test are not available, and a positive ECG may be induced in 4–5% of normal subjects. 286 , 361 In the pediatric age group, a negative test may convert to a positive test after puberty. 362 Long QT syndrome may also be unmasked by standing, exercise test, epinephrine test, or mental stress test. 266 , 269 , 270 , 275 , 288 In first-degree relatives of young SUD victims with no manifest abnormalities during the initial examination, the risk of developing manifest inherited cardiac disease or cardiac events during follow-up is low. 205

Many of the recommendations in this document seem intuitive, obvious, and straightforward; however, much of what is being recommended within this document is seldom routinely performed even in well-resourced countries. Many decedents of SUD never receive an autopsy, and the evaluation of first-degree relatives of an SUD victim or an SCA survivor ranges widely from no evaluation to ordering a multitude of tests that are then repeated regularly and indefinitely. The extensive variability in practice indicates that developing common sense processes and multidisciplinary teams remains a considerable challenge in many areas and guidance is required. Developing these processes and teams involves leaders challenging many medico-political barriers that obstruct and delay best medical practice. We hope that this document will empower those who wish to achieve such changes for the better, including efforts for continuous improvement of practice. So, assuming the recommendations in this document are embraced and implemented, what are the next steps and future directions in this field?

Firstly, unlike cancer statistics and even SCD in the elderly due to coronary artery disease, the precise prevalence, epidemiology, and etiologies of either SUD or SCA in the young remain obscure in most countries. Only if these conditions become a notifiable event will the true scale and scope ever be captured.

Secondly, communities, states/provinces, and countries must advocate for and expect that a true comprehensive autopsy, including postmortem genetic testing (ie, molecular autopsy), occurs whenever an SUD occurs in a young person. The current dismal rate of autopsy must be reversed. Only when this becomes the standard of care will the true epidemiology/etiology of SCD be determined.

Thirdly, the basic occurrence and subsequent extent of an evaluation of the living, whether she/he is an SCA survivor or the first-degree relative of an SUD victim or an SCA survivor, must become standard of care. After initial cardiological evaluation (examination, ECGs, stress test, and echocardiogram) has been completed, the true contribution of SCA-predisposing genetic heart disease will be exposed. Furthermore, the composition and contribution of advanced investigations ranging from sodium channel drug provocation studies, to MRI, to genetic testing requires further study, as does the recommended interval for a repeat cardiological evaluation of the first-degree relatives when their first evaluation is either normal or inconclusive.

Finally, when such investigations are commenced, future studies to minimize the collateral damage from uncertain clinical findings and genetic “variants of uncertain significance” will be needed. While it is recognized that the correct necropsy diagnosis of the SUD decedent and the correct diagnosis of the SCA survivor may give patients and their families some answers and resolution to the event, the premature and erroneous diagnosis due to excessive confidence in or overinterpretation of clinical or genetic findings of uncertain significance can cause remarkable harm. If we are to “first do no harm,” the model of multidisciplinary teams with the expertise to correctly evaluate all the investigations and potential (mis)diagnoses should be made accessible to all.

Developed in partnership with and endorsed by the Asia Pacific Heart Rhythm Society (APHRS) and the Heart Rhythm Society (HRS). Developed in collaboration with and endorsed by the Association for European Cardiovascular Pathology (AECVP), the European Society of Human Genetics (ESHG), the Latin American Heart Rhythm Society (LAHRS), the National Society of Genetic Counselors (NSGC), and the Pediatric and Congenital Electrophysiology Society (PACES). Developed in collaboration with the European Heart Rhythm Association (EHRA).

Supplementary data (Appendix 3) associated with this article can be found in the online version at <https://doi.org/10.1016/j.hrthm.2020.10.010>.

Document Reviewers: Takeshi Aiba, MD, PhD; Allison L. Cirino, MS, CGC; Florence Fellmann, MD, PhD; Michael H. Gollob, MD; Yung-Kuo Lin, MD, PhD; Joaquín S. Lucena, MD, PhD; Ciorsti MacIntyre, MD; Jens Cosedis Nielsen, MD, DMSc, PhD, FESC, FEHRA; Juan David Ramirez, MD; Gregory Webster, MD, MPH, CCDS

Number value: 0 = \$0; 1 = ≤ \$10,000; 2 = > \$10,000 to ≤ \$25,000; 3 = > \$25,000 to ≤ \$50,000; 4 = > \$50,000 to ≤ \$100,000; 5 = > \$100,000.

Research and fellowship support are classed as programmatic support. Sources of programmatic support are disclosed but are not regarded as a relevant relationship with industry for writing group members or reviewers. AHA = American Heart Association; AECVP = Association for European Cardiovascular Pathology; NIH = National Institutes of Health; NSGC = National Society of Genetic Counselors.

Number value: 0 = \$0; 1 = ≤ \$10,000; 2 = > \$10,000 to ≤ \$25,000; 3 = > \$25,000 to ≤ \$50,000; 4 = > \$50,000 to ≤ \$100,000; 5 = > \$100,000.

Research and fellowship support are classed as programmatic support. Sources of programmatic support are disclosed but are not regarded as a relevant relationship with industry for writing group members or reviewers. AHA = American Heart Association; NIH = National Institutes of Health.

cardiovascular implantable electronic device

catecholaminergic polymorphic ventricular tachycardia

relationship with industry and other entities

26409257 10.1016/j.jacc.2015.09.001 2. Fellmann F , van El CG , Charron P , European recommendations integrating genetic testing into multidisciplinary management of sudden cardiac death . Eur J Hum Genet 2019 ; 27 : 1763 – 1773 . 31235869 10.1038/s41431-019-0445-y PMC6870982 3. Cronin EM , Bogun FM , Maury P , 2019 HRS/EHRA/APHRS/LAHS expert consensus statement on catheter ablation of ventricular arrhythmias . Heart Rhythm 2020 ; 17 : e2 – e154 . 31085023 10.1016/j.hrthm.2019.03.002 PMC8453449 4. Towbin JA , McKenna WJ , Abrams DJ , 2019 HRS expert consensus statement on evaluation, risk stratification, and management of arrhythmogenic cardiomyopathy . Heart Rhythm 2019 ; 16 : e301 – e372 . 31078652 10.1016/j.hrthm.2019.05.007 5. Brignole M , Moya A , de Lange FJ , 2018 ESC guidelines for the diagnosis and management of syncope . Eur Heart J 2018 ; 39 : 1883 – 1948 . 29562304 10.1093/eurheartj/ehy037 6. Al-Khatib SM , Stevenson WG , Ackerman MJ , 2017 AHA/ACC/HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death . Heart Rhythm 2018 ; 15 : e73 – e189 . 29097319 10.1016/j.hrthm.2017.10.036 7. Mont L , Pelliccia A , Sharma S , Pre-participation cardiovascular evaluation for athletic participants to prevent sudden death: position paper from the EHRA and the EACPR, branches of the ESC . Europace 2017 ; 19 : 139 – 163 . 27815371 10.1093/europace/euw243 8. Priori SG , Blomstrom-Lundqvist C , Mazzanti A , 2015 ESC guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death . Eur Heart J 2015 ; 36 : 2793 – 2867 . 26745817 10.1093/eurheartj/ehv445 9. Pedersen CT , Kay GN , Kalman J , EHRA/HRS/APHRS expert consensus on ventricular arrhythmias . Heart Rhythm 2014 ; 11 : e166 – e196 . 25179489 10.1016/j.hrthm.2014.07.024 10. Priori SG , Wilde AA , Horie M , HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes . Heart Rhythm 2013 ; 10 : 1932 – 1963 . 24011539 10.1016/j.hrthm.2013.05.014 11. Ackerman MJ , Priori SG , Willems S , HRS/EHRA expert consensus statement on the state of genetic testing for the channelopathies and cardiomyopathies . Heart Rhythm 2011 ; 8 : 1308 – 1339 . 21787999 10.1016/j.hrthm.2011.05.020 12. Zipes DP , Camm AJ , Borggrefe M , ACC/AHA/ESC 2006 guidelines for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: a report of the American College of Cardiology/American Heart Association Task Force and the European Society of Cardiology Committee for Practice Guidelines (writing committee to develop Guidelines for Management of Patients With Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death) . Circulation 2006 ; 114 : e385 – e484 . 16935995 10.1161/CIRCULATIONAHA.106.178233 13. Fishman GI , Chugh SS , Dimarco JP , Sudden cardiac death prediction and prevention: report from a National Heart, Lung, and Blood Institute and Heart Rhythm Society Workshop . Circulation 2010 ; 122 : 2335 – 2348 . 21147730 10.1161/CIRCULATIONAHA.110.976092 PMC3016224 14. Risgaard B , Lynge TH , Wissenberg M , Risk factors and causes of sudden noncardiac death: a nationwide cohort study in Denmark . Heart Rhythm 2015 ; 12 : 968 – 974 . 25614248 10.1016/j.hrthm.2015.01.024 15. Winkel BG , Holst AG , Theilade J , Nationwide study of sudden cardiac death in persons aged 1–35 years . Eur Heart J 2011 ; 32 : 983 – 990 . 21131293 10.1093/eurheartj/ehq428 16. Sanchez O , Campuzano O , Fernandez-Falgueras A , Natural and undetermined sudden death: value of post-mortem genetic investigation . PLoS One 2016 ; 11 : e0167358 . 27930701 10.1371/journal.pone.0167358 PMC5145162 17. Nichol G , Thomas E , Callaway CW , Regional variation in out-of-hospital cardiac arrest incidence and outcome . JAMA 2008 ; 300 : 1423 – 1431 . 18812533 10.1001/jama.300.12.1423 PMC3187919 18. Chugh SS , Jui J , Gunson K , Current burden of sudden cardiac death: multiple source surveillance versus retrospective death certificate-based review in a large U.S. community . J Am Coll Cardiol 2004 ; 44 : 1268 – 1275 . 15364331 10.1016/j.jacc.2004.06.029 19. de Vreede-Swagemakers JJ , Gorgels AP , Dubois-Arbouw WI , Out-of-hospital cardiac arrest in the 1990's: a population-based study in the Maastricht area on incidence, characteristics and survival . J Am Coll Cardiol 1997 ; 30 : 1500 – 1505 . 9362408 10.1016/s0735-1097(97)00355-0 20. Stecker EC , Reinier K , Marijon E , Public

health burden of sudden cardiac death in the United States . *Circ Arrhythm Electrophysiol* 2014 ; 7 : 212 – 217 . 24610738 10.1161/CIRCEP.113.001034 PMC4041478 21. Wong CX , Brown A , Lau DH , Epidemiology of sudden cardiac death: global and regional perspectives . *Heart Lung Circ* 2019 ; 28 : 6 – 14 . 30482683 10.1016/j.hlc.2018.08.026 22. Kong MH , Fonarow GC , Peterson ED , Systematic review of the incidence of sudden cardiac death in the United States . *J Am Coll Cardiol* 2011 ; 57 : 794 – 801 . 21310315 10.1016/j.jacc.2010.09.064 PMC3612019 23. Kitamura T , Iwami T , Kawamura T , Nationwide improvements in survival from out-of-hospital cardiac arrest in Japan . *Circulation* 2012 ; 126 : 2834 – 2843 . 23035209 10.1161/CIRCULATIONAHA.112.109496 24. Gillum RF . Sudden coronary death in the United States: 1980–1985 . *Circulation* 1989 ; 79 : 756 – 765 . 2924409 10.1161/01.cir.79.4.756 25. Bagnall RD , Weintraub RG , Ingles J , A prospective study of sudden cardiac death among children and young adults . *N Engl J Med* 2016 ; 374 : 2441 – 2452 . 27332903 10.1056/NEJMoa1510687 26. Wisten A , Krantz P , Stattin EL . Sudden cardiac death among the young in Sweden from 2000 to 2010: an autopsy-based study . *Europace* 2017 ; 19 : 1327 – 1334 . 28873959 10.1093/europace/euw249 27. Risgaard B , Winkel BG , Jabbari R , Burden of sudden cardiac death in persons aged 1 to 49 years: nationwide study in Denmark . *Circ Arrhythm Electrophysiol* 2014 ; 7 : 205 – 211 . 24604905 10.1161/CIRCEP.113.001421 28. Byrne R , Constant O , Smyth Y , Multiple source surveillance incidence and aetiology of out-of-hospital sudden cardiac death in a rural population in the West of Ireland . *Eur Heart J* 2008 ; 29 : 1418 – 1423 . 18424446 10.1093/eurheartj/ehn155 29. Winkel BG , Risgaard B , Sadjadieh G , Bundgaard H , Haunso S , Tfelt-Hansen J . Sudden cardiac death in children (1–18 years): symptoms and causes of death in a nationwide setting . *Eur Heart J* 2014 ; 35 : 868 – 875 . 24344190 10.1093/eurheartj/ehu509 30. Burns KM , Cottengim C , Dykstra H , Epidemiology of sudden death in a population-based study of infants and children . *J Pediatr: X* 2020 ; 2 : 100023 . 32743542 10.1016/j.ympdx.2020.100023 PMC7394394 31. Eckart RE , Shry EA , Burke AP , Sudden death in young adults: an autopsy-based series of a population undergoing active surveillance . *J Am Coll Cardiol* 2011 ; 58 : 1254 – 1261 . 21903060 10.1016/j.jacc.2011.01.049 32. Thiene G , Corrado D , Basso C . The Northeast Italy, Veneto Region experience . In: Thiene G , Corrado D , Basso C , eds. *Sudden Cardiac Death in the Young and Athletes* . Springer-Verlag Milan ; 2016 . p. 171 – 181 . 33. Chappex N , Schlaepfer J , Fellmann F , Bhuiyan ZA , Wilhelm M , Michaud K . Sudden cardiac death among general population and sport related population in forensic experience . *J Forensic Leg Med* 2015 ; 35 : 62 – 68 . 26344462 10.1016/j.jflm.2015.07.004 34. Deo R , Albert CM . Epidemiology and genetics of sudden cardiac death . *Circulation* 2012 ; 125 : 620 – 637 . 22294707 10.1161/CIRCULATIONAHA.111.023838 PMC3399522 35. Becker LB , Han BH , Meyer PM , Racial differences in the incidence of cardiac arrest and subsequent survival: the CPR Chicago Project . *N Engl J Med* 1993 ; 329 : 600 – 606 . 8341333 10.1056/NEJM199308263290902 36. Zhao D , Post WS , Blasco-Colmenares E , Racial differences in sudden cardiac death . *Circulation* 2019 ; 139 : 1688 – 1697 . 30712378 10.1161/CIRCULATIONAHA.118.036553 PMC6443438 37. Behr ER , Casey A , Sheppard M , Sudden arrhythmic death syndrome: a national survey of sudden unexplained cardiac death . *Heart* 2007 ; 93 : 601 – 605 . 17237131 10.1136/hrt.2006.099598 PMC1955564 38. Griffis H , Wu L , Naim MY , Characteristics and outcomes of AED use in pediatric cardiac arrest in public settings: the influence of neighborhood characteristics . *Resuscitation* 2020 ; 146 : 126 – 131 . 31785372 10.1016/j.resuscitation.2019.09.038 39. Starks MA , Schmicker RH , Peterson ED , Association of neighborhood demographics with out-of-hospital cardiac arrest treatment and outcomes: where you live may matter . *JAMA Cardiol* 2017 ; 2 : 1110 – 1118 . 28854308 10.1001/jamacardio.2017.2671 PMC5710447 40. Basso C , Calabrese F , Corrado D , Thiene G . Postmortem diagnosis in sudden cardiac death victims: macroscopic, microscopic and molecular findings . *Cardiovasc Res* 2001 ; 50 : 290 – 300 . 11334833 10.1016/S0008-6363(01)00261-9 PMC7202452 41. Sasson C , Rogers MA , Dahl J , Kellermann AL . Predictors of survival from out-of-hospital cardiac arrest: a systematic review and meta-analysis . *Circ Cardiovasc Qual Outcomes* 2010 ; 3 : 63 – 81 . 20123673 10.1161/CIRCOUTCOMES.109.889576 42. Berdowski J , Berg RA , Tijssen JG , Koster RW . Global incidences of out-of-hospital cardiac arrest and survival rates: systematic review of 67 prospective studies . *Resuscitation* 2010 ; 81 : 1479 – 1487 . 20828914 10.1016/j.resuscitation.2010.08.006 43. Otsuki S , Aiba T , Tahara Y , Intra-day change in occurrence of out-of-hospital ventricular fibrillation in Japan: the JCS-ReSS study . *Int J Cardiol* 2020 ; 318 : 54 – 60 . 32569698 10.1016/j.ijcard.2020.06.014 44. Zive D , Koprowicz K , Schmidt T

, Variation in out-of-hospital cardiac arrest resuscitation and transport practices in the Resuscitation Outcomes Consortium: ROC Epistery-Cardiac Arrest . Resuscitation 2011 ; 82 : 277 – 284 . 21159416 10.1016/j.resuscitation.2010.10.022 PMC3039085 45. Girotra S , van Diepen S , Nallamothu BK , Regional variation in out-of-hospital cardiac arrest survival in the United States . Circulation 2016 ; 133 : 2159 – 2168 . 27081119 10.1161/CIRCULATIONAHA.115.018175 PMC4889467 46. Atwood C , Eisenberg MS , Herlitz J , Rea TD . Incidence of EMS-treated out-of-hospital cardiac arrest in Europe . Resuscitation 2005 ; 67 : 75 – 80 . 16199289 10.1016/j.resuscitation.2005.03.021 47. Hawkes C , Booth S , Ji C , Epidemiology and outcomes from out-of-hospital cardiac arrests in England . Resuscitation 2017 ; 110 : 133 – 140 . 27865775 10.1016/j.resuscitation.2016.10.030 48. Rea TD , Eisenberg MS , Sinibaldi G , White RD . Incidence of EMS-treated out-of-hospital cardiac arrest in the United States . Resuscitation 2004 ; 63 : 17 – 24 . 15451582 10.1016/j.resuscitation.2004.03.025 49. Becker L , Gold LS , Eisenberg M , White L , Hearne T , Rea T . Ventricular fibrillation in King County, Washington: a 30-year perspective . Resuscitation 2008 ; 79 : 22 – 27 . 18687513 10.1016/j.resuscitation.2008.06.019 50. Bunch TJ , White RD , Gersh BJ , Long-term outcomes of out-of-hospital cardiac arrest after successful early defibrillation . N Engl J Med 2003 ; 348 : 2626 – 2633 . 12826637 10.1056/NEJMoa023053 51. Cummins RO , Ornato JP , Thies WH , Pepe PE . Improving survival from sudden cardiac arrest: the “chain of survival” concept. A statement for health professionals from the Advanced Cardiac Life Support Subcommittee and the Emergency Cardiac Care Committee, American Heart Association . Circulation 1991 ; 83 : 1832 – 1847 . 2022039 10.1161/01.cir.83.5.1832 52. Iwami T , Nichol G , Hiraide A , Continuous improvements in “chain of survival” increased survival after out-of-hospital cardiac arrests: a large-scale population-based study . Circulation 2009 ; 119 : 728 – 734 . 19171854 10.1161/CIRCULATIONAHA.108.802058 53. Lund-Kordahl I , Olasveengen TM , Lorentz T , Samdal M , Wik L , Sunde K . Improving outcome after out-of-hospital cardiac arrest by strengthening weak links of the local Chain of Survival; quality of advanced life support and post-resuscitation care . Resuscitation 2010 ; 81 : 422 – 426 . 20122786 10.1016/j.resuscitation.2009.12.020 54. Wissenberg M , Lippert FK , Folke F , Association of national initiatives to improve cardiac arrest management with rates of bystander intervention and patient survival after out-of-hospital cardiac arrest . JAMA 2013 ; 310 : 1377 – 1384 . 24084923 10.1001/jama.2013.278483 55. Moriwaki Y , Tahara Y , Kosuge T , Suzuki N . Etiology of out-of-hospital cardiac arrest diagnosed via detailed examinations including perimortem computed tomography . J Emerg Trauma Shock 2013 ; 6 : 87 – 94 . 23723616 10.4103/0974-2700.110752 PMC3665077 56. Engdahl J , Holmberg M , Karlson BW , Luepker R , Herlitz J . The epidemiology of out-of-hospital ‘sudden’ cardiac arrest . Resuscitation 2002 ; 52 : 235 – 245 . 11886728 10.1016/s0300-9572(01)00464-6 57. D'Silva A , Sharma S . Management of young competitive athletes with cardiovascular conditions . Heart 2017 ; 103 : 463 – 473 . 27888210 10.1136/heartjnl-2016-309435 58. Asatryan B , Vital C , Kellerhals C , Sports-related sudden cardiac deaths in the young population of Switzerland . PLoS One 2017 ; 12 : e0174434 . 28350812 10.1371/journal.pone.0174434 PMC5370100 59. Landry CH , Allan KS , Connelly KA , Cunningham K , Morrison LJ , Dorian P . Sudden cardiac arrest during participation in competitive sports . N Engl J Med 2017 ; 377 : 1943 – 1953 . 29141175 10.1056/NEJMoa1615710 PMC5726886 60. McNally B , Robb R , Mehta M , Out-of-hospital cardiac arrest surveillance: Cardiac Arrest Registry to Enhance Survival (CARES), United States, October 1, 2005-December 31, 2010 . MMWR Surveill Summ 2011 ; 60 : 1 – 19 . 21796098 61. Hallstrom AP , Ornato JP , Weisfeldt M , Public-access defibrillation and survival after out-of-hospital cardiac arrest . N Engl J Med 2004 ; 351 : 637 – 646 . 15306665 10.1056/NEJMoa040566 62. Capucci A , Aschieri D , Piepoli MF , Bardy GH , Iacono E , Arvedi M . Tripling survival from sudden cardiac arrest via early defibrillation without traditional education in cardiopulmonary resuscitation . Circulation 2002 ; 106 : 1065 – 1070 . 12196330 10.1161/01.cir.0000028148.62305.69 63. Blom MT , Beesems SG , Homma PC , Improved survival after out-of-hospital cardiac arrest and use of automated external defibrillators . Circulation 2014 ; 130 : 1868 – 1875 . 25399395 10.1161/CIRCULATIONAHA.114.010905 64. Myat A , Song KJ , Rea T . Out-of-hospital cardiac arrest: current concepts . Lancet 2018 ; 391 : 970 – 979 . 29536861 10.1016/S0140-6736(18)30472-0 65. Skinner JR , Duflou JA , Semsarian C . Reducing sudden death in young people in Australia and New Zealand: the TRAGADY initiative . Med J Aust 2008 ; 189 : 539 – 540 . 19012547 10.5694/j.1326-5377.2008.tb02173.x 66. Ahmad F , McNally EM , Ackerman MJ , Establishment of specialized clinical cardiovascular genetics programs: recognizing

the need and meeting standards: a scientific statement from the American Heart Association . *Circ Genom Precis Med* 2019 ; 12 : e000054 . 31117808 10.1161/HCG.0000000000000054 67. Basso C , Aguilera B , Banner J , Guidelines for autopsy investigation of sudden cardiac death: 2017 update from the Association for European Cardiovascular Pathology . *Virchows Arch* 2017 ; 471 : 691 – 705 . 28889247 10.1007/s00428-017-2221-0 PMC5711979 68. Nancarrow SA , Booth A , Ariss S , Smith T , Enderby P , Roots A . Ten principles of good interdisciplinary team work . *Hum Resour Health* 2013 ; 11 : 19 . 23663329 10.1186/1478-4491-11-19 PMC3662612 69. Earle NJ , Crawford J , Hayes I , Development of a cardiac inherited disease service and clinical registry: a 15-year perspective . *Am Heart J* 2019 ; 209 : 126 – 130 . 30686478 10.1016/j.ahj.2018.11.013 70. Lahrouchi N , Raju H , Lodder EM , Utility of post-mortem genetic testing in cases of sudden arrhythmic death syndrome . *J Am Coll Cardiol* 2017 ; 69 : 2134 – 2145 . 28449774 10.1016/j.jacc.2017.02.046 PMC5405216 71. Marcondes L , Crawford J , Earle N , Long QT molecular autopsy in sudden unexplained death in the young (1–40 years old): lessons learnt from an eight year experience in New Zealand . *PLoS One* 2018 ; 13 : e0196078 . 29672598 10.1371/journal.pone.0196078 PMC5909669 72. Yeates L , Hunt L , Saleh M , Semsarian C , Ingles J . Poor psychological wellbeing particularly in mothers following sudden cardiac death in the young . *Eur J Cardiovasc Nurs* 2013 ; 12 : 484 – 491 . 23568895 10.1177/1474515113485510 73. van der Werf C , Onderwater AT , van Langen IM , Smets EM . Experiences, considerations and emotions relating to cardiogenetic evaluation in relatives of young sudden cardiac death victims . *Eur J Hum Genet* 2014 ; 22 : 192 – 196 . 23736216 10.1038/ejhg.2013.126 PMC3895645 74. Caleshu C , Kasparian NA , Edwards KS , Interdisciplinary psychosocial care for families with inherited cardiovascular diseases . *Trends Cardiovasc Med* 2016 ; 26 : 647 – 653 . 27256036 10.1016/j.tcm.2016.04.010 75. Rhodes A , Rosman L , Cahill J , Minding the genes: a multidisciplinary approach towards genetic assessment of cardiovascular disease . *J Genet Couns* 2017 ; 26 : 224 – 231 . 27695998 10.1007/s10897-016-0017-z 76. Waddell-Smith K , Donoghue T , Graham M , The inpatient cardiology visit: missing the opportunity to detect inherited heart conditions . *Heart Lung Circ* 2014 ; 23 ( Suppl 2 ) : e18 – e19 . 77. Waddell-Smith KE , Donoghue T , Oates S , Inpatient detection of cardiac-inherited disease: the impact of improving family history taking . *Open Heart* 2016 ; 3 : e000329 . 26925241 10.1136/openhrt-2015-000329 PMC4762189 78. Earle N , Crawford J , Gibson K , Detection of sudden death syndromes in New Zealand . *N Z Med J* 2016 ; 129 : 67 – 74 . 27857240 79. Ingles J , Lind JM , Phongsavan P , Semsarian C . Psychosocial impact of specialized cardiac genetic clinics for hypertrophic cardiomyopathy . *Genet Med* 2008 ; 10 : 117 – 120 . 18281919 10.1097/GIM.0b013e3181612cc7 80. Tan HL , Hofman N , van Langen IM , van der Wal AC , Wilde AA . Sudden unexplained death: heritability and diagnostic yield of cardiological and genetic examination in surviving relatives . *Circulation* 2005 ; 112 : 207 – 213 . 15998675 10.1161/CIRCULATIONAHA.104.522581 81. Behr E , Wood DA , Wright M , Cardiological assessment of first-degree relatives in sudden arrhythmic death syndrome . *Lancet* 2003 ; 362 : 1457 – 1459 . 14602442 10.1016/s0140-6736(03)14692-2 82. Kumar S , Peters S , Thompson T , Familial cardiological and targeted genetic evaluation: low yield in sudden unexplained death and high yield in unexplained cardiac arrest syndromes . *Heart Rhythm* 2013 ; 10 : 1653 – 1660 . 23973953 10.1016/j.hrthm.2013.08.022 83. Giudicessi JR , Lieve KVV , Rohatgi RK , Assessment and validation of a phenotype-enhanced variant classification framework to promote or demote RYR2 missense variants of uncertain significance . *Circ Genom Precis Med* 2019 ; 12 : e002510 . 31112425 10.1161/CIRCGEN.119.002510 84. Mellor G , Laksman ZWM , Tadros R , Genetic testing in the evaluation of unexplained cardiac arrest . *Circ Cardiovasc Genet* 2017 ; 10 : e001686 . 28600387 10.1161/CIRCGENETICS.116.001686 85. van der Werf C , Hofman N , Tan HL , Diagnostic yield in sudden unexplained death and aborted cardiac arrest in the young: the experience of a tertiary referral center in the Netherlands . *Heart Rhythm* 2010 ; 7 : 1383 – 1389 . 20646679 10.1016/j.hrthm.2010.05.036 86. Wilms HR , Midgley DJ , Morrow P , Stables S , Crawford J , Skinner JR . Evaluation of autopsy and police reports in the investigation of sudden unexplained death in the young . *Forensic Sci Med Pathol* 2012 ; 8 : 380 – 389 . 22615068 10.1007/s12024-012-9340-3 87. Biesecker B . Goals of genetic counselling . *Clinical Genetics* 2001 ; 60 : 323 – 330 . 11903329 10.1034/j.1399-0004.2001.600501.x 88. Resta R , Biesecker BB , Bennett RL , A new definition of Genetic Counseling: National Society of Genetic Counselors' Task Force report . *J Genet Couns* 2006 ; 15 : 77 – 83 . 16761103 10.1007/s10897-005-9014-3 89. Charron P , Arad M , Arbustini E , Genetic counselling and testing in cardiomyopathies: a position statement of the European Society of Cardiology Working Group on Myocardial and Pericardial

Diseases . Eur Heart J 2010 ; 31 : 2715 – 2726 . 20823110 10.1093/eurheartj/ehq271 90. Abacan M , Alsubaie L , Barlow-Stewart K , The global state of the genetic counseling profession . Eur J Hum Genet 2019 ; 27 : 183 – 197 . 30291341 10.1038/s41431-018-0252-x PMC6336871 91. Laurino MY , Leppig KA , Abad PJ , A report on ten Asia Pacific countries on current status and future directions of the genetic counseling profession: the establishment of the Professional Society of Genetic Counselors in Asia. J Genet Couns 2018 ; 27 : 21 – 32 . 28699126 10.1007/s10897-017-0115-6 92. Ingles J , Yeates L , Semsarian C . The emerging role of the cardiac genetic counselor . Heart Rhythm 2011 ; 8 : 1958 – 1962 . 21767518 10.1016/j.hrthm.2011.07.017 93. Hershberger RE , Givertz M , Ho CY , Genetic evaluation of cardiomyopathy: a Heart Failure Society of America Practice Guideline . J Card Fail 2018 ; 24 : 281 – 302 . 29567486 10.1016/j.cardfail.2018.03.004 PMC9903357 94. Skrzynia C , Demo EM , Baxter SM . Genetic counseling and testing for hypertrophic cardiomyopathy: an adult perspective . J Cardiovasc Transl Res 2009 ; 2 : 493 – 499 . 20560007 10.1007/s12265-009-9127-4 95. Ingles J . Psychological issues in managing families with inherited cardiovascular diseases . Cold Spring Harb Perspect Med 2019 ; 9 : 23 : a036558 . 10.1101/cshperspect.a036558 PMC7461768 31548222 96. Aatre RD , Day SM . Psychological issues in genetic testing for inherited cardiovascular diseases . Circ Cardiovasc Genet 2011 ; 4 : 81 – 90 . 21325164 10.1161/CIRCGENETICS.110.957365 97. Sturm AC . Genetic testing in the contemporary diagnosis of cardiomyopathy . Curr Heart Fail Rep 2013 ; 10 : 63 – 72 . 23135967 10.1007/s11897-012-0124-6 98. Burns C , McGaughan J , Davis A , Semsarian C , Ingles J . Factors influencing uptake of familial long QT syndrome genetic testing . Am J Med Genet A 2016 ; 170A : 418 – 425 . 26544151 10.1002/ajmg.a.37455 99. Ingles J , James C . Psychosocial care and cardiac genetic counseling following sudden cardiac death in the young . Prog Pediatr Cardiol 2017 ; 45 : 31 – 36 . 100. Wong EK , Bartels K , Hathaway J , Perceptions of genetic variant reclassification in patients with inherited cardiac disease . Eur J Hum Genet 2019 ; 27 : 1134 – 1142 . 30903112 10.1038/s41431-019-0377-6 PMC6777462 101. Bates K , Sweeting J , Yeates L , McDonald K , Semsarian C , Ingles J . Psychological adaptation to molecular autopsy findings following sudden cardiac death in the young . Genet Med 2019 ; 21 : 1452 – 1456 . 30327538 10.1038/s41436-018-0338-4 102. Meiser B , Irlé J , Lobb E , Barlow-Stewart K . Assessment of the content and process of genetic counseling: a critical review of empirical studies . J Genet Couns 2008 ; 17 : 434 – 451 . 18791813 10.1007/s10897-008-9173-0 103. Edwards A , Gray J , Clarke A , Interventions to improve risk communication in clinical genetics: systematic review . Patient Educ Couns 2008 ; 71 : 4 – 25 . 18207694 10.1016/j.pec.2007.11.026 104. Austin J , Semaka A , Hadjipavlou G . Conceptualizing genetic counseling as psychotherapy in the era of genomic medicine . J Genet Couns 2014 ; 23 : 903 – 909 . 24841456 10.1007/s10897-014-9728-1 PMC4318692 105. Accreditation Council for Genetic Counseling . Practice based competencies for genetic counselors . Available from [https://www.gceducation.org/wp-content/uploads/2019/02/ACG-C-Core-Competencies-Brochure\\_15\\_Web.pdf](https://www.gceducation.org/wp-content/uploads/2019/02/ACG-C-Core-Competencies-Brochure_15_Web.pdf) . Accessed May 5, 2020 . 10.1007/s10897-016-9984-3 27333894 106. Michie S , Marteau TM , Bobrow M . Genetic counselling: the psychological impact of meeting patients' expectations . J Med Genet 1997 ; 34 : 237 – 241 . 9132497 10.1136/jmg.34.3.237 PMC1050900 107. Ison HE , Ware SM , Schwantes-An TH , Freeze S , Elmore L , Spoonamore KG . The impact of cardiovascular genetic counseling on patient empowerment . J Genet Couns 2019 ; 28 : 570 – 577 . 30680842 10.1002/jgc4.1050 108. Burns C , James C , Ingles J . Communication of genetic information to families with inherited rhythm disorders . Heart Rhythm 2018 ; 15 : 780 – 786 . 29175646 10.1016/j.hrthm.2017.11.024 109. Liu G , MacLeod H , Webster G , McNally EM , O'Neill SM , Dellefave-Castillo L . Genetic counselors' approach to postmortem genetic testing after sudden death: an exploratory study . Acad Forensic Pathol 2018 ; 8 : 738 – 751 . 31240068 10.1177/1925362118797748 PMC6490591 110. Burns C , Yeates L , Spinks C , Semsarian C , Ingles J . Attitudes, knowledge and consequences of uncertain genetic findings in hypertrophic cardiomyopathy . Eur J Hum Genet 2017 ; 25 : 809 – 815 . 28594412 10.1038/ejhg.2017.66 PMC5520078 111. Ingles J , Semsarian C . Conveying a probabilistic genetic test result to families with an inherited heart disease . Heart Rhythm 2014 ; 11 : 1073 – 1078 . 24632221 10.1016/j.hrthm.2014.03.017 112. Reuter C , Grove ME , Orland K , Spoonamore K , Caleshu C . Clinical cardiovascular genetic counselors take a leading role in team-based variant classification . J Genet Couns 2018 ; 27 : 751 – 760 . 29234989 10.1007/s10897-017-0175-7 113. Bland A , Harrington EA , Dunn K , Clinically impactful differences in variant interpretation between clinicians and testing laboratories: a single-center

experience . Genet Med 2018 ; 20 : 369 – 373 . 29240077 10.1038/gim.2017.212 114. Furqan A , Arscott P , Girolami F , Care in specialized centers and data sharing increase agreement in hypertrophic cardiomyopathy genetic test interpretation . Circ Cardiovasc Genet 2017 ; 10 : e001700 . 28986452 10.1161/CIRCGENETICS.116.001700 115. Das KJ , Ingles J , Bagnall RD , Semsarian C . Determining pathogenicity of genetic variants in hypertrophic cardiomyopathy: importance of periodic reassessment . Genet Med 2014 ; 16 : 286 – 293 . 24113344 10.1038/gim.2013.138 116. Dunn KE , Caleshu C , Cirino AL , Ho CY , Ashley EA . A clinical approach to inherited hypertrophy: the use of family history in diagnosis, risk assessment, and management . Circ Cardiovasc Genet 2013 ; 6 : 118 – 131 . 23424256 10.1161/CIRCGENETICS.110.959387 PMC3898734 117. Shear MK . Getting straight about grief . Depress Anxiety 2012 ; 29 : 461 – 464 . 22730310 10.1002/da.21963 118. Simon NM , Shear KM , Thompson EH . The prevalence and correlates of psychiatric comorbidity in individuals with complicated grief . Compr Psychiatry 2007 ; 48 : 395 – 399 . 17707245 10.1016/j.comppsy.2007.05.002 119. American Psychiatric Association . Diagnostic and Statistical Manual of Mental Disorders (DSM-5) , Fifth ed. Washington, DC : American Psychiatric Association Publishing ; 2013 . 120. Simon NM . Treating complicated grief . JAMA 2013 ; 310 : 416 – 423 . 23917292 10.1001/jama.2013.8614 PMC4530627 121. Ingles J , Spinks C , Yeates L , McGeechan K , Kasparian N , Semsarian C . Post-traumatic stress and prolonged grief after the sudden cardiac death of a young relative . JAMA Intern Med 2016 ; 176 : 402 – 405 . 26809585 10.1001/jamainternmed.2015.7808 122. Shalev A , Liberzon I , Marmar C . Post-traumatic stress disorder . N Engl J Med 2017 ; 376 : 2459 – 2469 . 28636846 10.1056/NEJMra1612499 123. Latham AE , Prigerson HG . Suicidality and bereavement: complicated grief as psychiatric disorder presenting greatest risk for suicidality . Suicide Life Threat Behav 2004 ; 34 : 350 – 362 . 15585457 10.1521/suli.34.4.350.53737 PMC1459278 124. Prigerson HG , Bierhals AJ , Kasl SV . Traumatic grief as a risk factor for mental and physical morbidity . Am J Psychiatry 1997 ; 154 : 616 – 623 . 9137115 10.1176/ajp.154.5.616 125. Karam N , Jabre P , Narayanan K , Psychological support and medical screening of first-degree relatives of sudden cardiac arrest victims . JACC Clin Electrophysiol 2020 ; 6 : 586 – 587 . 32439046 10.1016/j.jacep.2020.02.002 126. McDonald K , Sharpe L , Yeates L , Semsarian C , Ingles J . A needs analysis of parents following sudden cardiac death in the young . Open Heart 2020 ; 7 : e001120 . 32709698 10.1136/openhrt-2019-001120 PMC7380729 127. Wisten A , Zingmark K . Supportive needs of parents confronted with sudden cardiac death: a qualitative study . Resuscitation 2007 ; 74 : 68 – 74 . 17353083 10.1016/j.resuscitation.2006.11.014 128. Dent A , Condon L , Blair P , Fleming P . A study of bereavement care after a sudden and unexpected death . Arch Dis Child 1996 ; 74 : 522 – 526 . 8758129 10.1136/adc.74.6.522 PMC1511561 129. Merlevede E , Spooren D , Henderick H , Perceptions, needs and mourning reactions of bereaved relatives confronted with a sudden unexpected death . Resuscitation 2004 ; 61 : 341 – 348 . 15172714 10.1016/j.resuscitation.2004.01.024 130. Steffen EM , Timotijevic L , Coyle A . A qualitative analysis of psychosocial needs and support impacts in families affected by young sudden cardiac death: the role of community and peer support [published online ahead of print May 5, 2020] . Eur J Cardiovasc Nurs 2020 ; 10.1177/1474515120922347.1474515120922347 . PMC7817993 32370539 131. Solomon P . Peer support/peer provided services underlying processes, benefits, and critical ingredients . Psychiatr Rehabil J 2004 ; 27 : 392 – 401 . 15222150 10.2975/27.2004.392.401 132. Landers GM , Zhou M . An analysis of relationships among peer support, psychiatric hospitalization, and crisis stabilization . Community Ment Health J 2011 ; 47 : 106 – 112 . 19551502 10.1007/s10597-009-9218-3 133. Bartone PT , Bartone JV , Violanti JM , Gileno ZM . Peer support services for bereaved survivors: a systematic review . Omega (Westport) 2019 ; 80 : 137 – 166 . 28871835 10.1177/0030222817728204 134. Farnsworth MM , Fosyth D , Haglund C , Ackerman MJ . When I go in to wake them ... I wonder: parental perceptions about congenital long QT syndrome . J Am Acad Nurse Pract 2006 ; 18 : 284 – 290 . 16719847 10.1111/j.1745-7599.2006.00132.x 135. Risgaard B , Winkel BG , Jabbari R , Sudden cardiac death: pharmacotherapy and proarrhythmic drugs: a nationwide cohort study in Denmark . JACC Clin Electrophysiol 2017 ; 3 : 473 – 481 . 29759603 10.1016/j.jacep.2016.12.023 136. Michowitz Y , Milman A , Sarquella-Brugada G , Fever-related arrhythmic events in the multicenter Survey on Arrhythmic Events in Brugada Syndrome . Heart Rhythm 2018 ; 15 : 1394 – 1401 . 29649615 10.1016/j.hrthm.2018.04.007 137. Amin AS , Herfst LJ , Delisle BP , Fever-induced QTc prolongation and ventricular arrhythmias in individuals with type 2 congenital long QT syndrome . J

Clin Invest 2008 ; 118 : 2552 – 2561 . 18551196 10.1172/JCI35337 PMC2423868 138. Probst V , Denjoy I , Meregalli PG , Clinical aspects and prognosis of Brugada syndrome in children . Circulation 2007 ; 115 : 2042 – 2048 . 17404158 10.1161/CIRCULATIONAHA.106.664219 139. Glinge C , Jabbari R , Risgaard B , Symptoms before sudden arrhythmic death syndrome: a nationwide study among the young in Denmark . J Cardiovasc Electrophysiol 2015 ; 26 : 761 – 767 . 25807988 10.1111/jce.12674 140. Maruthappu T , Posafalvi A , Castelletti S , Loss-of-function desmoplakin I and II mutations underlie dominant arrhythmogenic cardiomyopathy with a hair and skin phenotype . Br J Dermatol 2019 ; 180 : 1114 – 1122 . 30382575 10.1111/bjd.17388 PMC6318013 141. Lieve KVV , Verhagen JMA , Wei J , Linking the heart and the brain: neurodevelopmental disorders in patients with catecholaminergic polymorphic ventricular tachycardia . Heart Rhythm 2019 ; 16 : 220 – 228 . 30170228 10.1016/j.hrthm.2018.08.025 142. Byers PH , Belmont J , Black J , Diagnosis, natural history, and management in vascular Ehlers-Danlos syndrome . Am J Med Genet C Semin Med Genet 2017 ; 175 : 40 – 47 . 28306228 10.1002/ajmg.c.31553 143. Behr ER , Dalageorgou C , Christiansen M , Sudden arrhythmic death syndrome: familial evaluation identifies inheritable heart disease in the majority of families . Eur Heart J 2008 ; 29 : 1670 – 1680 . 18508782 10.1093/eurheartj/ehn219 144. Steinberg C , Padfield GJ , Champagne J , Cardiac abnormalities in first-degree relatives of unexplained cardiac arrest victims: a report from the Cardiac Arrest Survivors with Preserved Ejection Fraction Registry . Circ Arrhythm Electrophysiol 2016 ; 9 : e004274 . 27635072 10.1161/CIRCEP.115.004274 145. Raju H , Papadakis M , Govindan M , Low prevalence of risk markers in cases of sudden death due to Brugada syndrome relevance to risk stratification in Brugada syndrome . J Am Coll Cardiol 2011 ; 57 : 2340 – 2345 . 21636035 10.1016/j.jacc.2010.11.067 146. Johnson JN , Hofman N , Haglund CM , Cascino GD , Wilde AA , Ackerman MJ . Identification of a possible pathogenic link between congenital long QT syndrome and epilepsy . Neurology 2009 ; 72 : 224 – 231 . 19038855 10.1212/01.wnl.0000335760.02995.ca PMC2677528 147. Chahal CAA , Salloum MN , Alahdab F , Systematic review of the genetics of sudden unexpected death in epilepsy: potential overlap with sudden cardiac death and arrhythmia-related genes . J Am Heart Assoc 2020 ; 9 : e012264 . 10.1161/JAHA.119.012264 PMC6988156 31865891 148. Algra A , Tijssen JG , Roelandt JR , Pool J , Lubsen J . QTc prolongation measured by standard 12-lead electrocardiography is an independent risk factor for sudden death due to cardiac arrest . Circulation 1991 ; 83 : 1888 – 1894 . 2040041 10.1161/01.cir.83.6.1888 149. Brugada J , Brugada R , Brugada P . Right bundle-branch block and ST-segment elevation in leads V1 through V3: a marker for sudden death in patients without demonstrable structural heart disease . Circulation 1998 ; 97 : 457 – 460 . 9490240 10.1161/01.cir.97.5.457 150. Haissaguerre M , Derval N , Sacher F , Sudden cardiac arrest associated with early repolarization . N Engl J Med 2008 ; 358 : 2016 – 2023 . 18463377 10.1056/NEJMoa071968 151. Priori SG , Wilde AA , Horie M , Executive summary: HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes . Heart Rhythm 2013 ; 10 : e85 – e108 . 23916535 10.1016/j.hrthm.2013.07.021 152. Antzelevitch C , Yan GX , Ackerman MJ , J-Wave syndromes expert consensus conference report: emerging concepts and gaps in knowledge . Heart Rhythm 2016 ; 13 : e295 – e324 . 27423412 10.1016/j.hrthm.2016.05.024 PMC5035208 153. Conte G , Belhassen B , Lambiase P , Out-of-hospital cardiac arrest due to idiopathic ventricular fibrillation in patients with normal electrocardiograms: results from a multicentre long-term registry . Europace 2019 ; 21 : 1670 – 1677 . 31504477 10.1093/europace/euz221 PMC6826207 154. Haukilahti MAE , Holmstrom L , Vahatalo J , Sudden cardiac death in women . Circulation 2019 ; 139 : 1012 – 1021 . 30779638 10.1161/CIRCULATIONAHA.118.037702 155. Abdalla IS , Prineas RJ , Neaton JD , Jacobs DR Jr , Crow RS . Relation between ventricular premature complexes and sudden cardiac death in apparently healthy men . Am J Cardiol 1987 ; 60 : 1036 – 1042 . 3673904 10.1016/0002-9149(87)90348-1 156. Ataklte F , Erqou S , Laukkanen J , Kaptoge S . Meta-analysis of ventricular premature complexes and their relation to cardiac mortality in general populations . Am J Cardiol 2013 ; 112 : 1263 – 1270 . 23927786 10.1016/j.amjcard.2013.05.065 157. Noda T , Shimizu W , Taguchi A , Malignant entity of idiopathic ventricular fibrillation and polymorphic ventricular tachycardia initiated by premature extrasystoles originating from the right ventricular outflow tract . J Am Coll Cardiol 2005 ; 46 : 1288 – 1294 . 16198845 10.1016/j.jacc.2005.05.077 158. Aizawa Y , Sato A , Watanabe H , Dynamicity of the J-wave in idiopathic ventricular fibrillation with a special reference to pause-dependent augmentation of the J-wave . J Am Coll Cardiol 2012 ; 59 : 1948 – 1953 . 22624834 10.1016/j.jacc.2012.02.028 159. Kawata H , Noda T , Yamada Y ,

Effect of sodium-channel blockade on early repolarization in inferior/lateral leads in patients with idiopathic ventricular fibrillation and Brugada syndrome . *Heart Rhythm* 2012 ; 9 : 77 – 83 . 21855521 10.1016/j.hrthm.2011.08.017 160. Gandjbakhch E , Redheuil A , Pousset F , Charron P , Frank R . Clinical diagnosis, imaging, and genetics of arrhythmogenic right ventricular cardiomyopathy/dysplasia: JACC state-of-the-art review . *J Am Coll Cardiol* 2018 ; 72 : 784 – 804 . 30092956 10.1016/j.jacc.2018.05.065 161. Manolis TA , Manolis AA , Melita H , Manolis AS . Sudden unexpected death in epilepsy: the neuro-cardio-respiratory connection . *Seizure* 2019 ; 64 : 65 – 73 . 30566897 10.1016/j.seizure.2018.12.007 162. MacCormick JM , McAlister H , Crawford J , Misdiagnosis of long QT syndrome as epilepsy at first presentation . *Ann Emerg Med* 2009 ; 54 : 26 – 32 . 19282063 10.1016/j.annemergmed.2009.01.031 163. Lacour P , Buschmann C , Storm C , Cardiac implantable electronic device interrogation at forensic autopsy: an underestimated resource? *Circulation* 2018 ; 137 : 2730 – 2740 . 29915100 10.1161/CIRCULATIONAHA.117.032367 164. Bayes de Luna A , Coumel P , Leclercq JF . Ambulatory sudden cardiac death: mechanisms of production of fatal arrhythmia on the basis of data from 157 cases . *Am Heart J* 1989 ; 117 : 151 – 159 . 2911968 10.1016/0002-8703(89)90670-4 165. Gladding PA , Evans CA , Crawford J , Posthumous diagnosis of long QT syndrome from neonatal screening cards . *Heart Rhythm* 2010 ; 7 : 481 – 486 . 20167303 10.1016/j.hrthm.2009.12.023 166. Carturan E , Tester DJ , Brost BC , Basso C , Thiene G , Ackerman MJ . Postmortem genetic testing for conventional autopsy-negative sudden unexplained death: an evaluation of different DNA extraction protocols and the feasibility of mutational analysis from archival paraffin-embedded heart tissue . *Am J Clin Pathol* 2008 ; 129 : 391 – 397 . 18285261 10.1309/VLA7TT9EQ05FFVN4 PMC3332542 167. Searles Nielsen S , Mueller BA , De Roos AJ , Checkoway H . Newborn screening archives as a specimen source for epidemiologic studies: feasibility and potential for bias . *Ann Epidemiol* 2008 ; 18 : 58 – 64 . 18063239 10.1016/j.annepidem.2007.06.002 PMC2190083 168. Skinner JR , Chong B , Fawcner M , Webster DR , Hegde M . Use of the newborn screening card to define cause of death in a 12-year-old diagnosed with epilepsy . *J Paediatr Child Health* 2004 ; 40 : 651 – 653 . 15469540 10.1111/j.1440-1754.2004.00498.x 169. Johnson JN , Tester DJ , Bass NE , Ackerman MJ . Cardiac channel molecular autopsy for sudden unexpected death in epilepsy . *J Child Neurol* 2010 ; 25 : 916 – 921 . 20395638 10.1177/0883073809343722 170. Tsuda T , Geary EM , Temple J . Significance of automated external defibrillator in identifying lethal ventricular arrhythmias . *Eur J Pediatr* 2019 ; 178 : 1333 – 1342 . 31297625 10.1007/s00431-019-03421-9 171. Riesinger L , Fichtner S , Schuhmann CG , Postmortem interrogation of cardiac implantable electrical devices may clarify time and cause of death . *Int J Legal Med* 2019 ; 133 : 883 – 888 . 30238160 10.1007/s00414-018-1932-3 172. Sinha SK , Crain B , Flickinger K , Clinical inferences of cardiovascular implantable electronic device analysis at autopsy . *J Am Coll Cardiol* 2016 ; 68 : 1255 – 1264 . 27634115 10.1016/j.jacc.2016.06.052 173. Hansen BL , Jacobsen EM , Kjerrumgaard A , Diagnostic yield in victims of sudden cardiac death and their relatives . *Europace* 2020 ; 22 : 964 – 971 . 32307520 10.1093/europace/euaa056 174. Peterson GF , Clark SC , National Association of Medical Examiners. Forensic autopsy performance standards . *Am J Forensic Med Pathol* 2006 ; 27 : 200 – 225 . 16936503 10.1097/01.paf.0000243580.43150.3c 175. Gulino SP , Burns K , Gunther WM , MacLeod H . Improving forensic pathologic investigation of sudden death in the young: tools, guidance, and methods of cardiovascular dissection from the sudden death in the young case registry . *Academic Forensic Pathology* 2018 ; 8 : 347 – 391 . 31240048 10.1177/1925362118782077 PMC6490127 176. de Noronha SV , Behr ER , Papadakis M , The importance of specialist cardiac histopathological examination in the investigation of young sudden cardiac deaths . *Europace* 2014 ; 16 : 899 – 907 . 24148315 10.1093/europace/eut329 177. Finocchiaro G , Papadakis M , Robertus JL , Etiology of sudden death in sports: insights from a United Kingdom regional registry . *J Am Coll Cardiol* 2016 ; 67 : 2108 – 2115 . 27151341 10.1016/j.jacc.2016.02.062 178. Brinkmann B . Harmonization of medico-legal autopsy rules: Committee of Ministers. Council of Europe . *Int J Legal Med* 1999 ; 113 : 1 – 14 . 10654232 10.1007/s004140050271 179. Blokker BM , Weustink AC , Wagenveld IM , Conventional autopsy versus minimally invasive autopsy with postmortem MRI, CT, and CT-guided biopsy: comparison of diagnostic performance . *Radiology* 2018 ; 289 : 658 – 667 . 30251930 10.1148/radiol.2018180924 180. Bjune T , Risgaard B , Kruckow L , Post-mortem toxicology in young sudden cardiac death victims: a nationwide cohort study . *Europace* 2018 ; 20 : 614 – 621 . 28339816 10.1093/europace/euw435 181. Middleton O , Atherton D , Bundock E , National Association of

Medical Examiners position paper: recommendations for the investigation and certification of deaths in people with epilepsy . *Epilepsia* 2018 ; 59 : 530 – 543 . 29492970 10.1111/epi.14030 PMC6084455 182. Mellor G , Raju H , de Noronha SV , Clinical characteristics and circumstances of death in the sudden arrhythmic death syndrome . *Circ Arrhythm Electrophysiol* 2014 ; 7 : 1078 – 1083 . 25262685 10.1161/CIRCEP.114.001854 183. Sorkin T , Sheppard MN . Sudden unexplained death in alcohol misuse (SUDAM) patients have different characteristics to those who died from sudden arrhythmic death syndrome (SADS) . *Forensic Sci Med Pathol* 2017 ; 13 : 278 – 283 . 28668989 10.1007/s12024-017-9877-2 PMC5554285 184. Papadakis M , Raju H , Behr ER , Sudden cardiac death with autopsy findings of uncertain significance: potential for erroneous interpretation . *Circ Arrhythm Electrophysiol* 2013 ; 6 : 588 – 596 . 23671135 10.1161/CIRCEP.113.000111 185. Tester DJ , Medeiros-Domingo A , Will ML , Haglund CM , Ackerman MJ . Cardiac channel molecular autopsy: insights from 173 consecutive cases of autopsy-negative sudden unexplained death referred for postmortem genetic testing . *Mayo Clin Proc* 2012 ; 87 : 524 – 539 . 22677073 10.1016/j.mayocp.2012.02.017 PMC3498431 186. Gollob MH , Jones DL , Krahm AD , Somatic mutations in the connexin 40 gene (GJA5) in atrial fibrillation . *N Engl J Med* 2006 ; 354 : 2677 – 2688 . 16790700 10.1056/NEJMoa052800 187. Leong IUS , Dryland PA , Prosser DO , Splice site variants in the KCNQ1 and SCN5A genes: transcript analysis as a tool in supporting pathogenicity . *J Clin Med Res* 2017 ; 9 : 709 – 718 . 28725320 10.14740/jocmr2894w PMC5505308 188. Hofman N , Tan HL , Alders M , Yield of molecular and clinical testing for arrhythmia syndromes: report of 15 years' experience . *Circulation* 2013 ; 128 : 1513 – 1521 . 23963746 10.1161/CIRCULATIONAHA.112.000091 189. Papadakis M , Papatheodorou E , Mellor G , The diagnostic yield of Brugada syndrome after sudden death with normal autopsy . *J Am Coll Cardiol* 2018 ; 71 : 1204 – 1214 . 29544603 10.1016/j.jacc.2018.01.031 190. Elger BS , Michaud K , Fellmann F , Mangin P . Sudden death: ethical and legal problems of post-mortem forensic genetic testing for hereditary cardiac diseases . *Clin Genet* 2010 ; 77 : 287 – 292 . 19863545 10.1111/j.1399-0004.2009.01293.x 191. Michaud K , Fellmann F , Abriel H , Beckmann JS , Mangin P , Elger BS . Molecular autopsy in sudden cardiac death and its implication for families: discussion of the practical, legal and ethical aspects of the multidisciplinary collaboration . *Swiss Med Wkly* 2009 ; 139 : 712 – 718 . 20047134 10.4414/smw.2009.12837 192. Lahrouchi N , Raju H , Lodder EM , The yield of postmortem genetic testing in sudden death cases with structural findings at autopsy . *Eur J Hum Genet* 2020 ; 28 : 17 – 22 . 31534214 10.1038/s41431-019-0500-8 PMC6906523 193. Hofman N , Tan HL , Alders M , van Langen IM , Wilde AA . Active cascade screening in primary inherited arrhythmia syndromes: does it lead to prophylactic treatment? *J Am Coll Cardiol* 2010 ; 55 : 2570 – 2576 . 20513597 10.1016/j.jacc.2009.12.063 194. Manrai AK , Funke BH , Rehm HL , Genetic misdiagnoses and the potential for health disparities . *N Engl J Med* 2016 ; 375 : 655 – 665 . 27532831 10.1056/NEJMsa1507092 PMC5292722 195. Hosseini SM , Kim R , Udupa S , Reappraisal of reported genes for sudden arrhythmic death . *Circulation* 2018 ; 138 : 1195 – 1205 . 29959160 10.1161/CIRCULATIONAHA.118.035070 PMC6147087 196. Adler A , Novelli V , Amin AS , An international, multicentered, evidence-based reappraisal of genes reported to cause congenital long QT syndrome . *Circulation* 2020 ; 141 : 418 – 428 . 31983240 10.1161/CIRCULATIONAHA.119.043132 PMC7017940 197. Ackerman MJ , Tester DJ , Jones GS , Will ML , Burrow CR , Curran ME . Ethnic differences in cardiac potassium channel variants: implications for genetic susceptibility to sudden cardiac death and genetic testing for congenital long QT syndrome . *Mayo Clin Proc* 2003 ; 78 : 1479 – 1487 . 14661677 10.4065/78.12.1479 198. Ackerman MJ , Splawski I , Makielski JC , Spectrum and prevalence of cardiac sodium channel variants among black, white, Asian, and Hispanic individuals: implications for arrhythmogenic susceptibility and Brugada/long QT syndrome genetic testing . *Heart Rhythm* 2004 ; 1 : 600 – 607 . 15851227 10.1016/j.hrthm.2004.07.013 199. Kapa S , Tester DJ , Salisbury BA , Genetic testing for long-QT syndrome: distinguishing pathogenic mutations from benign variants . *Circulation* 2009 ; 120 : 1752 – 1760 . 19841300 10.1161/CIRCULATIONAHA.109.863076 PMC3025752 200. Giudicessi JR , Roden DM , Wilde AAM , Ackerman MJ . Classification and reporting of potentially proarrhythmic common genetic variation in long QT syndrome genetic testing . *Circulation* 2018 ; 137 : 619 – 630 . 29431662 10.1161/CIRCULATIONAHA.117.030142 PMC6383807 201. Giudicessi JR , Wilde AAM , Ackerman MJ . The genetic architecture of long QT syndrome: a critical reappraisal . *Trends Cardiovasc Med* 2018 ; 28 : 453 – 464 . 29661707 10.1016/j.tcm.2018.03.003 PMC6590899 202. Ingles J , Goldstein J , Thaxton C , Evaluating the

clinical validity of hypertrophic cardiomyopathy genes . *Circ Genom Precis Med* 2019 ; 12 : e002460 . 30681346 10.1161/CIRCGEN.119.002460 PMC6410971 203. Ackerman JP , Bartos DC , Kapplinger JD , Tester DJ , Delisle BP , Ackerman MJ . The promise and peril of precision medicine: phenotyping still matters most . *Mayo Clin Proc* 2016 ; 91 : 1606 – 1616 . 10.1016/j.mayocp.2016.08.008 PMC6365209 27810088 204. Carrieri D , Howard HC , Benjamin C , Recontacting patients in clinical genetics services: recommendations of the European Society of Human Genetics . *Eur J Hum Genet* 2019 ; 27 : 169 – 182 . 30310124 10.1038/s41431-018-0285-1 PMC6336881 205. van der Werf C , Stiekema L , Tan HL , Low rate of cardiac events in first-degree relatives of diagnosis-negative young sudden unexplained death syndrome victims during follow-up . *Heart Rhythm* 2014 ; 11 : 1728 – 1732 . 24882506 10.1016/j.hrthm.2014.05.028 206. van der Werf C , Hendrix A , Birnie E , Improving usual care after sudden death in the young with focus on inherited cardiac diseases (the CAREFUL study): a community-based intervention study . *Europace* 2016 ; 18 : 592 – 601 . 25833117 10.1093/europace/euv059 207. Tester DJ , Spoon DB , Valdivia HH , Makielski JC , Ackerman MJ . Targeted mutational analysis of the RyR2-encoded cardiac ryanodine receptor in sudden unexplained death: a molecular autopsy of 49 medical examiner/coroner's cases . *Mayo Clin Proc* 2004 ; 79 : 1380 – 1384 . 15544015 10.4065/79.11.1380 208. Anderson JH , Tester DJ , Will ML , Ackerman MJ . Whole-exome molecular autopsy after exertion-related sudden unexplained death in the young . *Circ Cardiovasc Genet* 2016 ; 9 : 259 – 265 . 27114410 10.1161/CIRCGENETICS.115.001370 209. Giudicessi JR , Ackerman MJ . Role of genetic heart disease in sentinel sudden cardiac arrest survivors across the age spectrum . *Int J Cardiol* 2018 ; 270 : 214 – 220 . 29884292 10.1016/j.ijcard.2018.05.100 PMC6364980 210. Shanks GW , Tester DJ , Ackerman JP , Importance of variant interpretation in whole-exome molecular autopsy: population-based case series . *Circulation* 2018 ; 137 : 2705 – 2715 . 29915097 10.1161/CIRCULATIONAHA.117.031053 211. Narula N , Tester DJ , Paulmichl A , Maleszewski JJ , Ackerman MJ . Post-mortem whole exome sequencing with gene-specific analysis for autopsy-negative sudden unexplained death in the young: a case series . *Pediatr Cardiol* 2015 ; 36 : 768 – 778 . 25500949 10.1007/s00246-014-1082-4 PMC4907366 212. Beckmann BM , Wilde AA , Kaab S . Dual inheritance of sudden death from cardiovascular causes . *N Engl J Med* 2008 ; 358 : 2077 – 2078 . 18463390 10.1056/NEJMc0708596 213. Ackerman MJ . Genetic purgatory and the cardiac channelopathies: exposing the variants of uncertain/unknown significance issue . *Heart Rhythm* 2015 ; 12 : 2325 – 2331 . 26144349 10.1016/j.hrthm.2015.07.002 214. Clift K , Macklin S , Halverson C , McCormick JB , Abu Dabrh AM , Hines S . Patients' views on variants of uncertain significance across indications . *J Community Genet* 2020 ; 11 : 139 – 145 . 31432391 10.1007/s12687-019-00434-7 PMC7062975 215. Van Driest SL , Wells QS , Stallings S , Association of arrhythmia-related genetic variants with phenotypes documented in electronic medical records . *JAMA* 2016 ; 315 : 47 – 57 . 26746457 10.1001/jama.2015.17701 PMC4758131 216. Krahn AD , Healey JS , Chauhan V , Systematic assessment of patients with unexplained cardiac arrest: Cardiac Arrest Survivors with Preserved Ejection Fraction Registry (CASPER) . *Circulation* 2009 ; 120 : 278 – 285 . 19597050 10.1161/CIRCULATIONAHA.109.853143 217. Marstrand P , Corell P , Henriksen FL , Pehrson S , Bundgaard H , Theilade J . Clinical evaluation of unselected cardiac arrest survivors in a tertiary center over a 1-year period (the LAZARUS study) . *J Electrocardiol* 2016 ; 49 : 707 – 713 . 27237785 10.1016/j.jelectrocard.2016.05.008 218. Waldmann V , Bougouin W , Karam N , Characteristics and clinical assessment of unexplained sudden cardiac arrest in the real-world setting: focus on idiopathic ventricular fibrillation . *Eur Heart J* 2018 ; 39 : 1981 – 1987 . 29566157 10.1093/eurheartj/ehy098 PMC5982722 219. Allan KS , Morrison LJ , Pinter A , Tu JV , Dorian P , Rescu I . Unexpected high prevalence of cardiovascular disease risk factors and psychiatric disease among young people with sudden cardiac arrest . *J Am Heart Assoc* 2019 ; 8 : e010330 . 30661423 10.1161/JAHA.118.010330 PMC6497342 220. Herman AR , Cheung C , Gerull B , Outcome of apparently unexplained cardiac arrest: results from investigation and follow-up of the prospective Cardiac Arrest Survivors with Preserved Ejection Fraction Registry . *Circ Arrhythm Electrophysiol* 2016 ; 9 : e003619 . 26783233 10.1161/CIRCEP.115.003619 221. Finocchiaro G , Papadakis M , Dhutia H , Obesity and sudden cardiac death in the young: clinical and pathological insights from a large national registry . *Eur J Prev Cardiol* 2018 ; 25 : 395 – 401 . 29319343 10.1177/2047487317751291 222. Mangili LC , Miname MH , Silva PRS , Achilles tendon xanthomas are associated with the presence and burden of subclinical coronary atherosclerosis in heterozygous familial hypercholesterolemia: a pilot study . *Atherosclerosis* 2017 ; 263 : 393 – 397 . 28499609 10.1016/j.atherosclerosis.2017.04.025 223.

Wang S , Peng D . Cardiac involvement in Emery-Dreifuss muscular dystrophy and related management strategies . *Int Heart J* 2019 ; 60 : 12 – 18 . 30518714 10.1536/ihj.17-604 224.

Clemens DJ , Tester DJ , Giudicessi JR , International Triadin Knockout Syndrome Registry . *Circ Genom Precis Med* 2019 ; 12 : e002419 . 30649896 10.1161/CIRCGEN.118.002419 225.

Walsh MA , Turner C , Timothy KW , A multicentre study of patients with Timothy syndrome . *Europace* 2018 ; 20 : 377 – 385 . 28371864 10.1093/europace/euw433 226.

Bitterman AD , Sponseller PD . Marfan syndrome: a clinical update . *J Am Acad Orthop Surg* 2017 ; 25 : 603 – 609 . 28837453 10.5435/JAAOS-D-16-00143 227.

Abrams DJ . How to develop a clinic for sudden cardiac arrest survivors and families of non-survivors . *Cardiol Young* 2017 ; 27 : S3 – S9 . 10.1017/S104795111600216X 28084954 228.

Tseng ZH , Olgin JE , Vittinghoff E , Prospective Countywide Surveillance and Autopsy Characterization of Sudden Cardiac Death: POST SCD Study . *Circulation* 2018 ; 137 : 2689 – 2700 . 29915095 10.1161/CIRCULATIONAHA.117.033427 PMC6013842 229.

Colquhoun MC , Chamberlain DA , Newcombe RG , A national scheme for public access defibrillation in England and Wales: early results . *Resuscitation* 2008 ; 78 : 275 – 280 . 18562074 10.1016/j.resuscitation.2008.03.226 230.

Zijlstra JA , Bekkers LE , Hulleman M , Beesems SG , Koster RW . Automated external defibrillator and operator performance in out-of-hospital cardiac arrest . *Resuscitation* 2017 ; 118 : 140 – 146 . 28526495 10.1016/j.resuscitation.2017.05.017 231.

Yamamoto T , Takayama M , Sato N , Inappropriate analyses of automated external defibrillators used during in-hospital ventricular fibrillation . *Circ J* 2008 ; 72 : 679 – 681 . 18362445 10.1253/circj.72.679 232.

Savastano S , Rordorf R , Vicentini A , A comprehensive electrocardiographic, molecular, and echocardiographic study of Brugada syndrome: validation of the 2013 diagnostic criteria . *Heart Rhythm* 2014 ; 11 : 1176 – 1183 . 24721456 10.1016/j.hrthm.2014.04.010 233.

Govindan M , Batchvarov VN , Raju H , Utility of high and standard right precordial leads during ajmaline testing for the diagnosis of Brugada syndrome . *Heart* 2010 ; 96 : 1904 – 1908 . 20962343 10.1136/hrt.2010.201244 234.

Curcio A , Mazzanti A , Bloise R , Clinical presentation and outcome of Brugada syndrome diagnosed with the new 2013 criteria . *J Cardiovasc Electrophysiol* 2016 ; 27 : 937 – 943 . 27098113 10.1111/jce.12997 235.

Miyamoto K , Yokokawa M , Tanaka K , Diagnostic and prognostic value of a type 1 Brugada electrocardiogram at higher (third or second) V1 to V2 recording in men with Brugada syndrome . *Am J Cardiol* 2007 ; 99 : 53 – 57 . 17196462 10.1016/j.amjcard.2006.07.062 236.

Haïssaguerre M , Shah DC , Jaïs P , Role of Purkinje conducting system in triggering of idiopathic ventricular fibrillation . *Lancet* 2002 ; 359 : 677 – 678 . 11879868 10.1016/S0140-6736(02)07807-8 237.

Divakaran S , Stewart GC , Lakdawala NK , Diagnostic accuracy of advanced imaging in cardiac sarcoidosis . *Circ Cardiovasc Imaging* 2019 ; 12 : e008975 . 31177817 10.1161/CIRCIMAGING.118.008975 PMC6653689 238.

White JA , Fine NM , Gula L , Utility of cardiovascular magnetic resonance in identifying substrate for malignant ventricular arrhythmias . *Circ Cardiovasc Imaging* 2012 ; 5 : 12 – 20 . 22038987 10.1161/CIRCIMAGING.111.966085 239.

Basso C , Perazzolo Marra M , Rizzo S , Arrhythmic mitral valve prolapse and sudden cardiac death . *Circulation* 2015 ; 132 : 556 – 566 . 26160859 10.1161/CIRCULATIONAHA.115.016291 240.

Basso C , Iliceto S , Thiene G , Perazzolo Marra M . Mitral valve prolapse, ventricular arrhythmias, and sudden death . *Circulation* 2019 ; 140 : 952 – 964 . 31498700 10.1161/CIRCULATIONAHA.118.034075 241.

Eckart RE , Scoville SL , Campbell CL , Sudden death in young adults: a 25-year review of autopsies in military recruits . *Ann Intern Med* 2004 ; 141 : 829 – 834 . 15583223 10.7326/0003-4819-141-11-200412070-00005 242.

Neumar RW , Shuster M , Callaway CW , Part 1: Executive Summary: 2015 American Heart Association guidelines update for cardiopulmonary resuscitation and emergency cardiovascular care . *Circulation* 2015 ; 132 : S315 – S367 . 26472989 10.1161/CIR.0000000000000252 243.

Avari Silva JN , Bromberg BI , Emge FK , Bowman TM , Van Hare GF . Implantable loop recorder monitoring for refining management of children with inherited arrhythmia syndromes . *J Am Heart Assoc* 2016 ; 5 : e003632 . 27231019 10.1161/JAHA.116.003632 PMC4937287 244.

Placidi S , Drago F , Milioni M , Miniaturized implantable loop recorder in small patients: an effective approach to the evaluation of subjects at risk of sudden death . *Pacing Clin Electrophysiol* 2016 ; 39 : 669 – 674 . 27062386 10.1111/pace.12866 245.

Sweeney MO , Ruetz LL , Belk P , Mullen TJ , Johnson JW , Sheldon T . Bradycardia pacing-induced short-long-short sequences at the onset of ventricular tachyarrhythmias: a possible mechanism of proarrhythmia? *J Am Coll Cardiol* 2007 ; 50 : 614 – 622 . 17692746 10.1016/j.jacc.2007.02.077 246.

Jiménez-Jáimez J , Peinado R , Grima EZ , Diagnostic approach to unexplained cardiac arrest (from the FIVI-Gen Study) . *Am J Cardiol* 2015 ; 116 : 894 –

899 . 26189708 10.1016/j.amjcard.2015.06.030 247. Kamath GS , Zareba W , Delaney J , Value of the signal-averaged electrocardiogram in arrhythmogenic right ventricular cardiomyopathy/dysplasia . *Heart Rhythm* 2011 ; 8 : 256 – 262 . 20933608 10.1016/j.hrthm.2010.10.007 PMC3077127 248. Rodrigues P , Joshi A , Williams H , Diagnosis and prognosis in sudden cardiac arrest survivors without coronary artery disease: utility of a clinical approach using cardiac magnetic resonance imaging . *Circ Cardiovasc Imaging* 2017 ; 10 : e006709 . 29237609 10.1161/CIRCIMAGING.117.006709 249. Ali-Ahmed F , Dalgaard F , Al-Khatib SM . Sudden cardiac death in patients with myocarditis: evaluation, risk stratification, and management . *Am Heart J* 2020 ; 220 : 29 – 40 . 31765933 10.1016/j.ahj.2019.08.007 250. Chatterjee D , Pieroni M , Fatah M , An autoantibody profile detects Brugada syndrome and identifies abnormally expressed myocardial proteins . *Eur Heart J* 2020 ; 41 : 2878 – 2890 . 32533187 10.1093/eurheartj/ehaa383 251. Wilde AAM , Lodder EM . A highly specific biomarker for Brugada syndrome. Also too good to be true? *Eur Heart J* 2020 ; 41 : 2891 – 2893 . 32558881 10.1093/eurheartj/ehaa468 252. Tschabrunn CM , Haqqani HM , Santangeli P , Zado ES , Marchlinski FE . 12-Lead electrocardiogram to localize region of abnormal electroanatomic substrate in arrhythmogenic right ventricular cardiomyopathy . *JACC Clin Electrophysiol* 2017 ; 3 : 654 – 665 . 29759533 10.1016/j.jacep.2017.01.009 253. Müller D , Schnitzer L , Brandt J , Arntz HR . The accuracy of an out-of-hospital 12-lead ECG for the detection of ST-elevation myocardial infarction immediately after resuscitation . *Ann Emerg Med* 2008 ; 52 : 658 – 664 . 18722690 10.1016/j.annemergmed.2008.06.469 254. Stær-Jensen H , Nakstad ER , Fossum E , Post-resuscitation ECG for selection of patients for immediate coronary angiography in out-of-hospital cardiac arrest . *Circ Cardiovasc Interv* 2015 ; 8 : e002784 . 26453688 10.1161/CIRCINTERVENTIONS.115.002784 255. Eysmann SB , Marchlinski FE , Buxton AE , Josephson ME . Electrocardiographic changes after cardioversion of ventricular arrhythmias . *Circulation* 1986 ; 73 : 73 – 81 . 3940671 10.1161/01.cir.73.1.73 256. Kim YJ , Min SY , Lee DH , The role of post-resuscitation electrocardiogram in patients with ST-segment changes in the immediate post-cardiac arrest period . *JACC Cardiovasc Interv* 2017 ; 10 : 451 – 459 . 28279312 10.1016/j.jcin.2016.11.046 257. Rolfast CL , Lust EJ , de Cock CC . Electrocardiographic changes in therapeutic hypothermia . *Crit Care* 2012 ; 16 : R100 . 22673196 10.1186/cc11369 PMC3580650 258. Salinas P , Lopez-de-Sa E , Pena-Conde L , Electrocardiographic changes during induced therapeutic hypothermia in comatose survivors after cardiac arrest . *World J Cardiol* 2015 ; 7 : 423 – 430 . 26225204 10.4330/wjc.v7.i7.423 PMC4513495 259. Bière L , Niro M , Pouliquen H , Risk of ventricular arrhythmia in patients with myocardial infarction and non-obstructive coronary arteries and normal ejection fraction . *World J Cardiol* 2017 ; 9 : 268 – 276 . 28400924 10.4330/wjc.v9.i3.268 PMC5368677 260. Marcus FI , McKenna WJ , Sherrill D , Diagnosis of arrhythmogenic right ventricular cardiomyopathy/dysplasia: proposed modification of the task force criteria . *Circulation* 2010 ; 121 : 1533 – 1541 . 20172911 10.1161/CIRCULATIONAHA.108.840827 PMC2860804 261. Hamid MS , Norman M , Quraishi A , Prospective evaluation of relatives for familial arrhythmogenic right ventricular cardiomyopathy/dysplasia reveals a need to broaden diagnostic criteria . *J Am Coll Cardiol* 2002 ; 40 : 1445 – 1450 . 12392835 10.1016/s0735-1097(02)02307-0 262. Nava A , Bauce B , Basso C , Clinical profile and long-term follow-up of 37 families with arrhythmogenic right ventricular cardiomyopathy . *J Am Coll Cardiol* 2000 ; 36 : 2226 – 2233 . 11127465 10.1016/s0735-1097(00)00997-9 263. Perrin MJ , Angaran P , Laksman Z , Exercise testing in asymptomatic gene carriers exposes a latent electrical substrate of arrhythmogenic right ventricular cardiomyopathy . *J Am Coll Cardiol* 2013 ; 62 : 1772 – 1779 . 23810883 10.1016/j.jacc.2013.04.084 264. Amin AS , de Groot EA , Ruijter JM , Wilde AA , Tan HL . Exercise-induced ECG changes in Brugada syndrome . *Circ Arrhythm Electrophysiol* 2009 ; 2 : 531 – 539 . 19843921 10.1161/CIRCEP.109.862441 265. Priori SG , Napolitano C , Memmi M , Clinical and molecular characterization of patients with catecholaminergic polymorphic ventricular tachycardia . *Circulation* 2002 ; 106 : 69 – 74 . 12093772 10.1161/01.cir.0000020013.73106.d8 266. Adler A , van der Werf C , Postema PG , The phenomenon of “QT stunning”: the abnormal QT prolongation provoked by standing persists even as the heart rate returns to normal in patients with long QT syndrome . *Heart Rhythm* 2012 ; 9 : 901 – 908 . 22300664 10.1016/j.hrthm.2012.01.026 PMC3478123 267. Dionne A , Fournier A , Dahdah N , Abrams D , Khairy P , Abadir S . Dynamic QT interval changes from supine to standing in healthy children . *Can J Cardiol* 2018 ; 34 : 66 – 72 . 29275885 10.1016/j.cjca.2017.10.016 268. Filippini L , Postema PG , Zoubin K , The brisk-standing-test for long QT syndrome in prepubertal school children: defining normal . *Europace*

2018 ; 20 : f108 – f112 . 29036559 10.1093/europace/eux259 269. Viskin S , Postema PG , Bhuiyan ZA , The response of the QT interval to the brief tachycardia provoked by standing: a bedside test for diagnosing long QT syndrome . J Am Coll Cardiol 2010 ; 55 : 1955 – 1961 . 20116193 10.1016/j.jacc.2009.12.015 PMC2862092 270. Sy RW , van der Werf C , Chattha IS , Derivation and validation of a simple exercise-based algorithm for prediction of genetic testing in relatives of LQTS probands . Circulation 2011 ; 124 : 2187 – 2194 . 22042885 10.1161/CIRCULATIONAHA.111.028258 271. Wong JA , Gula LJ , Klein GJ , Yee R , Skanes AC , Krahn AD . Utility of treadmill testing in identification and genotype prediction in long-QT syndrome . Circ Arrhythm Electrophysiol 2010 ; 3 : 120 – 125 . 20071715 10.1161/CIRCEP.109.907865 272. Schwartz PJ , Crotti L . QTc behavior during exercise and genetic testing for the long-QT syndrome . Circulation 2011 ; 124 : 2181 – 2184 . 22083145 10.1161/CIRCULATIONAHA.111.062182 273. Shimizu W , Noda T , Takaki H , Diagnostic value of epinephrine test for genotyping LQT1, LQT2, and LQT3 forms of congenital long QT syndrome . Heart Rhythm 2004 ; 1 : 276 – 283 . 15851169 10.1016/j.hrthm.2004.04.021 274. Ackerman MJ , Khositseth A , Tester DJ , Hejlik JB , Shen WK , Porter CB . Epinephrine-induced QT interval prolongation: a gene-specific paradoxical response in congenital long QT syndrome . Mayo Clin Proc 2002 ; 77 : 413 – 421 . 12004990 10.4065/77.5.413 275. Krahn AD , Healey JS , Chauhan VS , Epinephrine infusion in the evaluation of unexplained cardiac arrest and familial sudden death: from the Cardiac Arrest Survivors with Preserved Ejection Fraction Registry . Circ Arrhythm Electrophysiol 2012 ; 5 : 933 – 940 . 22944906 10.1161/CIRCEP.112.973230 276. Magnano AR , Talathoti N , Hallur R , Bloomfield DM , Garan H . Sympathomimetic infusion and cardiac repolarization: the normative effects of epinephrine and isoproterenol in healthy subjects . J Cardiovasc Electrophysiol 2006 ; 17 : 983 – 989 . 16879629 10.1111/j.1540-8167.2006.00555.x 277. Marjamaa A , Hiippala A , Arrhenius B , Intravenous epinephrine infusion test in diagnosis of catecholaminergic polymorphic ventricular tachycardia . J Cardiovasc Electrophysiol 2012 ; 23 : 194 – 199 . 21954897 10.1111/j.1540-8167.2011.02188.x 278. Denis A , Sacher F , Derval N , Diagnostic value of isoproterenol testing in arrhythmogenic right ventricular cardiomyopathy . Circ Arrhythm Electrophysiol 2014 ; 7 : 590 – 597 . 24970294 10.1161/CIRCEP.113.001224 279. Meregalli PG , Ruijter JM , Hofman N , Bezzina CR , Wilde AA , Tan HL . Diagnostic value of flecainide testing in unmasking SCN5A-related Brugada syndrome . J Cardiovasc Electrophysiol 2006 ; 17 : 857 – 864 . 16764707 10.1111/j.1540-8167.2006.00531.x 280. Hong K , Brugada J , Oliva A , Value of electrocardiographic parameters and ajmaline test in the diagnosis of Brugada syndrome caused by SCN5A mutations . Circulation 2004 ; 110 : 3023 – 3027 . 15520322 10.1161/01.CIR.0000144299.17008.07 PMC1513622 281. Brugada R , Brugada J , Antzelevitch C , Sodium channel blockers identify risk for sudden death in patients with ST-segment elevation and right bundle branch block but structurally normal hearts . Circulation 2000 ; 101 : 510 – 515 . 10662748 10.1161/01.cir.101.5.510 282. Krahn AD , Gollob M , Yee R , Diagnosis of unexplained cardiac arrest: role of adrenaline and procainamide infusion . Circulation 2005 ; 112 : 2228 – 2234 . 16203906 10.1161/CIRCULATIONAHA.105.552166 283. Sangwatanaroj S , Prechawat S , Sunsaneewitayakul B , Sitthisook S , Tosukhowong P , Tungsanga K . New electrocardiographic leads and the procainamide test for the detection of the Brugada sign in sudden unexplained death syndrome survivors and their relatives . Eur Heart J 2001 ; 22 : 2290 – 2296 . 11728150 10.1053/euhj.2001.2691 284. Somani R , Krahn AD , Healey JS , Procainamide infusion in the evaluation of unexplained cardiac arrest: from the Cardiac Arrest Survivors with Preserved Ejection Fraction Registry (CASPER) . Heart Rhythm 2014 ; 11 : 1047 – 1054 . 24657429 10.1016/j.hrthm.2014.03.022 285. Cheung CC , Mellor G , Deyell MW , Comparison of ajmaline and procainamide provocation tests in the diagnosis of Brugada syndrome . JACC Clin Electrophysiol 2019 ; 5 : 504 – 512 . 31000106 10.1016/j.jacep.2019.01.026 286. Hasdemir C , Payzin S , Kocabas U , High prevalence of concealed Brugada syndrome in patients with atrioventricular nodal reentrant tachycardia . Heart Rhythm 2015 ; 12 : 1584 – 1594 . 25998140 10.1016/j.hrthm.2015.03.015 287. Ikeda T , Abe A , Yusu S , The full stomach test as a novel diagnostic technique for identifying patients at risk of Brugada syndrome . J Cardiovasc Electrophysiol 2006 ; 17 : 602 – 607 . 16836706 10.1111/j.1540-8167.2006.00424.x 288. Huchet F , Kyndt F , Barc J , Familial catecholamine-induced QT prolongation in unexplained sudden cardiac death . J Am Coll Cardiol 2017 ; 69 : 1642 – 1643 . 28335847 10.1016/j.jacc.2017.01.030 289. Kannankeril PJ , Roden DM , Norris KJ , Whalen SP , George AL Jr , Murray KT . Genetic susceptibility to acquired long QT syndrome: pharmacologic challenge in first-degree relatives . Heart Rhythm 2005 ; 2 : 134 – 140 . 15851285 10.1016/j.hrthm.2004.10.039 290. Etienne P ,

Huchet F , Gaborit N , Mental stress test: a rapid, simple, and efficient test to unmask long QT syndrome . *Europace* 2018 ; 20 : 2014 – 2020 . 29688407 10.1093/europace/euy078 291. Kaab S , Hinterseer M , Nabauer M , Steinbeck G . Sotalol testing unmasks altered repolarization in patients with suspected acquired long-QT-syndrome: a case-control pilot study using i.v. sotalol . *Eur Heart J* 2003 ; 24 : 649 – 657 . 12657223 10.1016/s0195-668x(02)00806-0 292. Kamakura T , Wada M , Ishibashi K , Significance of coronary artery spasm diagnosis in patients with early repolarization syndrome . *J Am Heart Assoc* 2018 ; 7 : e007942 . 29436392 10.1161/JAHA.117.007942 PMC5850201 293. Beltrame JF , Crea F , Kaski JC , International standardization of diagnostic criteria for vasospastic angina . *Eur Heart J* 2017 ; 38 : 2565 – 2568 . 26245334 10.1093/eurheartj/ehv351 294. Nakao K , Ohgushi M , Yoshimura M , Hyperventilation as a specific test for diagnosis of coronary artery spasm . *Am J Cardiol* 1997 ; 80 : 545 – 549 . 9294979 10.1016/s0002-9149(97)00419-0 295. Waldmann V , Bougouin W , Karam N , Coronary vasospasm-related sudden cardiac arrest in the community . *J Am Coll Cardiol* 2018 ; 72 : 814 – 815 . 30092958 10.1016/j.jacc.2018.05.051 296. Garratt CJ , Antoniou A , Griffith MJ , Ward DE , Camm AJ . Use of intravenous adenosine in sinus rhythm as a diagnostic test for latent preexcitation . *Am J Cardiol* 1990 ; 65 : 868 – 873 . 2321537 10.1016/0002-9149(90)91428-9 297. Foo FS , Stiles MK , Heaven D . Unmasking latent preexcitation of a right-sided accessory pathway with intravenous adenosine after unexplained sudden cardiac arrest . *J Arrhythm* 2020 ; 36 : 939 – 941 . 33024474 10.1002/joa3.12408 PMC7532269 298. Wang YS , Scheinman MM , Chien WW , Cohen TJ , Lesh MD , Griffin JC . Patients with supraventricular tachycardia presenting with aborted sudden death: incidence, mechanism and long-term follow-up . *J Am Coll Cardiol* 1991 ; 18 : 1711 – 1719 . 1960318 10.1016/0735-1097(91)90508-7 299. Roberts JD , Gollob MH , Young C , Bundle branch re-entrant ventricular tachycardia: novel genetic mechanisms in a life-threatening arrhythmia . *JACC Clin Electrophysiol* 2017 ; 3 : 276 – 288 . 29759522 10.1016/j.jacep.2016.09.019 300. Santangeli P , Hamilton-Craig C , Dello Russo A , Imaging of scar in patients with ventricular arrhythmias of right ventricular origin: cardiac magnetic resonance versus electroanatomic mapping . *J Cardiovasc Electrophysiol* 2011 ; 22 : 1359 – 1366 . 21736658 10.1111/j.1540-8167.2011.02127.x 301. Haissaguerre M , Hocini M , Cheniti G , Localized structural alterations underlying a subset of unexplained sudden cardiac death . *Circ Arrhythm Electrophysiol* 2018 ; 11 : e006120 . 30002064 10.1161/CIRCEP.117.006120 PMC7661047 302. Horner JM , Horner MM , Ackerman MJ . The diagnostic utility of recovery phase QTc during treadmill exercise stress testing in the evaluation of long QT syndrome . *Heart Rhythm* 2011 ; 8 : 1698 – 1704 . 21699858 10.1016/j.hrthm.2011.05.018 303. Giudicessi JR , Ackerman MJ . Exercise testing oversights underlie missed and delayed diagnosis of catecholaminergic polymorphic ventricular tachycardia in young sudden cardiac arrest survivors . *Heart Rhythm* 2019 ; 16 : 1232 – 1239 . 30763784 10.1016/j.hrthm.2019.02.012 304. Vyas H , Hejlik J , Ackerman MJ . Epinephrine QT stress testing in the evaluation of congenital long-QT syndrome: diagnostic accuracy of the paradoxical QT response . *Circulation* 2006 ; 113 : 1385 – 1392 . 16534005 10.1161/CIRCULATIONAHA.105.600445 305. Rehm HL , Berg JS , Brooks LD , ClinGen: the clinical genome resource . *N Engl J Med* 2015 ; 372 : 2235 – 2242 . 26014595 10.1056/NEJMSr1406261 PMC4474187 306. Richards S , Aziz N , Bale S , Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology . *Genet Med* 2015 ; 17 : 405 – 424 . 25741868 10.1038/gim.2015.30 PMC4544753 307. Gersh BJ , Maron BJ , Bonow RO , 2011 ACCF/AHA guideline for the diagnosis and treatment of hypertrophic cardiomyopathy: executive summary: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines . *Circulation* 2011 ; 124 : 2761 – 2796 . 22068435 10.1161/CIR.0b013e318223e230 308. Burns C , Bagnall RD , Lam L , Semsarian C , Ingles J . Multiple gene variants in hypertrophic cardiomyopathy in the era of next-generation sequencing . *Circ Cardiovasc Genet* 2017 ; 10 : e001666 . 28790153 10.1161/CIRCGENETICS.116.001666 309. Amendola LM , Dorschner MO , Robertson PD , Actionable exomic incidental findings in 6503 participants: challenges of variant classification . *Genome Res* 2015 ; 25 : 305 – 315 . 25637381 10.1101/gr.183483.114 PMC4352885 310. Mohammed S , Lim Z , Dean PH , Genetic insurance discrimination in sudden arrhythmia death syndromes: empirical evidence from a cross-sectional survey in North America . *Circ Cardiovasc Genet* 2017 ; 10 : e001442 . 28119333 10.1161/CIRCGENETICS.116.001442 311. Moss AJ , Zareba W , Hall WJ , Effectiveness and limitations of beta-blocker therapy in congenital long-QT

syndrome . *Circulation* 2000 ; 101 : 616 – 623 . 10673253 10.1161/01.cir.101.6.616 312. Schwartz PJ , Priori SG , Locati EH , Long QT syndrome patients with mutations of the SCN5A and HERG genes have differential responses to Na<sup>+</sup> channel blockade and to increases in heart rate: implications for gene-specific therapy . *Circulation* 1995 ; 92 : 3381 – 3386 . 8521555 10.1161/01.cir.92.12.3381 313. Moss AJ , Windle JR , Hall WJ , Safety and efficacy of flecainide in subjects with Long QT-3 syndrome (DeltaKPQ mutation): a randomized, double-blind, placebo-controlled clinical trial . *Ann Noninvasive Electrocardiol* 2005 ; 10 : 59 – 66 . 16274417 10.1111/j.1542-474X.2005.00077.x PMC6932190 314. Moss AJ , Zareba W , Schwarz KQ , Rosero S , McNitt S , Robinson JL . Ranolazine shortens repolarization in patients with sustained inward sodium current due to type-3 long-QT syndrome . *J Cardiovasc Electrophysiol* 2008 ; 19 : 1289 – 1293 . 18662191 10.1111/j.1540-8167.2008.01246.x PMC2614458 315. Kannankeril PJ , Moore JP , Cerrone M , Efficacy of flecainide in the treatment of catecholaminergic polymorphic ventricular tachycardia: a randomized clinical trial . *JAMA Cardiol* 2017 ; 2 : 759 – 766 . 28492868 10.1001/jamacardio.2017.1320 PMC5548393 316. James CA , Bhonsale A , Tichnell C , Exercise increases age-related penetrance and arrhythmic risk in arrhythmogenic right ventricular dysplasia/cardiomyopathy-associated desmosomal mutation carriers . *J Am Coll Cardiol* 2013 ; 62 : 1290 – 1297 . 23871885 10.1016/j.jacc.2013.06.033 PMC3809992 317. Ruwald AC , Marcus F , Estes NA 3rd , Association of competitive and recreational sport participation with cardiac events in patients with arrhythmogenic right ventricular cardiomyopathy: results from the North American multidisciplinary study of arrhythmogenic right ventricular cardiomyopathy . *Eur Heart J* 2015 ; 36 : 1735 – 1743 . 25896080 10.1093/eurheartj/ehv110 PMC4500847 318. van Rijsingen IA , van der Zwaag PA , Groeneweg JA , Outcome in phospholamban R14del carriers: results of a large multicentre cohort study . *Circ Cardiovasc Genet* 2014 ; 7 : 455 – 465 . 24909667 10.1161/CIRCGENETICS.113.000374 319. van Rijsingen IA , Arbustini E , Elliott PM , Risk factors for malignant ventricular arrhythmias in Lamin A/C mutation carriers: a European cohort study . *J Am Coll Cardiol* 2012 ; 59 : 493 – 500 . 22281253 10.1016/j.jacc.2011.08.078 320. De Ferrari GM , Dusi V , Spazzolini C , Clinical management of catecholaminergic polymorphic ventricular tachycardia: the role of left cardiac sympathetic denervation . *Circulation* 2015 ; 131 : 2185 – 2193 . 26019152 10.1161/CIRCULATIONAHA.115.015731 321. Coleman MA , Bos JM , Johnson JN , Videoscopic left cardiac sympathetic denervation for patients with recurrent ventricular fibrillation/malignant ventricular arrhythmia syndromes besides congenital long-QT syndrome . *Circ Arrhythm Electrophysiol* 2012 ; 5 : 782 – 788 . 22787014 10.1161/CIRCEP.112.971754 PMC3912558 322. Bos JM , Crotti L , Rohatgi RK , Mexiletine shortens the QT interval in patients with potassium channel-mediated type 2 long QT syndrome . *Circ Arrhythm Electrophysiol* 2019 ; 12 : e007280 . 31006312 10.1161/CIRCEP.118.007280 323. Asatryan B , Schaller A , Seiler J , Usefulness of genetic testing in sudden cardiac arrest survivors with or without previous clinical evidence of heart disease . *Am J Cardiol* 2019 ; 123 : 2031 – 2038 . 30975432 10.1016/j.amjcard.2019.02.061 324. Ashar FN , Mitchell RN , Albert CM , A comprehensive evaluation of the genetic architecture of sudden cardiac arrest . *Eur Heart J* 2018 ; 39 : 3961 – 3969 . 30169657 10.1093/eurheartj/ehy474 PMC6247663 325. Milano A , Blom MT , Lodder EM , Sudden cardiac arrest and rare genetic variants in the community . *Circ Cardiovasc Genet* 2016 ; 9 : 147 – 153 . 26800703 10.1161/CIRCGENETICS.115.001263 326. Visser M , Dooijes D , van der Smagt JJ , Next-generation sequencing of a large gene panel in patients initially diagnosed with idiopathic ventricular fibrillation . *Heart Rhythm* 2017 ; 14 : 1035 – 1040 . 28087426 10.1016/j.hrthm.2017.01.010 327. Visser M , van der Heijden JF , van der Smagt JJ , Long-term outcome of patients initially diagnosed with idiopathic ventricular fibrillation: a descriptive study . *Circ Arrhythm Electrophysiol* 2016 ; 9 : e004258 . 27733492 10.1161/CIRCEP.116.004258 328. Ingles J , Bagnall RD , Yeates L , Concealed arrhythmogenic right ventricular cardiomyopathy in sudden unexplained cardiac death events . *Circ Genom Precis Med* 2018 ; 11 : e002355 . 30571190 10.1161/CIRCGEN.118.002355 329. Yang P , Kanki H , Drolet B , Allelic variants in long-QT disease genes in patients with drug-associated torsades de pointes . *Circulation* 2002 ; 105 : 1943 – 1948 . 11997281 10.1161/01.cir.0000014448.19052.4c 330. Tester DJ , Ackerman JP , Giudicessi JR , Plakophilin-2 truncation variants in patients clinically diagnosed with catecholaminergic polymorphic ventricular tachycardia and decedents with exercise-associated autopsy negative sudden unexplained death in the young . *JACC Clin Electrophysiol* 2019 ; 5 : 120 – 127 . 30678776 10.1016/j.jacep.2018.09.010 PMC6394846 331. van der Werf C , Kannankeril PJ , Sacher F , Flecainide therapy reduces exercise-induced ventricular arrhythmias in patients with

catecholaminergic polymorphic ventricular tachycardia . J Am Coll Cardiol 2011 ; 57 : 2244 – 2254 . 21616285 10.1016/j.jacc.2011.01.026 PMC3495585 332. Semsarian C , Ingles J , Wilde AA . Sudden cardiac death in the young: the molecular autopsy and a practical approach to surviving relatives . Eur Heart J 2015 ; 36 : 1290 – 1296 . 25765769 10.1093/eurheartj/ehv063 333. Asatryan B , Medeiros-Domingo A . Molecular and genetic insights into progressive cardiac conduction disease . Europace 2019 ; 21 : 1145 – 1158 . 31087102 10.1093/europace/euz109 334. Quenin P , Kyndt F , Mabo P , Clinical yield of familial screening after sudden death in young subjects: the French experience . Circ Arrhythm Electrophysiol 2017 ; 10 : e005236 . 28912206 10.1161/CIRCEP.117.005236 335. Cornel MC , van El CG . Barriers and facilitating factors for implementation of genetic services: a public health perspective . Front Public Health 2017 ; 5 : 195 . 28824901 10.3389/fpubh.2017.00195 PMC5543075 336. Charron P , Heron D , Gargiulo M , Prenatal molecular diagnosis in hypertrophic cardiomyopathy: report of the first case . Prenat Diagn 2004 ; 24 : 701 – 703 . 15386449 10.1002/pd.969 337. Harper JC , Wilton L , Traeger-Synodinos J , The ESHRE PGD Consortium: 10 years of data collection . Hum Reprod Update 2012 ; 18 : 234 – 247 . 22343781 10.1093/humupd/dmr052 338. Kuliev A , Pomerantseva E , Polling D , Verlinsky O , Rechitsky S . PGD for inherited cardiac diseases . Reprod Biomed Online 2012 ; 24 : 443 – 453 . 22386593 10.1016/j.rbmo.2011.12.009 339. van Velzen HG , Schinkel AFL , Baart SJ , Outcomes of contemporary family screening in hypertrophic cardiomyopathy . Circ Genom Precis Med 2018 ; 11 : e001896 . 29661763 10.1161/CIRCGEN.117.001896 340. Medeiros Domingo A , Bolliger S , Grani C , Recommendations for genetic testing and counselling after sudden cardiac death: practical aspects for Swiss practice . Swiss Med Wkly 2018 ; 148 : w14638 . 30044475 10.57187/smw.2018.14638 341. Rootwelt-Norberg C , Lie OH , Dejgaard LA , Life-threatening arrhythmic presentation in patients with arrhythmogenic cardiomyopathy before and after entering the genomic era: a two-decade experience from a large volume center . Int J Cardiol 2019 ; 279 : 79 – 83 . 30638987 10.1016/j.ijcard.2018.12.066 342. Lafreniere-Roula M , Bolkier Y , Zahavich L , Family screening for hypertrophic cardiomyopathy: is it time to change practice guidelines? Eur Heart J 2019 ; 40 : 3672 – 3681 . 31170284 10.1093/eurheartj/ehz396 PMC6885133 343. Christian S , Atallah J , Clegg R , Uptake of predictive genetic testing and cardiac evaluation for children at risk for an inherited arrhythmia or cardiomyopathy . J Genet Couns 2018 ; 27 : 124 – 130 . 28699125 10.1007/s10897-017-0129-0 344. Christian S , Somerville M , Huculak C , Atallah J . Practice variation among an international group of genetic counselors on when to offer predictive genetic testing to children at risk of an inherited arrhythmia or cardiomyopathy . J Genet Couns 2019 ; 28 : 70 – 79 . 10.1007/s10897-018-0293-x 30132189 345. Harris S , Cirino AL , Carr CW , The uptake of family screening in hypertrophic cardiomyopathy and an online video intervention to facilitate family communication . Mol Genet Genomic Med 2019 ; 7 : e940 . 31482667 10.1002/mgg3.940 PMC6825857 346. Knight LM , Miller E , Kovach J , Genetic testing and cascade screening in pediatric long QT syndrome and hypertrophic cardiomyopathy . Heart Rhythm 2020 ; 17 : 106 – 112 . 31229680 10.1016/j.hrthm.2019.06.015 347. Shah LL , Daack-Hirsch S , Ersig AL , Paik A , Ahmad F , Williams J . Family relationships associated with communication and testing for inherited cardiac conditions . West J Nurs Res 2019 ; 41 : 1576 – 1601 . 30539690 10.1177/0193945918817039 PMC6561845 348. Adler A , Sadek MM , Chan AY , Patient outcomes from a specialized inherited arrhythmia clinic . Circ Arrhythm Electrophysiol 2016 ; 9 : e003440 . 26743238 10.1161/CIRCEP.115.003440 349. Boers SN , van Delden JJ , Knoers NV , Bredenoord AL . Postmortem disclosure of genetic information to family members: active or passive? Trends Mol Med 2015 ; 21 : 148 – 153 . 25743261 10.1016/j.molmed.2015.01.002 350. Fox D , Spencer E , Torkamani A . Returning results to family members: professional duties in genomics research in the United States . J Leg Med 2018 ; 38 : 201 – 219 . 30289737 10.1080/01947648.2017.1417934 351. van El CG , Baccolini V , Piko P , Cornel MC . Stakeholder views on active cascade screening for familial hypercholesterolemia . Healthcare (Basel) 2018 ; 6 : 108 . 10.3390/healthcare6030108 PMC6163326 30200297 352. Kaufenstein S , Herz N , Scheiper S , Relevance of molecular testing in patients with a family history of sudden death . Forensic Sci Int 2017 ; 276 : 18 – 23 . 28472724 10.1016/j.forsciint.2017.04.001 353. Roston TM , Dewar L , Franciosi S , The accessibility and utilization of genetic testing for inherited heart rhythm disorders: a Canadian cross-sectional survey study . J Community Genet 2018 ; 9 : 257 – 262 . 29170972 10.1007/s12687-017-0348-y PMC6002300 354. Shah LL , Daack-Hirsch S . Family communication about genetic risk of hereditary cardiomyopathies and arrhythmias: an integrative review . J Genet Couns 2018 ; 27 : 1022 – 1039 . 29492742 10.1007/s10897-018-0225-9 PMC6113126 355. Catchpool M , Ramchand

J , Martyn M , A cost-effectiveness model of genetic testing and periodical clinical screening for the evaluation of families with dilated cardiomyopathy . *Genet Med* 2019 ; 21 : 2815 – 2822 . 31222143 10.1038/s41436-019-0582-2 PMC6892743 356. Mak CM , Mok NS , Shum HC , Sudden arrhythmia death syndrome in young victims: a five-year retrospective review and two-year prospective molecular autopsy study by next-generation sequencing and clinical evaluation of their first-degree relatives . *Hong Kong Med J* 2019 ; 25 : 21 – 29 . 10.12809/hkmj187256 30670673 357. van den Heuvel LM , Huisinga MJ , Hoedemaekers YM , Informing relatives at risk of inherited cardiac conditions: experiences and attitudes of healthcare professionals and counselees . *Eur J Hum Genet* 2019 ; 27 : 1341 – 1350 . 31053782 10.1038/s41431-019-0410-9 PMC6777450 358. Broekhuizen K , van Poppel MN , Koppes LL , Kindt I , Brug J , van Mechelen W . No significant improvement of cardiovascular disease risk indicators by a lifestyle intervention in people with familial hypercholesterolemia compared to usual care: results of a randomised controlled trial . *BMC Res Notes* 2012 ; 5 : 181 . 22490761 10.1186/1756-0500-5-181 PMC3392724 359. Raju H , Parsons S , Thompson TN , Insights into sudden cardiac death: exploring the potential relevance of non-diagnostic autopsy findings . *Eur Heart J* 2019 ; 40 : 831 – 838 . 30380018 10.1093/eurheartj/ehy654 360. te Riele AS , James CA , Rastegar N , Yield of serial evaluation in at-risk family members of patients with ARVD/C . *J Am Coll Cardiol* 2014 ; 64 : 293 – 301 . 25034067 10.1016/j.jacc.2014.04.044 PMC4380221 361. Hasdemir C , Juang JJ , Kose S , Coexistence of atrioventricular accessory pathways and drug-induced type 1 Brugada pattern . *Pacing Clin Electrophysiol* 2018 ; 41 : 1078 – 1092 . 29953624 10.1111/pace.13414 PMC6168386 362. Conte G , de Asmundis C , Ciconte G , Follow-up from childhood to adulthood of individuals with family history of Brugada syndrome and normal electrocardiograms . *JAMA* 2014 ; 312 : 2039 – 2041 . 25399282 10.1001/jama.2014.13752

Figure 1 Distribution of causes of death among autopsied cases of sudden cardiac death (n = 753) according to age in persons aged 1–49 years in Denmark (J.T.-H., unpublished data). SADS = sudden arrhythmic death syndrome. \*Coronary artery disease, especially in young persons, may be due to inherited disease (eg, familial hypercholesterolemia).

Cumulative overall survival rates, by participating emergency medical services agency—Cardiac Arrest Registry to Enhance Survival (CARES), United States, October 1, 2005–December 31, 2010. Agencies sorted by total number of out-of-hospital cardiac arrest events in CARES (from low to high; range: 18–5,434). 60

Cumulative Utstein survival rates (patients alive when arriving to hospital) by participating emergency medical services agency—Cardiac Arrest Registry to Enhance Survival (CARES), United States, October 1, 2005–December 31, 2010. Agencies sorted by total number of out-of-hospital cardiac arrest events in CARES (from low to high). 60 †Utstein survival refers to survival to hospital discharge of persons whose cardiac arrest events were witnessed by a bystander and had an initial rhythm of ventricular fibrillation or pulseless ventricular tachycardia (range: 0–598).

Bystander treatments of patients with out-of-hospital cardiac arrest before emergency medical services arrival among neighborhoods by percentage of black residents. Reprinted with permission from the American Medical Association. 39 AED = automated external defibrillator; CPR = cardiopulmonary resuscitation.

Bystander-witnessed arrest, bystander cardiopulmonary resuscitation (CPR), shockable heart rhythm as first recorded rhythm, and survival on arrival at the hospital, Denmark, 2001–2010. Reprinted with permission from the American Medical Association. 54

Examples of complimentary bystander cardiopulmonary resuscitation (CPR) programs. Reprinted with permission from Elsevier. 64

Referral flow for cardiac genetic investigation of sudden cardiac death (SCD) or resuscitated sudden cardiac arrest (SCA). MDT = multidisciplinary team.

Participants in a cardiac genetic service. “Pathologists” includes forensic pathologists. Modified with permission from Elsevier. 69

Psychological care following a sudden cardiac arrest (SCA) or a sudden unexplained death (SUD) where a genetic cause is suspected. Colors correspond to the Class of Recommendation in Table 1 .

Investigation of sudden unexplained death: personal and family history. Colors correspond to the Class of Recommendation in Table 1 . SCD = sudden cardiac death.

Investigation of sudden unexplained death: the postmortem examination and imaging. Colors correspond to the Class of Recommendation in Table 1 .

Clinical and genetic evaluation after sudden death where a phenotype is known. Colors correspond to the Class of Recommendation in Table 1 . \*See Section 8.2 .

Investigation of sudden death: genetic evaluation where the phenotype is unknown. Colors correspond to the Class of Recommendation in Table 1 . SCD = sudden cardiac death.

Investigation of sudden death. Colors correspond to the Class of Recommendation in Table 1 . AED = automated external defibrillator; CIED = cardiovascular implantable electronic device; ECG = electrocardiogram; SCD = sudden cardiac death.

Investigation of sudden cardiac arrest (SCA) survivors: genetic evaluation. Colors correspond to the Class of Recommendation in Table 1 .

Investigation of sudden cardiac arrest survivors. Colors correspond to the Class of Recommendation in Table 1 . AED = automated external defibrillator; AF = atrial fibrillation; BBR-VT = bundle branch re-entry ventricular tachycardia; CIED = cardiovascular implantable electronic device; CMR = cardiac magnetic resonance imaging; CPVT = catecholaminergic polymorphic ventricular tachycardia; ECG = electrocardiogram; EP = electrophysiological; LQTS = long QT syndrome; SCA = sudden cardiac arrest; SVT = supraventricular tachycardia.

Investigation of the family affected by sudden cardiac arrest and/or sudden unexplained death when cause is identified or not identified. Colors correspond to the Class of Recommendation in Table 1 . CMR = cardiac magnetic resonance imaging; ECG = electrocardiogram; SCA = sudden cardiac arrest; SCD = sudden cardiac death; SUD = sudden unexplained death.

ACC/AHA recommendation system: Applying Class of Recommendation and Level of Evidence to clinical strategies, interventions, treatments, and diagnostic testing in patient care \*

COR and LOE are determined independently (any COR may be paired with any LOE).

A recommendation with LOE C does not imply that the recommendation is weak. Many important clinical questions addressed in guidelines do not lend themselves to clinical trials. Although RCTs are unavailable, there may be a very clear clinical consensus that a particular test or therapy is useful or effective.

The outcome or result of the intervention should be specified (an improved clinical outcome or increased diagnostic accuracy or incremental prognostic information).

For comparative-effectiveness recommendations (COR 1 and 2a; LOE A and B only), studies that support the use of comparator verbs should involve direct comparisons of the treatments or strategies being evaluated.

The method of assessing quality is evolving, including the application of standardized, widely-used, and preferably validated evidence grading tools; and for systematic reviews, the incorporation of an Evidence Review Committee.

COR indicates Class of Recommendation; EO, expert opinion; LD, limited data; LOE, Level of Evidence; NR, nonrandomized; R, randomized; and RCT, randomized controlled trial.

Key goals of genetic counseling following sudden cardiac death/resuscitated sudden cardiac arrest

Sudden cardiac death (SCD) is an important public health issue and warrants further study to better quantify its occurrence, its impact on society, and the opportunities for improving outcomes through public education and provision of automated external defibrillators and cardiopulmonary resuscitation (CPR) training.

For survivors of sudden cardiac arrest (SCA), victims of sudden unexplained death (SUD), and their relatives, a multidisciplinary team is central to thorough investigation, so as to maximize the opportunity to make a diagnosis. Where there has been an SCD or resuscitated SCA and a genetic cause is suspected, genetic testing and counseling is essential for families, to ensure that risks, benefits, results, and the clinical significance of genetic testing can be discussed.

The psychological care of families affected by SUD and survivors of SCA (and their families) should run in parallel with the investigation process. Assessment by professionals trained in psychological care should be offered, as well as grief counseling and peer support, where appropriate.

For the investigation of SUD, a detailed personal and family history is essential, with attention to sentinel symptoms during life such as syncope or seizures, witness accounts, premorbid investigations, and inspection of any cardiac rhythm monitoring around the time of death.

A comprehensive autopsy is an essential part of the investigation of SUD and should include collection and storage of tissue suitable for genetic analysis. When the autopsy suggests a possible genetic cause, or no cause and the heart is normal, referral to a multidisciplinary team for further investigation is indicated.

For victims of SCD or survivors of cardiac arrest where the phenotype is known, genetic testing of the proband focused on likely candidate genes, along with clinical evaluation of family members, aids in identifying family members with, or at risk of developing, the same condition.

For victims of SCD or survivors of cardiac arrest where the phenotype is not known, arrhythmia syndrome-focused genetic testing may help arrive at a secure diagnosis, whereas wider testing without careful consideration of the implications of indeterminate results by experienced clinicians may only serve to add uncertainty and lead to misinterpretation of results.

For the investigation of SCA survivors, essential inquiry includes detailed personal and family history, witness accounts, physical examination, multiple electrocardiograms (ECGs), and cardiac imaging. Ambulatory monitoring and/or provocative testing (exercise, pharmacological, and invasive electrophysiological) may provide additional useful information. A sample suitable for future DNA testing should be taken early in the patient's course and stored.

Genetic investigation of SCA survivors is best undertaken at a center with multidisciplinary care infrastructure and should focus on likely candidate genes known to be causally related to the suspected phenotype. In some cases, genetic evaluation without a suspected phenotype may be undertaken with appropriate genetic counseling, although genetic evaluation of patients with a known nongenetic cause of cardiac arrest is discouraged.

The investigation of the families of victims of SUD and survivors of SCA should include clinical and, if known, genetic cascade testing. If the cause of SUD (or rarely, SCA) is unknown, then clinical investigation of first-degree relatives may include physical examination, ECGs, cardiac imaging, ambulatory monitoring, and provocative testing (exercise, pharmacological, and rarely invasive electrophysiological) with multidisciplinary team supervision. Follow-up and periodic re-evaluation are important and are directed by initial findings.