



2025 Brazilian guidelines for the management of neuromyelitis optica spectrum disorder in adults and children

Samira Luisa dos Apóstolos-Pereira¹ Alfredo Damasceno² Ana Claudia Piccolo³
 Maria Fernanda Mendes⁴ Maria Lúcia Brito⁵ Regina Alvarenga⁶ Bruna Klein da Costa^{7,8}
 Douglas Kazutochi Sato^{9,10} Doralina Guimarães Brum¹¹ Elisa Mello¹² Felipe von Glehn¹³
 Giordani Passos^{14,15} Guilherme Diogo¹⁶ Herval Soares Neto^{17,18} Lis Campos^{19,20}
 Manuela Fragomeni²¹ Milena de Sales Pitombeira²² Rafael Paterno²³ Raquel Vassão²⁴
 Renata Barbosa Paolilo²⁵ Tarso Adoni¹ Thiago Fukuda²⁶ Vanessa Daccach²⁷
 Jefferson Becker⁷ Claudia Cristina Ferreira Vasconcelos⁶

¹ Universidade de São Paulo, Faculdade de Medicina, Hospital das Clínicas, Serviço de Neurologia, São Paulo SP, Brazil.

² Universidade Estadual de Campinas, Campinas SP, Brazil.

³ Universidade Municipal de São Caetano do Sul, São Paulo SP, Brazil.

⁴ Santa Casa de São Paulo, Faculdade de Ciências Médicas, Disciplina de Neurologia, São Paulo SP, Brazil.

⁵ Hospital da Restauração, Recife PE, Brazil.

⁶ Universidade Federal do Estado do Rio de Janeiro, Pós-Graduação Stricto Sensu em Neurologia, Rio de Janeiro RJ, Brazil.

⁷ Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre RS, Brazil.

⁸ Santa Casa de Porto Alegre, Hospital da Criança Santo Antônio, Porto Alegre RS, Brazil.

⁹ Hospital de Clínicas de Porto Alegre, Serviço de Neurologia, Porto Alegre RS, Brazil.

¹⁰ Universidade Federal do Rio Grande do Sul, Faculdade de Medicina, Departamento de Medicina Interna, Porto Alegre RS, Brazil.

¹¹ Universidade Estadual Paulista, Faculdade de Medicina de Botucatu, Botucatu SP, Brazil.

¹² Santa Casa de São Paulo, Faculdade de Ciências Médicas, São Paulo SP, Brazil.

¹³ Universidade de Brasília, Brasília DF, Brazil.

¹⁴ Pontifícia Universidade Católica do Rio Grande do Sul, Hospital São Lucas, Porto Alegre RS, Brazil.

¹⁵ Hospital Mãe de Deus, Porto Alegre RS, Brazil.

Address for correspondence Samira Luisa dos Apóstolos Pereira (email: samira.apostolos@hc.fm.usp.br)

¹⁶ Universidade de São Paulo, Faculdade de Medicina, Hospital das Clínicas, São Paulo SP, Brazil.

¹⁷ Hospital do Servidor Público Estadual de São Paulo, São Paulo SP, Brazil.

¹⁸ Hospital Israelita Albert Einstein, São Paulo SP, Brazil.

¹⁹ Universidade Federal de Sergipe, Aracaju SE, Brazil.

²⁰ Universidade Tiradentes, Aracaju SE, Brazil.

²¹ Hospital da Criança de Brasília, Brasília DF, Brazil.

²² Hospital Geral de Fortaleza, Divisão de Neurologia, Fortaleza CE, Brazil.

²³ Irmandade da Santa Casa de São Paulo, Serviço de Neurologia, São Paulo SP, Brazil.

²⁴ Santa Casa de Belo Horizonte, Clínica Neurológica, Belo Horizonte MG, Brazil.

²⁵ Universidade de São Paulo, Faculdade de Medicina, Hospital das Clínicas, Instituto da Criança, São Paulo SP, Brazil.

²⁶ Universidade Federal da Bahia, Hospital Universitário Professor Edgar Santos, Salvador BA, Brazil.

²⁷ Universidade de São Paulo, Faculdade de Medicina de Ribeirão Preto, Hospital das Clínicas, Ribeirão Preto SP, Brazil.

Arq. Neuro-Psiquiatr. 2025;83(11):s00451812471.

Abstract

Background Neuromyelitis optica spectrum disorder (NMOSD) is a debilitating recurrent inflammatory disease of the central nervous system. Establishing management guidelines is essential to optimize patient care in Brazil.

Objective To combine the Delphi and Grading of Recommendations Assessment, Development, and Evaluation (GRADE) approaches to validate evidence-based guideline statements for NMOSD treatment.

received
July 1, 2025
received in its final form
September 16, 2025
accepted
September 19, 2025

DOI <https://doi.org/10.1055/s-0045-1812471>.
ISSN 0004-282X.

Editor-in-Chief: Ayrton Roberto Massaro.
Associate Editor: Lívia Almeida Dutra.

© 2025. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by/4.0/>)
Thieme Revinter Publicações Ltda., Rua Rego Freitas, 175, loja 1, República, São Paulo, SP, CEP 01220-010, Brazil

Keywords

- Neuromyelitis Optica
- Aquaporin 4
- Disease Management
- Therapeutics
- Guideline
- Consensus

Methods These guidelines focused on providing recommendations for different scenarios: general management and diagnosis of NMOSD; acute and preventive treatments (including systemic immunosuppressants and anti-B cell, anti-interleukin-6, and anti-complement monoclonal antibodies); and therapeutic approaches in special groups (such as the pediatric population and pregnant women). A literature review based on the Population, Intervention, Comparator, and Outcome (PICO) framework was performed, and studies were evaluated using the GRADE system. An initial questionnaire containing 19 statements was sent to 44 specialists from all regions of Brazil.

Results Three voting rounds were necessary to reach consensus in all statements. After the first voting round, 17 statements reached consensus (level of agreement > 70%). Two statements failed to reach consensus and were, thus, revised. One of them was segmented into three different statements. After the second voting round, three out of the four revised statements reached consensus. Upon further revision, the last statement was again submitted to voting, reaching consensus in this third round. Overall, agreement was achieved on the 21 proposed statements.

Conclusion The primary objective in the management of NMOSD is to mitigate the severity of the attacks and prevent relapses, thereby minimizing the risk of irreversible neurological deficits. The statements in the current guideline offer evidence-based recommendations for such management within the Brazilian context.

INTRODUCTION

Neuromyelitis optica spectrum disorder (NMOSD) is a chronic, relapsing, and debilitating autoimmune inflammatory disease of the central nervous system. It is characterized by severe attacks of optic neuritis, longitudinally-extensive transverse myelitis, and demyelinating syndromes affecting the brain, diencephalon, and brainstem.¹ The disorder predominantly affects non-Caucasian female populations (female: male ratio = 9:1), and the average age of onset is around the fourth decade of life.^{2,3}

After the discovery of aquaporin-4-immunoglobulin G (AQP4-IgG), a key astrocytic water channel antibody, NMOSD was distinguished from multiple sclerosis (MS).⁴ In 2015, the International Panel for Neuromyelitis Optica (NMO) Diagnosis¹ (IPND) recommended AQP4-IgG detection as a fundamental diagnostic biomarker for NMOSD, since it is detected in approximately 75% of NMOSD patients.⁵ For those seronegative for AQP4-IgG, approximately 30 to 40% test positive for myelin oligodendrocyte glycoprotein-IgG (MOG-IgG) antibodies.⁶ Around 30% of clinically-diagnosed NMOSD patients remain seronegative for both AQP4- and MOG-IgG, the so-called *double-seronegative NMOSD* (DS-NMOSD).⁷ When characteristic clinical and neuroimaging findings are present and differential diagnoses are excluded, DS-NMOSD can be diagnosed using the 2015 IPND criteria.¹ Magnetic resonance imaging (MRI) assessment of the brain, optic nerve, and spinal cord is essential to confirm NMOSD-specific imaging and exclude alternative diagnoses such as MS and myelin oligodendrocyte glycoprotein antibody-associated disease (MOGAD).¹

Historically associated with high morbidity and mortality, NMOSD outcomes have improved due to early diagnosis and treatment, reducing mortality to approximately 10% in 10 years.^{8,9} Early intervention with acute attack management and immunotherapy has significantly reduced morbidity and improved long-term outcomes.¹⁰ However, irreversible disability remains a major concern due to severe attacks, leading to extensive testing, immediate treatment with high-dose steroids, plasmapheresis, and the frequent need for admission to the Intensive Care Unit.¹¹ Several recent randomized controlled trials^{12–15} (RCTs) have demonstrated the efficacy of novel monoclonal antibodies (mAbs) such as satralizumab, eculizumab, inebilizumab, and ravulizumab in the treatment of AQP4-IgG-positive NMOSD patients, leading to approvals by regulatory agencies worldwide. However, there are no direct clinical comparisons to establish the best order of treatment administration.¹⁶ Moreover, these drugs are currently not approved for patients living with DS-NMOSD.

Given the disease's complexity and variability, establishing consensus management guidelines is essential to optimize patient care in Brazil and globally.¹⁷ The Delphi method gathers expert opinions to establish guidelines, particularly for rare diseases lacking standardized treatments.¹⁶ The Grading of Recommendations, Assessment, Development, and Evaluation (GRADE) method evaluates the quality of evidence to support treatment decisions.¹⁸ We combined the Delphi¹⁶ and GRADE¹⁸ approaches to prioritize clinical questions and validate evidence-based guideline statements, thus ensuring best-practice recommendations for NMOSD treatment.

METHODS

A conventional Delphi methodological approach¹⁹ was employed to develop the current guideline. The qualitative study was conducted from March 2023 to October 2024 by the Brazilian Academy of Neurology (Academia Brasileira de Neurologia, ABN, in Portuguese).


A steering committee was formed, composed of six experts (ACP, AD, CCFV, JB, MFM, and SLAP) with experience in the diagnosis and treatment of NMOSD (►Figure 1). The present guideline focused on providing recommendations for different scenarios: general management and diagnosis of NMOSD; acute and preventive treatments, including systemic immunosuppressive therapy (IST), and anti-B cell, anti-interleukin-6 (anti-IL-6), and anti-complement mAbs; and therapeutic approaches in special groups (such as the pediatric population and pregnant women).

A rapid systematic literature review (RSLR) was conducted to develop statements related to each aspect. To this end, six groups were formed, each consisting of a steering committee member and three specialists with expertise in NMOSD²⁰ (►Table 1, ►Figure 1). A structured search in the

PubMed, Embase, LILACS, and Google Scholar databases was performed to recover observational studies and double-blinded RCTs written in English, Spanish, and Portuguese, and published from 1999 to 2024. The search strategy was based on the Patient, Intervention, Comparator, Outcome (PICO) framework, as described elsewhere,²¹ and developed using the strategy described in ►Table 2. All studies considered for full reading were evaluated for the quality of evidence using the GRADE system, and classified accordingly into four levels: high, moderate, low, and very low.¹⁸

Generation of statements

The statements have four main characteristics: anonymity, interaction, controlled feedback, and statistical group response. They were developed based on the quality classification of the evidence and sent for critical review to a senior advisory group consisting of four neurologists with extensive clinical experience and scientific knowledge about NMOSD. Aspects such as risk of bias or limitations, publication bias, imprecision, inconsistency, and indirect evidence were analyzed in each selected study for the final quality classification of the evidence (online Supplementary Material). The



Abbreviations: GRADE, Grading of Recommendations, Assessment, Development, and Evaluation; NMOSD, neuromyelitis optica spectrum disorder.

Figure 1 Algorithm of the Delphi method steps used to prepare the 2025 Brazilian NMOSD Consensus.

Table 1 Eligibility criteria and credentials of the expert panel members

1. ≥ 10 years of specialty in neurology.
2. One of the following conditions:
 - a. Participation in reference services in a teaching hospital specialized in people living with neuromyelitis optica spectrum disorder (NMOSD; pwNMOSD);
 - b. Leading national cohort studies to investigate NMOSD; and
 - c. Experience in caring for at least 50 pwNMOSD in the last year.

comments and suggestions from the senior group were sent back to the steering committee, and an initial questionnaire containing 19 statements was organized in an electronic form (Google Forms, Alphabet Inc.), with 5 response options categorized by the Likert scale (*strongly disagree*, *disagree*, *neither agree nor disagree*, *agree*, and *strongly agree*). Expert opinions were added at the end of the analysis, after the voting rounds, considering the assessment of peculiarities according to the regions of Brazil.

Voting rounds

An online questionnaire (► **Supplementary Material**, available at <https://www.arquivosdeneuropsiquiatria.org/wp-content/uploads/2025/09/ANP-2025.0217-Supplementary-Material.docx>) was sent to 44 main specialists from all regions of Brazil, selected based on their requirements (► **Table 1**) to be part of the voting expert panel and to assist in the development of the final consensus. The steering committee members and the senior specialists did not participate in the voting rounds.

To achieve maximum agreement on each assertion, a maximum of four voting rounds was stipulated; participation in the first two rounds was also a requirement. Each response submission was extracted from Google Forms and transferred to Excel (Microsoft Corp.) spreadsheets, with

anonymization concerning the voting specialists, and qualitative and quantitative analyses were conducted. For each voting round, the statements that did not reach a predefined consensus threshold ($> 70\%$ of agreement) were revised and reformulated by the steering committee. The revised statements were sent back to the same voting panel, including feedback reports from the qualitative and quantitative analyses. The algorithm depicted in ► **Figure 1** summarizes the steps of the Delphi methodology followed. Standard protocol approvals, registrations, and patient consent were not required for this study. Additional data can be assessed in the Supplementary Material.

RESULTS

Overview

The targeted literature search yielded 21 articles on acute management of NMOSD attacks (out of 54 articles identified), 31 articles on oral immunosuppressants (out of 107 articles identified), 34 articles on B-cell depleting therapies (out of 565 articles identified), 9 articles on IL-6 receptor blockade (out of 25 articles identified), 8 articles on complement component 5 (C5) inhibitors (out of 149 articles identified), and 27 articles (out of 147 articles identified) on special situations (pediatric NMOSD, pregnancy, and breastfeeding). Following the GRADE

Table 2 PICO strategy for the 2025 Brazilian consensus on the treatment of neuromyelitis optica spectrum disorder (NMOSD) in the adult and pediatric populations

Acronym	Description
P - study population	All people living with NMOSD (pwNMOSD), according to diagnostic criteria published since 1999, ⁹ including: ● Adult patients (aged ≥ 18 years); ● Pediatric patients (aged ≤ 17 years); ● Pregnant women with NMOSD, regardless of the aquaporin-4-immunoglobulin G (AQP4-IgG) serostatus.
I - intervention	● Intervention for diagnosis, using the descriptor <i>AQP4-IgG antibody</i>; ● Intervention in the acute phase, using the descriptors <i>oral and/or intravenous corticosteroids</i>, <i>immunoglobulin</i>, and <i>therapeutic apheresis</i>; and ● Prophylactic intervention to prevent new attacks, using the following descriptors: <i>oral corticosteroids</i>, <i>azathioprine</i>, <i>mycophenolate mofetil</i>, <i>methotrexate</i>, <i>cyclophosphamide</i>, <i>rituximab</i>, <i>tocilizumab</i>, <i>satralizumab</i>, <i>inebilizumab</i>, <i>eculizumab</i>, or <i>ravulizumab</i>.
C - comparator	● Placebo; ● Corticosteroids; ● Systemic immunosuppressants; and ● Rituximab.
O - Outcome	● Efficacy outcomes: evaluation of the time until the first relapse, reduction in the annualized relapse rate, and disability progression assessed by the Expanded Disability Status Scale; ● Safety and tolerability; and ● Outcomes related to the route and frequency of treatment administration.

system, all articles included were evaluated (online Supplementary Material). Based on this evidence, 19 draft statements were developed.

Voting rounds

Three out of the four previously-stipulated voting rounds were necessary to reach consensus in all statements. All voting members ($n=44$) participated in all 3 rounds and voted in all statements. After the first voting round, 17 out of the 19 statements reached consensus (level of agreement $>70\%$). Two statements failed to reach consensus (levels of agreement of 65.9% and 63.6%). Therefore, these statements were revised, and one of them was segmented into three different statements. They were then submitted to a second voting round, upon which three out of the four revised statements reached consensus. The statement that did not reach consensus (level of agreement of 59.1%) was further revised and resubmitted to voting, reaching consensus in this third round (level of agreement of 70.5%).

Overall results

There was agreement on 21 consensus statements overall. These are discussed in more detail in the sections that follow, which are divided according to the topics of the targeted literature review.

General approach to patients with NMOSD diagnosis (► Table 3)


Statement 1 reached consensus in round 1 of voting (level of agreement of 95.4%), recommending that patients with clinical presentation suggestive of NMOSD must be tested for AQP4-IgG and, when negative, tested for MOG-IgG. Real-world data²² provide strong evidence that serum AQP4-IgG predicts NMOSD relapses after an initial attack. The 2015 IPND criteria improved NMOSD diagnosis by 40%, reclassifying 60% of cases and reducing the median time to diagnosis by 1 to 18 months.^{1,23} The detection of AQP4-IgG should be performed using appropriate methodologies, most notably cell-based assays (CBAs) employing either fixed or live cells, in serum samples ideally obtained prior to the initiation of

immunotherapy.¹ Live CBAs have consistently demonstrated¹ superior diagnostic performance compared to fixed-cell techniques. However, their implementation frequently relies on in-house protocols that can vary across laboratories, posing challenges for standardization and reproducibility—issues that are particularly pronounced in low-income countries, where technical resources and expertise may be limited.²⁴ Consequently, despite their lower sensitivity, fixed CBAs have been more broadly adopted in the routine clinical practice due to their greater availability and feasibility.²⁴ Despite optimal methods and timing of testing, approximately 30% of the patients with clinical presentation consistent with NMOSD lack AQP4-IgG (seronegative NMOSD).²⁵ In a significant proportion of these cases, MOG-IgG is detected.⁶ Therefore, it is strongly recommended that all patients under investigation for NMOSD who tested negative for AQP4-IgG should be investigated for MOG antibodies in serum samples using CBA to ensure accurate diagnosis and appropriate management.²⁶ Accurate reporting of MOG-IgG testing requires specifying both the antibody titer and the assay matrix. Titer levels provide valuable quantitative insights that can correlate with disease activity and prognosis, with higher titers more suggestive of pathogenic relevance. Early serum sampling is fundamental for a reliable MOG-IgG diagnosis, as antibody levels may decline over time.²⁵ While serum testing is generally preferred for its higher sensitivity, cerebrospinal fluid (CSF) analysis may offer additional diagnostic value in select cases, particularly in scenarios of encephalitis and possibly myelitis. Conversely, the usefulness of CSF testing in optic neuritis remains uncertain. Combining detailed titration data with information on assay methodology and sample source enhances diagnostic precision and clinical interpretation.^{25,26} (► Figure 2). Early antibody detection is essential for accurate diagnosis and the implementation of appropriate therapeutic strategies,²⁶ mainly in patients exhibiting optic neuritis and myelitis, since NMOSD and MOGAD have different immunological mechanisms and require different therapeutic approaches (► Figure 3). Despite advanced detection methods, including live CBA and flow cytometry analysis, approximately 25% of

Table 3 Brazilian Delphi consensus – statements on the general management and diagnosis of NMOSD

Consensus statement	Level of agreement
General approach to a patient with a NMOSD diagnosis	
1. We recommend that all patients under investigation for NMOSD be tested for AQP4-IgG antibodies. If negative, then MOG-IgG antibodies should be tested.	95.4%
2. We suggest that the treatment of pwNMOSD should be conducted within a specialized reference center.	79.5%
3. We recommend that: A. Vaccination status should be updated prior to the initiation of preventive therapies in all pwNMOSD; and B. Ongoing monitoring through clinical evaluation and laboratory tests (when indicated) should be performed in NMOSD patients receiving preventive therapy to assess treatment response, adverse events, and manage therapeutic adjustments when necessary.	100%

Abbreviations: AQP4-IgG, aquaporin-4-immunoglobulin G; MOG-IgG, myelin oligodendrocyte glycoprotein-immunoglobulin G; NMOSD, neuromyelitis optica spectrum disorder; pwNMOSD, people living with neuromyelitis optica spectrum disorder.




Abbreviations: AQP4-IgG, aquaporin-4-immunoglobulin G; CBA, cell-based assay; MOG-IgG, myelin oligodendrocyte glycoprotein-immunoglobulin G; NMOSD, neuromyelitis optica spectrum disorder; MOGAD, MOG-IgG-associated disease.

Figure 2 Diagnosis of NMOSD based on current NMOSD and MOGAD guidelines.

the patients clinically diagnosed with NMOSD remain seronegative for both AQP4 and MOG autoantibodies (DS-NMOSD).⁷

Statement 2 reached consensus (level of agreement of 79.5%) in round 1 of voting, recommending that patients diagnosed with NMOSD receive treatment within specialized reference centers dedicated to this condition. Qualitative real-world data²⁷ demonstrate that the journey of all people living with NMOSD (pwNMOSD) is challenging due to diagnostic errors and misinformation. Although only qualitative evidence is available, the panel postulated that receiving care in specialized centers may increase the accuracy of diagnosis and the timeliness of appropriate treatment for pwNMOSD, given the complexities associated with the differential diagnosis and the accurate application of diagnostic criteria in scenarios involving diverse populations.^{23,24} The strategy of treatment in pwNMOSD requires assessment of infection risk and up-to-date vaccination status, as well as of challenges related to selection, accessibility, and management of preventive therapies (such as serious adverse events).^{28,29} Infusion reactions and severe adverse events may occur in patients receiving intravenous MAb therapies, such as rituximab, inebilizumab, eculizumab, ravulizumab, and tocilizumab. It is recommended that these agents be administered in

specialized neuroimmunology infusion centers staffed by trained personnel capable of managing infusion-related reactions and monitoring for therapy-specific adverse effects. Implementation of premedication protocols and continuous monitoring during and after infusion are advised to mitigate reaction risk. Additionally, comprehensive preventive strategies should be instituted prior to treatment initiation and maintained throughout therapy. The panel suggested that vigilant clinical monitoring of infections (such as respiratory and urinary infections) and routine laboratory evaluation, including assessment of liver enzymes, complete blood count (including neutrophil and platelet counts), and immunoglobulin levels, are essential to detect cytopenia and hypogammaglobulinemia early, thereby optimizing patient safety and treatment outcomes.^{24,27–29} In specific clinical scenarios, such as concomitant infections or unexpected relapses, adjustments to the dosing strategy of B-cell-depleting therapies may be required. Rituximab, one of the most widely used and cost-effective immunobiological agents for NMOSD, can be administered at fixed intervals (such as every 6 months) or tailored according to lymphocyte subpopulation counts. Notably, early or unexpected relapses in pwNMOSD receiving rituximab have been associated with paradoxical increases



Abbreviations: Anti-AQP4, aquaporin-4-immunoglobulin G; anti-MOG, myelin oligodendrocyte glycoprotein antibody; NMOSD, neuromyelitis optica spectrum disorder; IVMP, intravenous methylprednisolone, IVIG, intravenous immunoglobulin; MOGAD, MOG-IgG myelin oligodendrocyte glycoprotein-immunoglobulin G-associated disease.

Figure 3 Diagnosis and treatment of NMOSD based on the AQP4-IgG serostatus, according to the 2025 Brazilian consensus. Treatment recommendations are depicted in the figure. Eculizumab is not included, since its use is not currently approved by the Brazilian Health Regulatory Agency (Agência Nacional de Vigilância Sanitária, ANVISA, in Portuguese) for NMOSD treatment in Brazil. The blue lines indicate the consensus for preventive treatment recommendations for AQP4-IgG-positive patients. The red dashed lines indicate the consensus for AQP4-IgG-negative patients. Considerations regarding the selection among available monoclonal antibodies for preventive treatment are presented in the green dotted box.

in B-cell-activating factor (BAFF) or premature B-cell repopulation. Monitoring peripheral B-cell counts—particularly those of CD19+ cells—has been shown to predict relapse risk, underscoring the importance of close specialist supervision to guide timely retreatment and optimize dosing schedules. Moreover, genetic variations in Fcγ receptors, particularly *FCGR3A* polymorphisms, have been reported to influence the clinical response to rituximab, suggesting a role of host immunogenetic factors in treatment outcomes.³⁰ Specifically, lipid (such as cholesterol and triglyceride) levels influenced by the IL-6 blockade must be evaluated, and fibrinogen levels, when deemed necessary.³¹

Statement 3 reached consensus (level of agreement of 100% in round 1 of voting), recommending that vaccination status should be up to date before the initiation of preventive therapies. Before starting on a lifelong preventive treatment, the vaccination status must be reviewed and the patient should be screened for infections, including those by the hepatitis B virus (HBV) and tuberculosis. Vaccines are generally safe for pwNMOSD, including those for coronavirus disease 2019 (COVID-19).³² For patients receiving B-cell-depleting therapies, it is recommended to vaccinate before the initiation of these therapies, particularly for HBV, pneumococcal, influenza, and recombinant adjuvanted zoster vaccines.³² Live vaccines are contraindicated during IST.³³ In addition, monitoring treatment response and adverse events through clinical evaluation and laboratory exams is mandatory in pwNMOSD under preventive therapy.

Acute management of NMOSD attacks

The panel discussed the acute management of NMOSD attacks (–Table 4) based on 3 approaches: 1 g/day of intravenous methylprednisolone (IVMP), therapeutic plasma exchange (TPE), and intravenous immunoglobulin (IVIG). Exacerbations in NMOSD often cause severe disability; thus, prompt, intensive treatment is crucial to suppress relapses and limit CNS damage for better recovery.^{34–36}

Statement 4 reached consensus (97.7% of agreement in round 1 of voting), recommending that NMOSD attacks must be treated with high doses of IVMP as early as possible. regarding the acute-phase treatment, eight studies were identified: two prospective^{36,37} and six retrospective studies.^{34,35,38–41} The main therapeutic strategies evaluated included high-dose IVMP, either alone or in combination with TPE, as well as IVIG administered alone or following IVMP in steroid-refractory cases. The panel members concurred that IVMP is the current standard to treat acute NMOSD attacks and is strongly recommended despite the low grade of evidence. There is a paucity of high-quality studies, and only one study⁴¹ investigated the effectiveness of IVMP as a monotherapy in the acute phase. This recommendation is based on its widespread availability, decades of clinical experience, reduced need for monitoring, and potential to accelerate recovery, particularly when administered early at a dose of 1 g for 3 to 7 consecutive days.^{34,41} In the clinical practice, after completing corticosteroid pulse therapy, it is common to maintain 1 mg/kg/day of oral corticosteroids,

Table 4 Brazilian Delphi consensus – statements on the acute management of NMOSD attacks

Consensus statement	Level of agreement
Acute management of NMOSD attacks	
4. We recommend administering high-dose intravenous methylprednisolone (IVMP; 1g/day for ≥ 5 days) as early as possible for the treatment of NMOSD attacks, regardless of the AQP4-IgG serostatus, to improve outcomes related to attack severity.	97.7%
5. We recommend initiating therapeutic plasma exchange (TPE; for ≥ 5 cycles) as early as possible for the treatment of NMOSD attacks, either alone or associated with high-dose IVMP, regardless of the AQP4-IgG serostatus, to reduce functional disability and increase the likelihood of recovery.	86.4%
6. We do not recommend the use of intravenous immunoglobulin (IVIG) for the treatment of NMOSD attacks, regardless of the AQP4-IgG serostatus, except in patients seropositive for MOG-IgG antibodies.	75%

Abbreviations: AQP4-IgG, aquaporin-4-immunoglobulin G; MOG-IgG, myelin oligodendrocyte glycoprotein-immunoglobulin G; NMOSD, neuromyelitis optica spectrum disorder.

followed by a gradual, individualized taper. This may vary according to the mechanism of action of the selected preventive therapy. Given the heterogeneity in the clinical approaches, definitive recommendations cannot yet be established. Recovery after an attack in pwNMOSD treated with steroids depends on factors such as early treatment, age, prior immunosuppressant use, elevated CSF protein, and MRI with gadolinium enhancement.⁴² Although IVMP is standard for relapses, full recovery is uncommon, and many attacks show limited or no response to therapy.^{34,41}

Statement 5 reached consensus (86.4% of agreement in round 1 of voting), recommending that NMOSD attacks must be treated with TPE (for ≥ 5 cycles) associated or not with high doses of IVMP and as early as possible (regardless of the AQP4-IgG serostatus). Therapeutic plasma exchange for NMOSD attacks is supported by low-to-moderate quality of evidence under the GRADE framework, but has a strong clinical rationale and observed benefits, leading to a practical recommendation favoring its use. It can be performed concurrently with IVMP or early, as the first-line therapy in severe cases (→ **Table 5**). When the response to pulse therapy is incomplete or absent, TPE is recommended for use on alternate days for 5 to 7 cycles, preferably within 5 days of the onset of NMOSD relapse.^{34–37} A retrospective study³⁴ reviewed 83 cases of NMOSD attacks on patients treated with IVMP alone versus IVMP plus TPE; A reduction in disability (according to the Expanded Disability Status Scale, EDSS) at follow-up occurred in 65% of the combined treatment group, but only in 35% of the group receiving IVMP alone. Two large retrospective cohort studies^{38,41} on TPE efficacy in functional recovery concluded that early inter-

vention yields better outcomes, with fewer than 5% of pwNMOSD presenting improvement when TPE was delayed for 3 weeks after the attack.³⁸ A retrospective cohort⁴¹ of 207 attacks in 105 pwNMOSD identified early TPE initiation and its use as first-line therapy as strong predictors of complete remission. Although TPE appears to be an effective approach to treat steroid-resistant NMOSD attacks and an early intervention improves outcomes, the optimal timing of initiation remains a critical, unresolved issue.^{39,43} The use of TPE without steroids was evaluated in a prospective observational study of 30 pwNMOSD,³⁷ showing functional improvement in 73.3% of the patients and a significant correlation between recovery and early TPE initiation. The AQP4-IgG status did not affect the outcomes. Current evidence supports a “the sooner, the better” approach for both high-dose IVMP and TPE. However, access to tertiary centers offering TPE remains a major barrier, especially in low-income regions. In practice, delayed healthcare access can hinder timely treatment, potentially worsening prognosis. Of note, immunoadsorption, a distinct therapeutic apheresis technique, is limited to ultraspecialized centers and was not considered in the present consensus.

Statement 6 reached consensus (75% of agreement in round 1 of voting) on the assumption that IVIG is not indicated for acute exacerbations of NMOSD. This recommendation aligns with the absence of high-quality evidence per the GRADE to support IVIG use. Alone or in combination, IVIG has not shown sufficient benefit in treating NMOSD relapses, regardless of the AQP4-IgG status. The data available consistently demonstrate that IVIG is less effective than steroids or TPE.⁴⁴ In steroid-resistant attacks, IVIG is inferior

Table 5 Severe relapse criteria in pwNMOSD^{43,115}

Severe relapses in pwNMOSD include:

1. Optic neuritis: bilateral optic neuritis or severe monocular visual loss (visual acuity worse than 20/200); and
2. Acute myelitis: clinical presentation consistent with transverse/complete myelitis, major motor deficits (muscle strength lower than II out of V on the MRC scale), or bladder dysfunction.

Abbreviations: MRC, Medical Research Council Scale for muscle strength; pwNMOSD, people living with neuromyelitis optica spectrum disorder.

Table 6 Brazilian Delphi consensus – Statements on the preventive management of NMOSD attacks in AQP4-IgG-positive patients

Consensus statement	Level of agreement
Prevention of NMOSD attacks in AQP4-IgG-seropositive patients	
7. We recommend rituximab monotherapy, initiated at diagnosis, after the first attack, or following relapse due to the failure of previous treatments, for the prevention of attacks and disability in adults and children with NMOSD, regardless of the AQP4-IgG serostatus.	77.3%
8. We recommend treatment with eculizumab, inebilizumab, ravulizumab, or satralizumab, as monotherapy, at diagnosis, after the first attack, or following relapse due to the failure of previous treatments, for the prevention of attacks and disability in adults (aged ≥ 18 years) with AQP4-IgG-positive NMOSD.	95.5%
9. We recommend that the choice among eculizumab, inebilizumab, ravulizumab, or satralizumab consider factors such as drug availability, comorbidities, other autoimmune disorders, and patient preference, since efficacy and safety have not been compared in head-to-head studies in AQP4-IgG-seropositive pwNMOSD.	97.7%
10. We do not recommend the combined therapy with two or more monoclonal antibodies (e.g., eculizumab, inebilizumab, ravulizumab, rituximab, satralizumab, or tocilizumab) in AQP4-IgG-seropositive NMOSD patients.	95.5%
11. We do not recommend azathioprine, cyclophosphamide, oral steroids, methotrexate, or mycophenolate mofetil, as monotherapy, for first-line preventive therapy in adults with AQP4-IgG-positive NMOSD.	75%*

Abbreviations: AQP4-IgG, aquaporin-4-immunoglobulin G; NMOSD, neuromyelitis optica spectrum disorder; pwNMOSD, people living with neuromyelitis optica spectrum disorder.

Note: *Agreement reached in round 2 of voting.

to TPE in reducing pathogenic antibody levels and improving disability outcomes.⁴⁵ Based on current evidence, the panel agreed that the most effective strategy for severe NMOSD relapses remains the combined use of high-dose corticosteroids and plasmapheresis.

NMOSD preventive treatment

The panel addressed preventive treatment across 18 scenarios: 6 for AQP4-IgG-positive NMOSD, 5 concerning DS-NMOSD patients meeting the 2015 IPND criteria, 1 for patients stable on long-term immunosuppression, 3 for pediatric cases, and 3 for women during pregnancy or breastfeeding. The preventive treatments were categorized by AQP4-IgG status, as robust evidence¹ supports NMOSD AQP4-IgG-positive as a distinct disease entity (► **Figure 3**), and treatment recommendations are now supported by high-level evidence under the GRADE framework (► **Table 6**). Patients fulfilling clinical and neuroimaging criteria for NMOSD but testing negative for AQP4-IgG should be re-evaluated for differential diagnoses, including MS and MOGAD (► **Figures 2–3**). Our recommendations for AQP4-IgG-negative NMOSD apply only to those who do not meet the latest MS criteria and are also negative for MOG-IgG. Treatment in this subgroup must be individualized due to diagnostic uncertainty, absence of reliable biomarkers for diagnosis, and a scarce literature on treatment (► **Tables 7–8**). Recommendations for special populations, irrespective of the AQP4-IgG status—such as pediatric patients and pregnant or breastfeeding women—are presented in ► **Tables 9–10**. These guidelines do not apply to MOG-IgG-positive individuals who are classified under MOGAD, a distinct clinical entity.

Preventive treatment in AQP4-IgG-positive NMOSD

Statements 7, 8, 9, 10, and 11 were developed to guide the preventive treatment in AQP4-IgG-positive NMOSD, and they are presented in ► **Table 6**.

Statement 7 reached the consensus (77.3% of agreement in round 1 of voting) that rituximab treatment is recommended at the moment the diagnosis is established, after the first attack, and after relapse due to the failure of previous treatments, for the prevention of attacks and disability in adult and children diagnosed with NMOSD, regardless of the AQP4-IgG serostatus. This recommendation is supported by a placebo-controlled randomized trial, called “A multi-center, randomized, double-blind, placebo-controlled trial to determine the efficacy of rituximab against a relapse of neuromyelitis optica spectrum disorders with anti-aquaporin 4 antibody” (RIN-1),⁴⁶ which enrolled 38 adults with AQP4-IgG-positive NMOSD, randomized 1:1 to receive rituximab (375 mg/m² weekly for 4 weeks, followed by 2 g every 6 months) or placebo. While 37% of the placebo group experienced relapses, none in the rituximab group relapsed. This trial provides high-level evidence, according to the GRADE framework, that rituximab significantly reduces relapse risk in adults with AQP4-IgG-positive NMOSD. Additional evidence of low-to-moderate quality from observational and open-label studies supports its effectiveness across other outcomes and subgroups (online Supplementary Material). For instance, the study called “An open-label extension study following the RIN-1 study to investigate the efficacy of rituximab against a relapse of neuromyelitis optica” (RIN-2),⁴⁷ which compared rituximab to the placebo arm of RIN-1, reported an annualized relapse rate (ARR) of 0.035 with rituximab versus 0.321 in the placebo group, with stable

Table 7 Brazilian Delphi Consensus – statements on preventive management of NMOSD attacks in AQP4-IgG-negative patients

Consensus statement	Level of agreement
Prevention of NMOSD attacks in AQP4-IgG-seronegative patients	
7. We recommend monotherapy with rituximab at diagnosis, after the first attack, or following relapse due to the failure of previous treatments, for the prevention of relapses and disability in adults or children with NMOSD, regardless of the AQP4-IgG serostatus.	77.3%
12. We suggest that azathioprine or mycophenolate mofetil may be used as initial preventive therapy in AQP4-IgG-negative patients (children and adults), only in the initial phase of the disease and when other options are not available. These oral therapies should be used with caution, given the absence of high-quality evidence supporting their application.	84.1%
13. We suggest tocilizumab monotherapy at diagnosis, after the first attack, or following relapse due to the failure of previous treatments, for the prevention of attacks and disability in AQP4-IgG-negative adults with NMOSD.	70.5%*
14. We do not recommend, at this time, treatment with other B-cell-depleting therapies (e.g., ocrelizumab, ofatumumab, or ublituximab) for pwNMOSD, given the lack of supporting evidence.	97.7%

Abbreviations: AQP4-IgG, aquaporin-4-immunoglobulin G; NMOSD, neuromyelitis optica spectrum disorder; pwNMOSD, people living with neuromyelitis optica spectrum disorder.

Note: *Agreement reached in round 3 of voting.

Table 8 Brazilian Delphi Consensus – statements on long-term preventive management of NMOSD attacks in clinically-stable patients

Consensus statement	Level of agreement
Long-term prevention of NMOSD attacks in clinically-stable patients	
15. We suggest that in all pwNMOSD (adults or children) who are relapse-free and without adverse events, the use of azathioprine, mycophenolate mofetil, or rituximab as preventive therapy may be maintained, regardless of the AQP4-IgG serostatus.	90.1%

Abbreviations: AQP4-IgG, aquaporin-4-immunoglobulin G; NMOSD, neuromyelitis optica spectrum disorder; pwNMOSD, people living with neuromyelitis optica spectrum disorder.

EDSS scores throughout the study period. Regarding safety, adverse events of grades ≥ 3 occurred in 4 to 6% of the patients in the RIN-1 and RIN-2 trials. A meta-analysis⁴⁸ of 36 studies (n = 1,542) reported infrequent, mostly mild-to-moderate adverse events, with infusion reactions and infec-

tions being the most common. These risks can be mitigated through strategies such as premedication, infusion rate adjustments, and vaccination protocols.

Statement 8 reached consensus (95.5% of agreement in round 1 of voting), recommending the treatment with

Table 9 Brazilian Delphi consensus – Statements on preventive management of NMOSD attacks in children – special situations

Consensus statement	Level of agreement
Preventive treatment in children with NMOSD	
7. We recommend rituximab as monotherapy at diagnosis, after the first attack, or after relapse due to the failure of previous treatments, for the prevention of relapses and disability in adults or children with NMOSD, regardless of the AQP4-IgG serostatus.	77.3%
16. We recommend satralizumab, at diagnosis, after the first attack, or after relapses due to the failure of previous treatments, as monotherapy or combined with oral immunosuppressants, for the prevention of relapses and disability in adolescents aged ≥ 12 years with NMOSD who are AQP4-IgG seropositive.	93.2%
17. We suggest that preventive therapy with azathioprine or mycophenolate mofetil be used with caution in children with NMOSD, regardless of the AQP4-IgG serostatus, and only when rituximab and satralizumab are contraindicated.	86.4%
18. We do not recommend monotherapy with cyclophosphamide, oral steroids, or methotrexate for relapse prevention in children with NMOSD, regardless of the AQP4-IgG serostatus.	81.8%

Abbreviations: AQP4-IgG, aquaporin-4-immunoglobulin G; NMOSD, neuromyelitis optica spectrum disorder.

Table 10 Brazilian Delphi consensus – statements on preventive management of NMOSD attacks according to AQP4-IgG status: special situations

Consensus statement	Level of agreement
Preventive treatment in NMOSD during pregnancy and breastfeeding	
19. We suggest the use of rituximab during pregnancy and breastfeeding in women with NMOSD, regardless of the AQP4-IgG serostatus, when the patient provides informed consent and is closely monitored in collaboration with the obstetrics team.	84.1%
20. We recommend that treatment with cyclophosphamide, methotrexate, or mycophenolate mofetil be avoided during pregnancy and breastfeeding in women with NMOSD, regardless of the AQP4-IgG serostatus, due to teratogenic and toxicity risks.	100%
21. We do not suggest the use of eculizumab, inebilizumab, ravulizumab, or satralizumab during pregnancy and breastfeeding in women with NMOSD, given the current lack of evidence to support their application in this context.	86.4%

Abbreviations: AQP4-IgG, aquaporin-4-immunoglobulin G; NMOSD, neuromyelitis optica spectrum disorder.

eculizumab, inebilizumab, ravulizumab, or satralizumab, as a monotherapy in AQP4-IgG-seropositive pwnNMOSD at the moment the diagnosis is established, after the first attack, and after relapse due to the failure of previous treatments, for the prevention of attacks and disability in adults (aged ≥ 18 years). This statement is supported by RCTs for eculizumab,^{14,49} inebilizumab, satralizumab,^{15,31,50} and ravulizumab,¹³ according to the GRADE framework (evidence of moderate-to-high quality).

The recommendation of eculizumab to prevent relapses in AQP4-IgG-positive NMOSD is based on the phase-3 trial called “A Randomized Controlled Trial of Eculizumab in AQP4 Antibody-positive Participants With NMO” (PREVENT),¹⁴ which enrolled 143 adult. After about 211 weeks, the relapse rate was of 3% in the eculizumab group compared to 43% in the placebo group (hazard ratio [HR] = 0.06; 95% CI = 0.02–0.20; $p < 0.001$). As a secondary endpoint, eculizumab monotherapy was associated with a significantly lower rate of worsening of the disability status according to the EDSS (4.8% versus 38.5% in the placebo group).⁴⁹

The recommendation of ravulizumab is supported by the open-label trial with an external comparator called “A Phase 3, External Placebo-Controlled, Open-Label, Multicenter Study to Evaluate the Efficacy and Safety of Ravulizumab in Adult Patients With Neuromyelitis Optica Spectrum Disorder (NMOSD)” (CHAMPION-NMOSD).¹³ A total of 58 AQP4-IgG-positive NMOSD adults received IV loading doses of ravulizumab ranging from 2,400 to 3,000 mg, followed by 3,000 to 3,600 mg every 8 weeks. No relapses occurred during the follow-up, although the single-arm design and use of an external comparator are study limitations.

The recommendation of satralizumab (120 mg via subcutaneous [SC] injection every 4 weeks [Q4w]; as monotherapy or in combination with oral ISTs) for relapse prevention in AQP4-IgG-seropositive NMOSD patients was supported by 2 RCTs and one observational study (moderate-quality evidence for ARR reduction and time until first relapse).^{15,31,50} The analysis of disability (EDSS) generally were included as secondary endpoints in the studies evaluated.

The recommendation of inebilizumab is supported by a single RCT (high-quality evidence): “A Double-masked, Pla-

cebo-controlled Study With Open-label Period to Evaluate the Efficacy and Safety of MEDI-551 in Adult Subjects With Neuromyelitis Optica and Neuromyelitis Optica Spectrum Disorders”¹² (N-Momentum), a phase-2/3 RCT that enrolled 230 patients, 93% of whom were AQP4-IgG-seropositive. The patients received 2 IV doses of inebilizumab (300 mg on days 1 and 15) or placebo. For the primary outcome—time until first relapse—inebilizumab demonstrated a 73% relative risk reduction compared to placebo (12% versus 39%; HR = 0.27; 95%CI = 0.15–0.50; $p < 0.0001$). Disability worsening, measured by the EDSS, was assessed as a secondary endpoint and showed less progression in the inebilizumab group compared to the placebo group. Additional long-term data from the open-label extension of N-MOMentum⁵¹ showed that 77% of the patients remained relapse-free after 4 years of continuous treatment. The limitations include the relatively-short duration of the blinded follow-up (28 weeks) and the underrepresentation of Black or African-American patients, who comprised only 9% of the study population.

Statement 9 reached consensus (97.3% of agreement in round 1 of voting), recommending that the choice among eculizumab, inebilizumab, ravulizumab, or satralizumab should consider factors such as drug availability, comorbidities or other autoimmune disorders, and patient preference, since efficacy and safety have not been compared in head-to-head studies. According to the GRADE framework, there is low-level evidence to support clinical decision-making, and previous publications⁵² were heterogeneous in their assessment of efficacy, using different endpoints and relapse definitions. Moreover, real-world evidence^{53,54} on the use of novel mAbs in individuals with NMOSD remains scarce, primarily due to significant disparities in therapeutic access across geographic regions and diverse populations. While these therapies are approved and available in many high-income countries, access is still limited or delayed in low- and middle-income countries, representing a major challenge and contributing to substantial global disparities in NMOSD care.¹⁷

From a clinical standpoint, mechanisms of action may assist in the selection of appropriate therapies in patients with comorbidities. Of note, eculizumab, ravulizumab, and inebilizumab have also been also approved for the treatment

of generalized myasthenia gravis (gMG), which can co-occur with NMOSD,⁵⁵ and, in such cases, they may offer dual therapeutic benefits.⁵⁶ Moreover, IL-6 receptor-blocking mAbs (that is, satralizumab and tocilizumab) may be advantageous in patients with NMOSD and coexisting autoimmune diseases—such as rheumatoid arthritis, giant cell arteritis, or systemic sclerosis—in which IL-6 signaling plays a pivotal role in pathogenesis.¹⁵

Safety profiles are a critical consideration. Complement inhibitors (eculizumab and ravulizumab) require meningococcal vaccination before initiation,¹⁴ whereas patients treated with inebilizumab or satralizumab require monitoring for infections and hematological abnormalities.^{12,15}

Dosing frequency and administration routes significantly impact adherence to treatment and quality of life. Ravulizumab is administered intravenously every 8 weeks, offering a significant advantage over eculizumab, which requires bi-weekly dosing.¹³ Inebilizumab, on the other hand, is administered intravenously every 6 months, a regimen that may be particularly appealing to patients residing in rural or remote areas, since it minimizes the need for frequent travel.¹² In contrast, satralizumab is administered monthly by SC injections, which enables self-injection at home and offers a significant advantage for patients with difficult venous access or those seeking to reduce hospital visits.⁵⁷ Finally, regarding adverse events, satralizumab treatment was generally well tolerated, with only a slightly higher risk of infections compared to placebo.⁵⁷ Thus, the panel agreed that therapeutic decisions in NMOSD should be individualized, taking into account not only efficacy and safety, but also comorbidities, drug accessibility, dosing convenience, and patient preference.

Statement 10 reached consensus (95.5% of agreement in round 1 of voting), not recommending the use of combined mAb therapy in patients diagnosed with NMOSD. Combining mAb therapy in NMOSD is not recommended due to insufficient evidence on its safety and efficacy. Most clinical trials were designed to evaluate monotherapy.¹⁶ At initial phases,⁵⁸ most studies kept pwNMOSD on a stable dose of previous IST, but not mAbs. In this regard, low-quality evidence describes⁵⁹ the use of IST associated with mAbs, even though combining these therapies could increase the risk of adverse effects such as infections. Future research may provide new insights into its potential benefits.

Statement 11 reached consensus (75% of agreement in round 2 of voting), recommending against the use of azathioprine, cyclophosphamide, oral steroids, methotrexate, or mycophenolate mofetil as first-line preventive monotherapy in adults with AQP4-IgG-seropositive NMOSD. The original version of this statement included adults and children, but it failed to reach consensus (63% of agreement) due to insufficient evidence. The final recommendation focused solely on adults, acknowledging the lack of pediatric data, which is addressed separately in statement 16.

Due to the lack of high-to-moderate-quality evidence regarding the efficacy and safety of these drugs, in contrast to the high-quality evidence for early use of mAbs such as rituximab, the use of immunosuppressants as monotherapy

as first-line treatment in adults with AQP4-IgG-seropositive NMOSD was not recommended. In addition, rituximab is associated with improvements in cost-efficacy⁶⁰ and quality of life.⁶¹ The key limitations of traditional immunosuppressants include: the lack of RCTs, with most studies being retrospective, with small sample sizes (especially in comparator studies), inconsistent treatment protocols, and a high risk of bias (example: concomitant corticosteroid use), and inferior efficacy compared to mAbs, as trials allowing background immunosuppressants (such as azathioprine, mycophenolate) showed worse outcomes than those using mAbs.³¹

The use of azathioprine is supported by seven observational studies ($n=641$) that showed reduced ARR,^{8,62–67} while confirming its inferiority to rituximab,⁶⁴ mycophenolate mofetil,⁶⁷ and cyclophosphamide.⁶⁶ Regarding disability, the EDSS scores improved by 0.98 points with rituximab versus 0.44 points with azathioprine ($p < 0.001$). A significantly higher proportion of patients achieved relapse-free status with rituximab (78.8%) compared to azathioprine (54.3%; $p=0.033$).⁶⁴ New relapses were observed in 33.5% of the patients, and the impact on disability was highly variable. A prospective cohort study compared azathioprine, mycophenolate mofetil, and lower doses of rituximab; no difference was found between azathioprine and mycophenolate mofetil, while low doses of rituximab were significantly more effective in reducing relapse rates, with higher relapse-free status and a better tolerability profile. The concomitant use of azathioprine and corticosteroids in some studies^{62,63,65,66} may have favored the results observed. There was a significant rate of adverse events (35%) and discontinuation (16%) while using azathioprine.⁸ The use of oral prednisolone is supported by two small Japanese retrospective observational studies^{68,69} with a small number of cases and classified as very low-quality evidence; both studies reported a significant reduction in ARR (0.49 versus 1.48 and 0.21 versus 0.98; $p < 0.01$) and stabilization of the EDSS scores (change: +0.02 versus +0.89; $p < 0.01$) if compared with no medication. Adverse events, ranging from mild to severe, were reported (online Supplementary Material). The use of methotrexate (after the previous use of other drugs) is supported by limited, very low-quality evidence from two retrospective observational studies^{70,71} showing a reduction in ARR (from 1.39 pretreatment to 0.18 posttreatment, $p < 0.005$; and from 0.97 to 0.00, $p < 0.05$), albeit relapses still occurred in 57% of the patients⁷⁰ (online Supplementary Material).

The evaluation on the use of mycophenolate mofetil is based on the results of 16 observational studies involving 786 pwNMOSD treated with it as monotherapy. Reduced relapse rates were observed in nine studies;^{72–80} however, the overall evidence remains insufficient to support a definitive treatment recommendation due to its predominantly low- to very low-quality (online Supplementary Material). In a cohort of 90 AQP4-IgG-positive patients receiving monotherapy, high and moderate doses of micophenolate mofetil led to greater reductions in AAR and EDSS scores.⁷³ In comparative analyses,^{67,73,77,81,82} the efficacy of micophenolate mofetil was lower than that of rituximab. Adding all patients from

the 16 studies, new relapses were reported in 23% of the total number of patients treated, and adverse effects, in 18%, ranging from mild to severe dose-dependent.⁷⁵

Preventive treatment for patients with AQP4-IgG-negative NMOSD

Statements 7 and 12 to 14 were developed to guide the preventive treatment in patients with AQP4-IgG-negative NMOSD, and they are shown in **Table 7**.

Statement 7 reached the consensus (77.3% of agreement in round 1 of voting) that rituximab treatment is recommended at the moment the diagnosis is established, after the first attack, and after relapse due to the failure of previous treatments, for the prevention of attacks and disability in adults and children diagnosed with NMOSD, regardless of the AQP4-IgG serostatus.

Evidence of preventive treatment in AQP4-IgG-negative NMOSD is scarce, according to the GRADE framework. Even so, there was no difference in relapse rates and disability scores between AQP4-IgG seropositive and seronegative NMOSD in a recent meta-analysis of patients treated with rituximab. Similar findings were reported in retrospective studies^{83,84} that included approximately 20% seronegative patients, which showed reductions in ARR and stabilization or improvement in disability scores. Observational evidence also suggests that rituximab has comparable efficacy in AQP4-IgG-negative patients⁸⁵ and in pediatric populations,⁸⁶ supporting its use across a broader spectrum of NMOSD phenotypes. Moreover, rituximab is associated with a better quality of life in pwnMOSD, and its early use is associated with a better cost-effectiveness profile in an economic model.^{60,61} While these findings are based on lower-certainty evidence, they reinforce rituximab's potential usefulness beyond AQP4-IgG-positive adult patients.

Key considerations for seronegative patients include the identification of clinical heterogeneity and the need for the exclusion of MOGAD and other disease mimics. Regarding comparative efficacy, rituximab remains the first-line therapy due to its consistent outcomes across NMOSD subtypes.⁸⁵

The panel recommended that, in AQP4-IgG-positive adults with NMOSD, azathioprine, mycophenolate, steroids, methotrexate, or cyclophosphamide should be avoided as monotherapy due to low-quality evidence regarding efficacy and safety. Rituximab was preferred in this population. In AQP4-IgG-negative NMOSD, rituximab was also recommended, despite the lower evidence quality, supported by real-world data.

Statement 12 reached consensus (84.1% of agreement in round 1 of voting), suggesting that azathioprine and mycophenolate mofetil should be used as preventive therapy in all AQP4-IgG-negative pwnMOSD (children and adults) only in the initial phase of the disease and when other drugs (rituximab and tocilizumab) are not possible. These IST should be prescribed with caution, given the absence of high-quality evidence.

Due to limited evidence and the lack of proven efficacy of mAbs in patients with AQP4-IgG-negative NMOSD, therapeutic options remain restricted. However, there is growing

support for the use of azathioprine and mycophenolate mofetil in seronegative patients. The panel's suggestion is supported by the significant reduction in relapse rates with azathioprine use, regardless of the AQP4-IgG serostatus.^{62,64} Moreover, four observational studies^{77-79,82} have reported comparable reductions in relapse rates with mycophenolate mofetil among seronegative individuals. Still, the evidence is limited by methodological constraints, and we advise cautious therapeutic decision-making.

Furthermore, three comparative observational studies support the superiority of rituximab^{64,82} and tocilizumab⁸⁷ over azathioprine. The evidence favoring rituximab is rated as moderate-quality by the GRADE, indicating a reasonable level of confidence. A multicenter, phase-2, open-label trial⁸⁷ with 59 patients demonstrated a significant reduction in the risk of relapse with the use of tocilizumab (HR=0.17; 95%CI=0.04-0.76), with 81% of the patients remaining relapse-free at 48 weeks, compared to 37% of those treated with azathioprine. This moderate certainty reflects a potential bias from the open-label design and small sample size, but the results suggest that tocilizumab is indeed a promising alternative for the treatment of high-risk NMOSD patients. Of note, adverse events were comparable between the groups, supporting an acceptable safety profile for tocilizumab.⁸⁷

Statement 13 reached consensus (70.5% of agreement in round 3 of voting), suggesting the use of tocilizumab as monotherapy—either at the moment the diagnosis is established, after the first attack, or following relapse due to the failure of previous treatments—for the prevention of attacks and disability in adults with AQP4-IgG-negative NMOSD. Four observational studies⁸⁸⁻⁹¹ of very low-quality evidence and one moderate-quality RCT⁸⁷ have indicated that tocilizumab (8 mg/kg IV Q4W) may reduce ARR in AQP4-IgG-positive and -negative NMOSD. The treatment was well tolerated, with a lower risk of adverse events (primarily infections and transaminase elevations) compared to azathioprine. The panel recommends diligent monitoring of the treatment (statement 3). Of note, an initial statement was designed, in which we suggested the administration of this treatment to adults (aged ≥ 18 years) with NMOSD, regardless of the AQP4 serostatus. However, due to low-quality evidence regarding the efficacy of tocilizumab when compared with evidence from mAbs evaluated in an RCT for AQP4-IgG-positive patients, this statement did not reach consensus after the first (agreement of 65.9%) or second (agreement of 59.1%) voting rounds. Therefore, this statement was revised and finally reached consensus (agreement of 70.5%) after the third voting round, with tocilizumab treatment being recommended only for AQP4-IgG-negative NMOSD patients.

Statement 14 reached consensus (97.7% of agreement in round 1 of voting), not recommending the use of other B-cell-depleting therapies (ocrelizumab, ofatumumab, or ublituximab) for patients diagnosed with NMOSD, regardless of the AQP4-IgG serostatus, given the current absence of supporting evidence. A study assessing ofatumumab in pwnMOSD was registered, but its completion date has

passed, and its status has not been updated in more than 2 years. To date, no studies have evaluated the efficacy of other B-cell-depleting therapies other than rituximab and inebilizumab in pwNMOSD.

Preventive treatment in patients with NMOSD: special clinical situations

Statement 15 was developed to guide the preventive management in patients with NMOSD who remain clinically stable and relapse-free over prolonged periods while under ongoing IST, regardless of the AQP4-IgG serostatus or age (–Table 8). Its significance arises from two considerations: first, stable and relapse-free patients are typically excluded from RCTs; and second, no biomarker is currently available to predict relapse in this population.

Statement 15 reached the consensus (90.1% of agreement in round 1 of voting) that preventive therapy with azathioprine, mycophenolate mofetil, or rituximab can be maintained in all patients (adults or children) already undergoing treatment and who are both relapse-free and present no adverse events, regardless of the AQP4-IgG serostatus. This statement is based on pivotal RCTs assessing mAbs in pwNMOSD, all of which required the inclusion of patients with active disease despite previous treatments. The RIN-1 trial⁴⁶ enrolled patients with documented relapses within the preceding 2 years. The PREVENT trial¹⁴ included patients with ≥ 2 relapses in the previous 12 months or ≥ 3 in the previous 24 months, including at least 1 in the last year. The open-label CHAMPION-NMOSD extension¹³ enrolled patients who presented at least 1 relapse in the preceding 12 months. In the SakuraSky study,⁵⁷ patients with ≥ 2 relapses within 24 months (including ≥ 1 in the last year) were enrolled while maintaining their previous immunosuppressants, whereas in the study called “A Multicenter, Randomized, Addition to Baseline Treatment, Double-Blind, Placebo-Controlled, Phase 3 Study to Evaluate the Efficacy and Safety of Satralizumab (SA237) in Patients With Neuro-myelitis Optica (NMO) and NMO Spectrum Disorder (NMOSD)” (SakuraStar),⁵⁷ the patients had experienced ≥ 1 relapse in the previous 12 months, discontinuing any previous immunosuppressants.

Given this context, there is no high-level-evidence, according to the GRADE framework, to guide switching from conventional ISTs to mAbs in patients with stable NMOSD. The panel members agreed that patients who are relapse-free and without adverse events on their current therapy do not need to be switched to a new one. While waiting for further real-world data and the identification of new biomarkers able to predict treatment failure, the panel acknowledged that therapeutic changes in this subgroup may be considered on a case-by-case basis.

Preventive treatment in children with NMOSD

Statements 7, 16, 17, and 18 were developed to the guide preventive treatment in children with NMOSD (–Table 9).

Statement 7 reached consensus (77.3% of agreement in round 1 of voting), recommending rituximab as monotherapy at the moment the diagnosis is established, after the first

attack, and after relapse due to the failure of previous treatments, for the prevention of attacks and disability in adults or children diagnosed with NMOSD, regardless of the AQP4-IgG serostatus. According to the GRADE framework, the use of rituximab has a high level of evidence for AQP4-IgG-positive NMOSD patients, and a low to moderate evidence for patients with DS-NMOSD and children. This statement reiterates the recommendation of rituximab for the prevention of relapses and disability in children with NMOSD, regardless of the AQP4 serostatus. This is supported by four retrospective observational studies involving 91 pediatric patients.^{92–95} Three studies^{92–94} analyzed the ARR, 2 of which observed a decrease in ARR either from 2.50 to 0.14 ($p < 0.001$),⁹² or from 1 to 0,⁹⁴ while a third noted a stable ARR of 0.25.⁹³ Two studies^{93,95} evaluated patients who remained relapse-free during treatment, with 87.5%⁹⁵ and 31.5%⁹³ of patients free from relapses over periods ranging from 6 to 150 months. No data were available on the time until the first relapse. Adverse events were only assessed in one study,⁹² which reported infusion reactions in three patients and one case of neutropenia. Given the reliance on observational data, caution is advised when using rituximab in pediatric patients, emphasizing the need for monitoring for infusion-related reactions and neutropenia, as per statement 3.

Statement 16 reached consensus (93.2% of agreement in round 1 of voting), recommending satralizumab as monotherapy or combined with oral immunosuppressants as treatment at the moment the diagnosis is established, after the first attack, and after relapse due to the failure of previous treatments, for the prevention of attacks and disability in adolescents (aged ≥ 12 years) with AQP4-IgG-positive NMOSD. This statement is based on a high-quality study¹⁵ that analyzed the efficacy and adverse effects of satralizumab treatment in pwNMOSD. Considering the rarity of NMOSD in children, conducting RCTs with sufficient statistical power exclusively for pediatric patients remains extremely challenging.

Statement 17 reached consensus (86.4% of agreement in round 1 of voting), suggesting that azathioprine and mycophenolate mofetil may be used as preventive therapies in children with NMOSD, regardless of the AQP4-IgG serostatus, only with caution and when rituximab and satralizumab are contraindicated. This recommendation is supported by five retrospective observational studies^{92–96} that evaluated the use of azathioprine in pediatric patients, all of which were of low-quality evidence regarding efficacy, according to the GRADE framework. Three studies reported a reduction in the ARR before and after azathioprine treatment, from 0.89 to 0.65 in 11 AQP4-IgG-positive patients,⁹⁴ from 1.69 to 0.59 in 39 AQP4-IgG-positive patients,⁹² and from 1.26 to 0.40 in 10 patients irrespective of the AQP4-IgG serostatus⁹⁶. Only two studies^{93,95} evaluated the proportion of relapse-free patients using azathioprine as monotherapy or in combination with oral corticosteroids: Umeton et al.⁹³ reported that 33% of the patients were relapse-free, while McKeon et al.⁹⁵ reported a rate of 50% of relapse-free patients. There is no available data regarding the time until the first relapse, relapse prevention, or changes in disability accumulation.

The use of mycophenolate is supported by four observational studies,^{92–95} all of very low-quality evidence according to the GRADE framework. Three studies^{92–94} evaluated the ARR regardless of the AQP4-IgG serostatus, and the mean ARR across the studies was of 0.44. In two studies this decreased after treatment, from 0.98 to 0.28,⁹⁴ and from 1.04 to 0.72.⁹² Another study⁹⁵ reported that 66% of the patients remained relapse-free over a 12- to 24-month period. In none of these studies disability accumulation or its reduction were reported. Although observational studies have shown reductions in the ARR and a low frequency of side effects, high- or moderate-quality evidence is currently lacking to confirm the efficacy of mycophenolate mofetil for relapse reduction, time until relapse, disability accumulation or reduction, or its safety in children and adolescents. Only low-quality studies have evaluated the safety of azathioprine and mycophenolate mofetil in pediatric patients with NMOSD. Among those treated with azathioprine, the reported adverse events include lymphopenia, infections, gastrointestinal disturbances, and elevated liver enzymes. Therefore, caution is recommended if azathioprine or mycophenolate are required to be used in pediatric populations, with particular attention to the increased risk of infections and gastrointestinal complications. Active monitoring for infections, vaccination status, and laboratory monitoring of liver enzymes and cytopenias are strongly advised during treatment with either medication.⁹⁷

Statement 18 reached consensus (81.8% of agreement in round 1 of voting), not recommending treatment with cyclophosphamide, oral steroids, or methotrexate as preventive monotherapy for children with NMOSD, regardless of the AQP4-IgG serostatus. The use of methotrexate was reported⁹⁵ in 1 patient who had no relapses over 36 months; in the same study, on the other hand, two children relapsed during treatment with cyclophosphamide.

Preventive treatment in NMOSD during pregnancy and breastfeeding (► Table 10)

Statement 19 reached consensus (84.1% of agreement in round 1 of voting), suggesting that rituximab may be considered during pregnancy and breastfeeding in women with NMOSD, regardless of the AQP4-IgG serostatus, as long as the patient consents and treatment is closely monitored in collaboration with the obstetrics team. In eight low-quality studies,^{98–105} the use of azathioprine, rituximab, and prednisolone during pregnancy and the postpartum period was described. These studies concluded that maintaining the IST throughout pregnancy and postpartum reduces the ARR and prevents disability accumulation. Data on relapse rates, disability progression, or adverse effects on the fetus or pregnant patients were not reported. These treatments should only be administered with the patient's full informed consent and under close monitoring by the neurology and obstetrics teams. Despite the lack of high- or moderate-evidence efficacy studies on the use of azathioprine in patients with NMOSD, its use is supported in statements 12 and 13 in cases in which rituximab is unavailable or in patients already receiving azathioprine with a favorable therapeutic response.

Teratogenicity has not been identified in humans and, based on safety data from pregnant patients with inflammatory bowel disease and organ transplant recipients, azathioprine has recently been recognized¹⁰⁶ as a safe immunosuppressant during pregnancy. Breastfeeding is compatible with azathioprine use, as drug transfer into breast milk is minimal.¹⁰⁷ Nevertheless, treatment should be accompanied by informed patient consent and close monitoring throughout the perinatal period.

Statement 20 reached consensus (100% of agreement in round 1 of voting), contraindicating treatment with cyclophosphamide, methotrexate, or mycophenolate mofetil during pregnancy and breastfeeding in women with NMOSD, regardless of the AQP4-IgG serostatus, due to teratogenic and toxicity risks. In safety studies in women of childbearing age and pregnant women with NMOSD, methotrexate was contraindicated due to teratogenic and abortifacient effects,¹⁰⁸ cyclophosphamide, due to known embryofetal toxicity,¹⁰⁹ and mycophenolate mofetil, due to the risk of spontaneous abortion and congenital anomalies.¹¹⁰ Cyclophosphamide was also contraindicated during breastfeeding because it is excreted in breast milk at toxic levels.¹¹¹

Statement 21 reached consensus (86.3% of agreement in round 1 of voting), not recommending treatment with eculizumab, inebilizumab, ravulizumab, or satralizumab during pregnancy and breastfeeding in women with NMOSD, given the absence of evidence to support such an indication. There is no safety data on the use of these medications in pregnant or lactating women. Agencies such as the Brazilian Health Regulatory Agency (Agência Nacional de Vigilância Sanitária, ANVISA, in Portuguese) or the United States Food and Drug Administration (FDA) have not yet classified the risks of these drugs during pregnancy and breastfeeding.

DISCUSSION

The current consensus represents the first initiative to capture expert perspectives on the management of NMOSD in a continental developing country. Developed using the Delphi methodology, a validated approach for scientific discussions among expert panels, the present guideline aims to generate knowledge on topics with limited scientific evidence, such as NMOSD. A selected group of neuroimmunology experts contributed to the development and validation of the consensus statements on the management of NMOSD, based on the quality of evidence to support therapeutic decision-making directed at people living with this disease.

Additionally, social determinants of health are known to impact outcomes in pwNMOSD. This may be worse in scenarios in which disparities between public and private healthcare systems affect access to timely diagnosis and treatment, such as the access to AQP4-IgG testing, which, in Brazil, is available in the private sector, but not within the public system routinely. In Brazil, the incorporation of new diagnostic tools and treatments is evaluated by the National Commission for the Incorporation of Technologies into the Unified Health System (Comissão Nacional de Incorporação

de Tecnologias no Sistema Único de Saúde, CONITEC, in Portuguese), which advises the Ministry of Health on the technologies and clinical guidelines that should be included into the Brazilian Unified Health System (Sistema Único de Saúde, SUS, in Portuguese). Currently, there are no therapeutic options for NMOSD available through either the SUS or private health insurance. This limitation derives from the absence of a defined protocol within SUS and the lack of regulatory definitions from the Brazilian National Health Agency (Agência Nacional de Saúde, ANS, in Portuguese), which prevents private-sector physicians from prescribing appropriate medications for NMOSD. The panel concurs that the development of an official national guideline for NMOSD management should be prioritized to ensure tailored care for patients in a country with a high prevalence of the disease.¹¹²

Given the complexity and distinct pathophysiology of immune-mediated neurological diseases, the present study specifically focused on NMOSD seropositive for AQP4-IgG and NMOSD seronegative for both AQP4-IgG and MOG-IgG, in agreement with the 2015 IPND diagnostic criteria.¹ It is noteworthy to emphasize that the current consensus does not address, and its recommendations are not intended for, other immune-mediated conditions such as MOGAD, once this is now recognized as distinct entity. Neither do our recommendations encompass high-risk syndromes suggestive of NMOSD, not fulfilling 2015 IPND diagnostic criteria,¹ which are seronegative for both AQP4-IgG and MOG-IgG, including isolated or recurrent optic neuritis, chronic relapsing idiopathic optic neuritis, isolated or recurrent transverse myelitis, and brainstem syndromes. Brazilian guidelines underscore the urgent need for early and widespread access to AQP4-IgG testing. In early 2025, CONITEC recommended its inclusion in the SUS for patients with clinical and radiological features of NMOSD. Despite regulatory approval, the test remains largely inaccessible in most public healthcare settings. This is expected to improve early diagnosis and pave the way for the formulation of treatment planning strategies in this country with continental dimensions.

The primary objective in the management of patients diagnosed with NMOSD is to mitigate the severity of the attacks and prevent relapses, thereby minimizing the risk of irreversible neurological deficits from recurrent episodes. Regarding acute-stage management, it was reemphasized that exacerbations in pNMOSD result in severe residual disability. Therefore, the treatment of relapses must begin promptly and aggressively, which is crucial to prevent severe disability. The goals of the acute-phase therapy include suppressing relapses, minimizing CNS damage, and promoting optimal functional recovery. The evidence supports the principle of 'the sooner, the better' for both high-dose methylprednisolone and apheresis. From a pragmatic perspective, in certain regions of Brazil, significant delays in accessing tertiary care centers where plasmapheresis may be available remain a challenge. Nonetheless, such delays should not preclude the administration of optimal treatment strategies during the acute phase. This emphasizes the critical importance of ensuring that these patients are adequately monitored and managed in specialized centers.

The accessibility and affordability of recently-approved treatments for NMOSD are likely to vary across countries and regions, significantly influencing clinical decisions regarding the timing of treatment initiation and the choice of the therapeutic agent.^{16,17,58} Recently, several prospective RCTs have led to the FDA approval of the first three immunotherapies for patients with AQP4-IgG-positive NMOSD: eculizumab in June 2019, inebilizumab in June 2020, and satralizumab in August 2020. In addition, rituximab was approved for NMOSD in Japan in June 2022 based on the results of an investigator-initiated phase-II/III clinical study; and in May 2023, the European Medicines Agency (EMA) approved ravulizumab for the treatment of AQP4-IgG-positive NMOSD. In Brazil, ravulizumab, satralizumab, and inebilizumab were respectively approved by ANVISA in 2022, 2020, 2023 respectively.²⁴ To date, there are no head-to-head trials or direct clinical evidence to guide the best treatment strategy for NMOSD, and the current recommendations are based on expert opinions and consensus.^{16,17,58,113,114} In upper-middle-income countries such as Brazil, the heterogeneous healthcare landscape—including limited access to serum diagnostics, acute-phase interventions like therapeutic apheresis, and high-efficacy monoclonal antibody therapies—compounds the challenges of clinical management and underscores the urgent unmet needs that this consensus aims to address.

In conclusion, these guidelines, developed through an evidence-based modified Delphi process, constitute a significant step towards the standardization of care. The statements herein proposed aim to address a critical gap by providing evidence-based recommendations for the management of patients with NMOSD within the Brazilian context.

Authors' Contributions

Conceptualization: SLAP, AD, ACP, MFM, MLB, RA, JB, CCFV; Data curation: SLAP, AD, ACP, MFM, BKC, DKS, DGB, EM, FvG, GP, GD, HSN, LC, MF, MSP, RP, RV, RBP, TA, TF, VD, JB, CCFV; Formal analysis: SLAP, AD, ACP, MFM, BKC, DKS, DGB, EM, FvG, GP, GD, HSN, LC, MF, MSP, RP, RV, RBP, TA, TF, VD, JB, CCFV; Writing - original draft: SLAP, AD, ACP, MFM, MLB, RA, JB, CCFV.

Data Availability Statement

The datasets generated and/or analyzed during the current study are available from the corresponding author upon request.

Funding

The authors declare that the present study was funded by the Brazilian Academy of Neurology (Academia Brasileira de Neurologia, ABN, in Portuguese).

Conflict of Interest

SLAP: received honoraria for lectures and for participating in advisory boards for Roche, AstraZeneca, Amgen, Merck, and Novartis. AD: received honoraria for being a member of advisory boards for Alexion, Amgen, AstraZeneca,

Horizon, Merck Serono, Novartis, and Roche; and for acting as an investigator for Roche. ACP: received honoraria and congress travel support for lectures and for being a member of advisory boards for Amgen, Roche, and AstraZeneca. MFM: received honoraria and congress travel support for lectures from Biogen, EMS, AstraZeneca, and Merck; and for acting as an investigator for Merck. MLB: received honoraria for lectures and for participating in advisory boards for EMS, Roche, Merck, and Biogen. RA: received honoraria from Horizon for being a member of an advisory board and for coordinating a round table during a symposium. BKC: received honoraria for lectures and for acting as an investigator from Roche, Amgen, and Novartis; BKC also received grants and fellowships from the Multiple Sclerosis International Federation (MSIF) and from ARSEP. DKS: received honoraria for lectures and for participating in advisory boards from Roche, AstraZeneca, Amgen, Merck, and Novartis. EM: received honoraria and congress travel support for lectures from Amgen, Merck, Roche, Novartis, and Biogen. FVG: received honoraria for lectures from Biogen, AstraZeneca, Amgen, Novartis, and Roche. GP: received honoraria and/or travel support for lectures and/or for participating in advisory boards from Alexion/AstraZeneca, Alnylam, Biogen, EMS, Horizon/Amgen, Merck, Novartis, Roche, and Sanofi Genzyme; GP also acted as an investigator for Alexion/AstraZeneca and Merck. GD: received honoraria for lectures from Novartis and for acting as a consultant for Amgen, and travel support from EMS (for winning the "Neuro-impacto" competition). HSN: received honoraria for lectures and/or for participating in advisory boards for AstraZeneca, Amgen, Roche, Merck, Sanofi, and Novartis. LC: received honoraria for lectures from AstraZeneca, Biogen, Novartis, and Roche; and received travel support from Merck and Amgen. MF: received honoraria for acting as a researcher for Roche. MSP: received honoraria for lectures from Amgen, AstraZeneca, Biogen, Merck, Novartis, and Roche; MSP also received honoraria for acting as an investigator for Roche. RP: received honoraria and travel support for lectures and/or for acting as an investigator for Alexion/AstraZeneca, Amgen, and Roche. RV: received honoraria and/or company shares as compensation for lectures, for acting as a consultant, and for event sponsorships from Horizon/Amgen, Alexion/AstraZeneca, and Roche. RBP: received honoraria and congress support for lectures and/or for acting as an investigator for Amgen, Novartis, Roche, and AstraZeneca. TA: received honoraria for lectures and for participating in advisory boards from AstraZeneca, Biogen, Merck, Novartis, and Roche. TF: received honoraria for lectures and/or for acting as an investigator for AstraZeneca, Roche, and Amgen. VD: received honoraria for lectures from AstraZeneca and Alexion, and for participating in advisory boards from AstraZeneca, Alexion, Amgen, and Roche. JB: received honoraria for lectures and/or for participating in advisory boards and/or for acting as an investigator for Roche, Amgen, and AstraZeneca. CCFV: received honoraria for lectures from Biogen; and for participating in advisory

boards from Roche, Novartis, Merck, Sanofi, Amgen, and AstraZeneca. DGB has no conflict of interest to declare. DGB has no conflict of interest to declare.

Acknowledgments

The authors would like to thank the Brazilian Academy of Neurology (Academia Brasileira de Neurologia, ABN, in Portuguese) for all the support that made this work possible, the NMOSD-Brazilian Study Group and Cristina Stephan, M.D., M.Sc., of StepMedical Affairs Consulting, for her support in the final review of the present paper.

References

- Wingerchuk DM, Banwell B, Bennett JL, Cabre P, Carroll W, Chitnis T, et al; International Panel for NMO Diagnosis. International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology* 2015;85(02):177–189. Doi: 10.1212/WNL.0000000000001729
- Hor JY, Asgari N, Nakashima I, Broadley SA, Leite MI, Kissani N, et al. Epidemiology of Neuromyelitis Optica Spectrum Disorder and Its Prevalence and Incidence Worldwide. *Front Neurol* 2020; 11:501. Doi: 10.3389/fneur.2020.00501
- Papp V, Magyari M, Aktas O, Berger T, Broadley SA, Cabre P, et al. Worldwide Incidence and Prevalence of Neuromyelitis Optica: A Systematic Review. *Neurology* 2021;96(02):59–77. Doi: 10.1212/WNL.00000000000011153
- Lennon VA, Wingerchuk DM, Kryzer TJ, Pittock SJ, Lucchinetti CF, Fujihara K, et al. A serum autoantibody marker of neuromyelitis optica: distinction from multiple sclerosis. *Lancet* 2004;364 (9451):2106–2112. Doi: 10.1016/S0140-6736(04)17551-X
- Waters PJ, McKeon A, Leite MI, Rajasekharan S, Lennon VA, Villalobos A, et al. Serologic diagnosis of NMO: a multicenter comparison of aquaporin-4-IgG assays. *Neurology* 2012;78(09): 665–671, discussion 669. Doi: 10.1212/WNL.0b013e318248dec1
- Sato DK, Callegaro D, Lana-Peixoto MA, Waters PJ, Jorge FMdH, Takahashi T, et al. Distinction between MOG antibody-positive and AQP4 antibody-positive NMO spectrum disorders. *Neurology* 2014;82(06):474–481. Doi: 10.1212/WNL.0000000000000101
- Wu Y, Gerales R, Juryńczyk M, Palace J. Double-negative neuromyelitis optica spectrum disorder. *Mult Scler* 2023;29(11-12):1353–1362. Doi: 10.1177/13524585231199819
- Bichuetti DB, Perin MMM, Souza NA, Oliveira EML. Treating neuromyelitis optica with azathioprine: 20-year clinical practice. *Mult Scler* 2019;25(08):1150–1161. Doi: 10.1177/1352458518776584
- Wingerchuk DM, Hogancamp WF, O'Brien PC, Weinshenker BG. The clinical course of neuromyelitis optica (Devic's syndrome). *Neurology* 1999;53(05):1107–1114. Doi: 10.1212/wnl.53.5.1107
- Francis A, Gibbons E, Yu J, Johnston K, Rochon H, Powell L, et al. Characterizing mortality in patients with AQP4-Ab+ neuromyelitis optica spectrum disorder. *Ann Clin Transl Neurol* 2024;11 (07):1942–1947. Doi: 10.1002/acn3.52092
- Royston M, Kielhorn A, Weycker D, Shaff M, Houde L, Tanvir I, et al. Neuromyelitis Optica Spectrum Disorder: Clinical Burden and Cost of Relapses and Disease-Related Care in US Clinical Practice. *Neurol Ther* 2021;10(02):767–783. Doi: 10.1007/s40120-021-00253-4
- Cree BAC, Bennett JL, Kim HJ, Weinshenker BG, Pittock SJ, Wingerchuk DM, et al; N-MOMentum study investigators. Inebilizumab for the treatment of neuromyelitis optica spectrum disorder (N-MOMentum): a double-blind, randomised placebo-controlled phase 2/3 trial. *Lancet* 2019;394(10206):1352–1363. Doi: 10.1016/S0140-6736(19)31817-3
- Pittock SJ, Barnett M, Bennett JL, Berthele A, Sèze Jd, Levy M, et al. Ravulizumab in Aquaporin-4-Positive Neuromyelitis Optica

- Spectrum Disorder. *Ann Neurol* 2023;93(06):1053–1068. Doi: 10.1002/ana.26626
- 14 Pittock SJ, Berthele A, Fujihara K, Kim HJ, Levy M, Palace J, et al. Eculizumab in Aquaporin-4-Positive Neuromyelitis Optica Spectrum Disorder. *N Engl J Med* 2019;381(07):614–625. Doi: 10.1056/NEJMoa1900866
 - 15 Yamamura T, Kleiter I, Fujihara K, Palace J, Greenberg B, Zakrzewska-Pniewska B, et al. Trial of Satralizumab in Neuromyelitis Optica Spectrum Disorder. *N Engl J Med* 2019;381(22):2114–2124. Doi: 10.1056/NEJMoa1901747
 - 16 Paul F, Marignier R, Palace J, Arrambide G, Asgari N, Bennett JL, et al. International Delphi Consensus on the Management of AQP4-IgG+ NMOSD: Recommendations for Eculizumab, Inebilizumab, and Satralizumab. *Neurol Neuroimmunol Neuroinflamm* 2023;10(04):e200124. Doi: 10.1212/NXI.0000000000200124
 - 17 Alonso R, Rivera V, Contentti EGC, Bedoya VFHDD, Ciampi E, Silva B, et al. Recommendations on NMOSD treatment in Latin America: Utilizing consensus-based RAND/UCLA methodology. *Mult Scler Relat Disord* 2025;94:106244. Doi: 10.1016/j.msard.2024.106244
 - 18 Balslem H, Helfand M, Schünemann HJ, Oxman AD, Kunz R, Brozek J, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol* 2011;64(04):401–406. Doi: 10.1016/j.jclinepi.2010.07.015
 - 19 Niederberger M, Spranger J. Delphi Technique in Health Sciences: A Map. *Front Public Health* 2020;8(457):457. Doi: 10.3389/fpubh.2020.00457
 - 20 Smela B, Toumi M, Świerk K, Francois C, Biernikiewicz M, Clay E, Boyer L. Rapid literature review: definition and methodology. *J Mark Access Health Policy* 2023;11(01):2241234. Doi: 10.1080/20016689.2023.2241234
 - 21 Richardson WS, Wilson MC, Nishikawa J, Hayward RS. The well-built clinical question: a key to evidence-based decisions. *ACPJ Club* 1995; 123(03):A12–A13. Doi: 10.7326/ACPJC-1995-123-3-A12
 - 22 Weinshenker BG, Wingerchuk DM, Vukusic S, Linbo L, Pittock SJ, Lucchinetti CF, Lennon VA. Neuromyelitis optica IgG predicts relapse after longitudinally extensive transverse myelitis. *Ann Neurol* 2006;59(03):566–569. Doi: 10.1002/ana.20770
 - 23 Contentti EC, Castillo ISd, Marques VD, López PA, Barreira AA, Armas E, et al. Application of the 2015 diagnostic criteria for neuromyelitis optica spectrum disorders in a cohort of Latin American patients. *Mult Scler Relat Disord* 2018;20:109–114. Doi: 10.1016/j.msard.2018.01.001
 - 24 Vassão-Araujo R, Apóstolos S, Jansen AM, Lana-Peixoto MA, Gomes Neto AP, Rico-Restrepo M, et al. A roadmap to increasing access to AQP4-Ig testing for NMOSD: expert recommendations. *Arq Neuro-Psiquiatr* 2025;83(02):1–11. Doi: 10.1055/s-0045-1801845
 - 25 Redenbaugh V, Montalvo M, Sechi E, Buciu M, Fryer JP, McKeon A, et al. Diagnostic value of aquaporin-4-IgG live cell based assay in neuromyelitis optica spectrum disorders. *Mult Scler J Exp Transl Clin* 2021;7(04):20552173211052656. Doi: 10.1177/2052173211052656
 - 26 Banwell B, Bennett JL, Marignier R, Kim HJ, Brilot F, Flanagan EP, et al. Diagnosis of myelin oligodendrocyte glycoprotein antibody-associated disease: International MOGAD Panel proposed criteria. *Lancet Neurol* 2023;22(03):268–282. Doi: 10.1016/S1474-4422(22)00431-8
 - 27 Delgado-García G, Lapidus S, Talero R, Levy M. The patient journey with NMOSD: From initial diagnosis to chronic condition. *Front Neurol* 2022;13:966428. Doi: 10.3389/fneur.2022.966428
 - 28 Gomes ABAGR, Feo LB, Silva GD, Disserol CCD, Paolillo RB, Lara AN, et al. Reducing infection risk in multiple sclerosis and neuromyelitis optica spectrum disorders: a Brazilian reference center's approach. *Arq Neuro-Psiquiatr* 2022;80(10):1057–1066. Doi: 10.1055/s-0042-1754348
 - 29 Mealy MA, Cook LJ, Pache F, Velez DL, Borisow N, Becker Det al. Vaccines and the association with relapses in patients with neuromyelitis optica spectrum disorder. *Mult Scler Relat Disord* 2018;23:78–82. Doi: 10.1016/j.msard.2018.05.003
 - 30 Ellrichmann G, Bolz J, Peschke M, Duscha A, Hellwig K, Lee D-H, et al. Peripheral CD19⁺ B-cell counts and infusion intervals as a surrogate for long-term B-cell depleting therapy in multiple sclerosis and neuromyelitis optica/neuromyelitis optica spectrum disorders. *J Neurol* 2019;266(01):57–67. Doi: 10.1007/s00415-018-9092-4
 - 31 Kleiter I, Traboulsee A, Palace J, Yamamura T, Fujihara K, Saiz A, et al. Long-term Efficacy of Satralizumab in AQP4-IgG-Seropositive Neuromyelitis Optica Spectrum Disorder From SAKuraSky and SAKuraStar. *Neurol Neuroimmunol Neuroinflamm* 2022;10(01):e200071. Doi: 10.1212/NXI.0000000000200071
 - 32 Kim S, Seok HY. Evaluation of the safety profile of COVID-19 vaccines in patients with MS, NMOSD, and MOGAD. *Neurol Sci* 2023;44(06):1841–1848. Doi: 10.1007/s10072-023-06676-1
 - 33 Cai H, Zhou R, Jiang F, Zeng Q, Yang H. Vaccination in neuromyelitis optica spectrum disorders: Friend or enemy? *Mult Scler Relat Disord* 2022;58:103394. Doi: 10.1016/j.msard.2021.103394
 - 34 Abboud H, Petrak A, Mealy M, Sasidharan S, Siddique L, Levy M. Treatment of acute relapses in neuromyelitis optica: Steroids alone versus steroids plus plasma exchange. *Mult Scler* 2016;22(02):185–192. Doi: 10.1177/1352458515581438
 - 35 Akaishi T, Takeshita T, Himori N, Takahashi T, Misu T, Ogawa R, et al. Rapid Administration of High-Dose Intravenous Methylprednisolone Improves Visual Outcomes After Optic Neuritis in Patients With AQP4-IgG-Positive NMOSD. *Front Neurol* 2020; 11:932. Doi: 10.3389/fneur.2020.00932
 - 36 Fu J, Wang Y, Li H, Zhou H, Song H, Sun M, et al. Efficacy of Plasma Exchange Treatment for Demyelinating Optic Neuritis Associated with Various Serum Antibodies: A Prospective Cohort Study. *Neurol Ther* 2022;11(02):797–813. Doi: 10.1007/s40120-022-00344-w
 - 37 Kumawat BL, Choudhary R, Sharma CM, Jain D, Hiremath A. Plasma Exchange as a First Line Therapy in Acute Attacks of Neuromyelitis Optica Spectrum Disorders. *Ann Indian Acad Neurol* 2019;22(04):389–394. Doi: 10.4103/aian.AIAN_365_19
 - 38 Bonnan M, Valentino R, Debeugny S, Merle H, Fergé J-L, Mehdaoui H, Cabre P. Short delay to initiate plasma exchange is the strongest predictor of outcome in severe attacks of NMO spectrum disorders. *J Neurol Neurosurg Psychiatry* 2018;89(04):346–351. Doi: 10.1136/jnnp-2017-316286
 - 39 Almeida GMRd, Araujo RSd, Castrillo BB, Silva GD, Fortini I, Gonçalves MRR, et al. Therapeutic plasma exchange for neuromyelitis optica attacks: Evidence and challenges from a real-world cohort from Brazil. *J Neuroimmunol* 2024;388:578295. Doi: 10.1016/j.jneuroim.2024.578295
 - 40 Kleiter I, Gahlen A, Borisow N, Fischer K, Wernecke K-D, Hellwig K, et al; NEMOS (Neuromyelitis Optica Study Group) Apheresis therapies for NMOSD attacks: A retrospective study of 207 therapeutic interventions. *Neurol Neuroimmunol Neuroinflamm* 2018;5(06):e504. Doi: 10.1212/NXI.0000000000000504
 - 41 Kleiter I, Gahlen A, Borisow N, Fischer K, Wernecke K-D, Wegner B, et al; Neuromyelitis Optica Study Group. Neuromyelitis optica: Evaluation of 871 attacks and 1,153 treatment courses. *Ann Neurol* 2016;79(02):206–216. Doi: 10.1002/ana.24554
 - 42 Qin C, Tao R, Zhang S-Q, Chen B, Chen M, Yu H-H, et al. Predictive Factors of Resistance to High-Dose Steroids Therapy in Acute Attacks of Neuromyelitis Optica Spectrum Disorder. *Front Neurol* 2020;11:585471. Doi: 10.3389/fneur.2020.585471
 - 43 Queiroz ALG, Soares Neto HR, Kobayashi TT, Silva SMCA. Plasma exchange in inflammatory demyelinating disorders of the central nervous system: reasonable use in the clinical practice. *Arq Neuro-Psiquiatr* 2023;81(03):296–307. Doi: 10.1055/s-0042-1758447
 - 44 Li X, Tian DC, Fan M, Xiu Y, Wang X, Li T, et al. Intravenous immunoglobulin for acute attacks in neuromyelitis optica

- spectrum disorders (NMOSD). *Mult Scler Relat Disord* 2020; 44:102325. Doi: 10.1016/j.msard.2020.102325
- 45 Siwach G, Hans R, Takkar A, Ahuja CK, Lamba DS, Lal V, Sharma RRL. Comparison of efficacy of plasma exchange versus intravenous immunoglobulin as an add-on therapy in acute attacks of neuromyelitis optica spectrum disorder. *J Clin Apher* 2024;39(03):e22129. Doi: 10.1002/jca.22129
 - 46 Tahara M, Oeda T, Okada K, Kiriya T, Ochi K, Maruyama H, et al. Safety and efficacy of rituximab in neuromyelitis optica spectrum disorders (RIN-1 study): a multicentre, randomised, double-blind, placebo-controlled trial. *Lancet Neurol* 2020;19(04):298–306. Doi: 10.1016/S1474-4422(20)30066-1
 - 47 Tahara M, Oeda T, Okada K, Ochi K, Maruyama H, Fukaura H, et al. Compassionate open-label use of rituximab following a randomised clinical trial against neuromyelitis optica (RIN-2 study): B cell monitoring-based administration. *Mult Scler Relat Disord* 2022;60:103730. Doi: 10.1016/j.msard.2022.103730
 - 48 Wang H, Zhou J, Li Y, Wei L, Xu X, Zhang J, et al. Adverse events of rituximab in neuromyelitis optica spectrum disorder: a systematic review and meta-analysis. *Ther Adv Neurol Disord* 2021; 14:17562864211056710. Doi: 10.1177/17562864211056710
 - 49 Pittock SJ, Fujihara K, Palace J, Berthele A, Kim HJ, Oreja-Guevara C, et al; PREVENT Study Group. Eculizumab monotherapy for NMOSD: Data from PREVENT and its open-label extension. *Mult Scler* 2022;28(03):480–486. Doi: 10.1177/13524585211038291
 - 50 Traboulsee A, Greenberg BM, Bennett JL, Szczechowski L, Fox E, Shkrobot S, et al. Safety and efficacy of satralizumab monotherapy in neuromyelitis optica spectrum disorder: a randomised, double-blind, multicentre, placebo-controlled phase 3 trial. *Lancet Neurol* 2020;19(05):402–412. Doi: 10.1016/S1474-4422(20)30078-8
 - 51 Cree BAC, Kim HJ, Weinshenker BG, Pittock SJ, Wingerchuk DM, Fujihara K, et al; N-MOMentum study investigators. Safety and efficacy of inebilizumab for the treatment of neuromyelitis optica spectrum disorder: end-of-study results from the open-label period of the N-MOMentum trial. *Lancet Neurol* 2024;23(06):588–602. Doi: 10.1016/S1474-4422(24)00077-2
 - 52 Rocha RB, Bomtempo FF, Tadinac ACB, Allioni GA, Silva GD, Telles JPM. Heterogenous relapse and efficacy endpoint definitions for neuromyelitis optica spectrum disorder studies: A systematic review. *Mult Scler Relat Disord* 2024;91:105868. Doi: 10.1016/j.msard.2024.105868
 - 53 Poisson KE, Nguyen L, Horn PS, Beck AF, Wu H, Wesselkamper K, et al. Racial, Ethnic, and Socioeconomic Disparities in Pediatric Aquaporin-4-Positive Neuromyelitis Optica Spectrum Disorder. *Pediatr Neurol* 2024;161:194–200. Doi: 10.1016/j.pediatrneurol.2024.09.028
 - 54 Tkachuk V, Aguirre MEB, Alonso R, Barboza A, Liwacki SDV, Mainella C, et al. Barriers to access and unmet needs to neuromyelitis optica spectrum disorders care in an Argentinean cohort. *Mult Scler Relat Disord* 2023;70:104485. Doi: 10.1016/j.msard.2022.104485
 - 55 Leite MI, Coutinho E, Lana-Peixoto M, Apostolos S, Waters P, Sato D, et al. Myasthenia gravis and neuromyelitis optica spectrum disorder: a multicenter study of 16 patients. *Neurology* 2012;78(20):1601–1607. Doi: 10.1212/WNL.0b013e31825644ff
 - 56 Song X, Chen J, Jin C, Peng Y, Sun Y, Zheng X. Inebilizumab treatment in a patient with co-occurring AQP4-IgG positive neuromyelitis optica spectrum disorder and myasthenia gravis: a case report and literature review. *Front Immunol* 2025; 15:1528989. Doi: 10.3389/fimmu.2024.1528989
 - 57 Yamamura T, Weinshenker B, Yeaman MR, De Seze J, Patti F, Lobo P, et al. Long-term safety of satralizumab in neuromyelitis optica spectrum disorder (NMOSD) from SAKuraSky and SAKuraStar. *Mult Scler Relat Disord* 2022;66:104025. Doi: 10.1016/j.msard.2022.104025
 - 58 Niino M, Isobe N, Araki M, Ohashi T, Okamoto T, Ogino M, et al. Clinical practice guidelines for multiple sclerosis, neuromyelitis optica spectrum disorder, and myelin oligodendrocyte glycoprotein antibody-associated disease 2023 in Japan. *Mult Scler Relat Disord* 2024;90:105829. Doi: 10.1016/j.msard.2024.105829
 - 59 Wingerchuk DM, Weinshenker BG, McCormick D, Barron S, Simone L, Jarzylo L. Aligning payer and provider strategies with the latest evidence to optimize clinical outcomes for patients with neuromyelitis optica spectrum disorder. *J Manag Care Spec Pharm* 2022;28(12-a, Suppl):S3–S27. Doi: 10.18553/jmcp.2022.28.12-a.s1
 - 60 Silva GD, Apóstolos-Pereira SL, Boaventura M, Paolilo RB, Matos A, Pitombeira MS, et al. Early rituximab versus escalating therapy in neuromyelitis optica: A cost and quality of life analysis. *Mult Scler Relat Disord* 2024;92:106160. Doi: 10.1016/j.msard.2024.106160
 - 61 Aungsumart S, Apiwattanakul M. Cost effectiveness of rituximab and mycophenolate mofetil for neuromyelitis optica spectrum disorder in Thailand: Economic evaluation and budget impact analysis. *PLoS One* 2020;15(02):e0229028. Doi: 10.1371/journal.pone.0229028
 - 62 Costanzi C, Matiello M, Lucchinetti CF, Weinshenker BG, Pittock SJ, Mandrekar J, et al. Azathioprine: tolerability, efficacy, and predictors of benefit in neuromyelitis optica. *Neurology* 2011;77(07):659–666. Doi: 10.1212/WNL.0b013e31822a2780
 - 63 Elson L, Kitley J, Luppe S, Lythgoe D, Mutch K, Jacob S, et al. Long-term efficacy, tolerability and retention rate of azathioprine in 103 aquaporin-4 antibody-positive neuromyelitis optica spectrum disorder patients: a multicentre retrospective observational study from the UK. *Mult Scler* 2014;20(11):1533–1540. Doi: 10.1177/1352458514525870
 - 64 Nikoo Z, Badihian S, Shaygannejad V, Asgari N, Ashtari F. Comparison of the efficacy of azathioprine and rituximab in neuromyelitis optica spectrum disorder: a randomized clinical trial. *J Neurol* 2017;264(09):2003–2009. Doi: 10.1007/s00415-017-8590-0
 - 65 Qiu W, Kermod AG, Li R, Dai Y, Wang Y, Wang J, et al. Azathioprine plus corticosteroid treatment in Chinese patients with neuromyelitis optica. *J Clin Neurosci* 2015;22(07):1178–1182. Doi: 10.1016/j.jocn.2015.01.028
 - 66 Xu Y, Wang Q, Ren HT, Qiao L, Zhang Y, Fei Y-Y, et al. Comparison of efficacy and tolerability of azathioprine, mycophenolate mofetil, and cyclophosphamide among patients with neuromyelitis optica spectrum disorder: A prospective cohort study. *J Neurol Sci* 2016;370:224–228. Doi: 10.1016/j.jns.2016.09.035
 - 67 Yang Y, Wang CJ, Wang BJ, Zeng ZL, Guo SG. Comparison of efficacy and tolerability of azathioprine, mycophenolate mofetil, and lower dosages of rituximab among patients with neuromyelitis optica spectrum disorder. *J Neurol Sci* 2018; 385:192–197. Doi: 10.1016/j.jns.2017.12.034
 - 68 Takai Y, Kuroda H, Misu T, Akaishi T, Nakashima I, Takahashi T, et al. Optimal management of neuromyelitis optica spectrum disorder with aquaporin-4 antibody by oral prednisolone maintenance therapy. *Mult Scler Relat Disord* 2021;49:102750. Doi: 10.1016/j.msard.2021.102750
 - 69 Watanabe S, Misu T, Miyazawa I, Nakashima I, Shiga Y, Fujihara K, Itoyama Y. Low-dose corticosteroids reduce relapses in neuromyelitis optica: a retrospective analysis. *Mult Scler* 2007;13(08):968–974. Doi: 10.1177/1352458507077189
 - 70 Kitley J, Elson L, George J, Waters P, Woodhall M, Vincent A, et al. Methotrexate is an alternative to azathioprine in neuromyelitis optica spectrum disorders with aquaporin-4 antibodies. *J Neurol Neurosurg Psychiatry* 2013;84(08):918–921. Doi: 10.1136/jnnp-2012-304774
 - 71 Ng ASL, Tan K. Methotrexate is effective for the treatment of neuromyelitis optica spectrum disorders in Asian patients. *Clin Exp Neuroimmunol* 2015;6:149–153. Doi: 10.1111/cen3.12189
 - 72 Chen H, Zhang Y, Shi Z, Feng H, Yao S, Xie J, Zhou H. The Efficacy and Tolerability of Mycophenolate Mofetil in Treating Neuromyelitis Optica and Neuromyelitis Optica Spectrum Disorder in Western China. *Clin Neuropharmacol* 2016;39(02):81–87. Doi: 10.1097/WNF.0000000000000131

- 73 Huang Q, Wang J, Zhou Y, Yang H, Wang Z, Yan Z, et al. Low-Dose Mycophenolate Mofetil for Treatment of Neuromyelitis Optica Spectrum Disorders: A Prospective Multicenter Study in South China. *Front Immunol* 2018;9:2066. Doi: 10.3389/fimmu.2018.02066
- 74 Huh SY, Kim SH, Hyun JW, Joung A-R, Park MS, Kim B-J, Kim HJ. Mycophenolate mofetil in the treatment of neuromyelitis optica spectrum disorder. *JAMA Neurol* 2014;71(11):1372–1378. Doi: 10.1001/jamaneurol.2014.2057
- 75 Jacob A, Matiello M, Weinshenker BG, Wingerchuk DM, Lucchinetti C, Shuster E, et al. Treatment of neuromyelitis optica with mycophenolate mofetil: retrospective analysis of 24 patients. *Arch Neurol* 2009;66(09):1128–1133. Doi: 10.1001/archneurol.2009.175
- 76 Jiao Y, Cui L, Zhang W, Zhang Y, Zhang X, Jiao J. Dose effects of mycophenolate mofetil in Chinese patients with neuromyelitis optica spectrum disorders: a case series study. *BMC Neurol* 2018;18(01):47. Doi: 10.1186/s12883-018-1056-x
- 77 Mealy MA, Wingerchuk DM, Palace J, Greenberg BM, Levy M. Comparison of relapse and treatment failure rates among patients with neuromyelitis optica: multicenter study of treatment efficacy. *JAMA Neurol* 2014;71(03):324–330. Doi: 10.1001/jamaneurol.2013.5699
- 78 Montcuquet A, Collongues N, Papeix C, Zephir H, Audoin B, Laplaud D, et al; NOMADMUS study group and the Observatoire Français de la Sclérose en Plaques (OFSEP) Effectiveness of mycophenolate mofetil as first-line therapy in AQP4-IgG, MOG-IgG, and seronegative neuromyelitis optica spectrum disorders. *Mult Scler* 2017;23(10):1377–1384. Doi: 10.1177/1352458516678474
- 79 Pathomrattanapibon C, Tisavipat N, Jitprapaikulsan J, Prayoon-wiwan N, Rattanathamsakul N, Siritho S. The efficacy and safety of mycophenolate mofetil in Thai neuromyelitis optica spectrum disorder patients. *Mult Scler Relat Disord* 2022;63:103882. Doi: 10.1016/j.msard.2022.103882
- 80 Zheng ZYSF, Liu X-R, Yang C-S, Sun L-X, Wang T, Yu C-Y. Efficacy of mycophenolate mofetil for AQP4 antibody-positive optic neuritis immunotherapy. *International Eye Science*. 2022;22(06):149–153
- 81 Huang W, Wang L, Xia J, Li W, Wang M, Yu J, et al; Pan-Yangtze River Delta Alliance for Demyelinating Disease. Efficacy and safety of azathioprine, mycophenolate mofetil, and reduced dose of rituximab in neuromyelitis optica spectrum disorder. *Eur J Neurol* 2022;29(08):2343–2354. Doi: 10.1111/ene.15355
- 82 Zhang L, Tian J, Dong X, Jia Z, Sun Y, Guo L, et al. Efficacy of azathioprine, mycophenolate mofetil, and rituximab in the treatment of neuromyelitis optica spectrum disorder and analysis of prognostic factors. *Neurol Sci* 2022;43(04):2651–2658. Doi: 10.1007/s10072-021-05609-0
- 83 Annovazzi P, Capobianco M, Moiola L, Patti F, Frau J, Uccelli A, et al. Rituximab in the treatment of Neuromyelitis optica: a multicentre Italian observational study. *J Neurol* 2016;263(09):1727–1735. Doi: 10.1007/s00415-016-8188-y
- 84 Bedi GS, Brown AD, Delgado SR, Usmani N, Lam BL, Sheremata WA. Impact of rituximab on relapse rate and disability in neuromyelitis optica. *Mult Scler* 2011;17(10):1225–1230. Doi: 10.1177/1352458511404586
- 85 Mealy MA, Kim SH, Schmidt F, López R, Arango JAJ, Paul F, et al. Aquaporin-4 serostatus does not predict response to immunotherapy in neuromyelitis optica spectrum disorders. *Mult Scler* 2018;24(13):1737–1742. Doi: 10.1177/1352458517730131
- 86 Longoni G, Banwell B, Filippi M, Yeh EA. Rituximab as a first-line preventive treatment in pediatric NMOSDs: Preliminary results in 5 children. *Neurol Neuroimmunol Neuroinflamm* 2014;1(04):e46. Doi: 10.1212/NXI.0000000000000046
- 87 Zhang C, Zhang M, Qiu W, Ma H, Zhang X, Zhu Z, et al; TANGO Study Investigators. Safety and efficacy of tocilizumab versus azathioprine in highly relapsing neuromyelitis optica spectrum disorder (TANGO): an open-label, multicentre, randomised, phase 2 trial. *Lancet Neurol* 2020;19(05):391–401. Doi: 10.1016/S1474-4422(20)30070-3
- 88 Araki M. Blockade of IL-6 signaling in neuromyelitis optica. *Neurochem Int* 2019;130:104315. Doi: 10.1016/j.neuint.2018.10.012
- 89 Du C, Zeng P, Han JR, Zhang T-X, Jia D, Shi F-D, Zhang C. Early Initiation of Tocilizumab Treatment Against Moderate-to-Severe Myelitis in Neuromyelitis Optica Spectrum Disorder. *Front Immunol* 2021;12:660230. Doi: 10.3389/fimmu.2021.660230
- 90 Lotan I, Charlson RW, Ryerson LZ, Levy M, Kister I. Effectiveness of subcutaneous tocilizumab in neuromyelitis optica spectrum disorders. *Mult Scler Relat Disord* 2020;39:101920. Doi: 10.1016/j.msard.2019.101920
- 91 Ringelstein M, Ayzenberg I, Lindenblatt G, Fischer K, Gahlen A, Novi G, et al; Neuromyelitis Optica Study Group (NEMOS) Interleukin-6 Receptor Blockade in Treatment-Refractory MOG-IgG-Associated Disease and Neuromyelitis Optica Spectrum Disorders. *Neurol Neuroimmunol Neuroinflamm* 2021;9(01):e1100. Doi: 10.1212/NXI.0000000000001100
- 92 Paolillo RB, Hachohen Y, Yazbeck E, Armangue T, Bruijstens A, Lechner C, et al. Treatment and outcome of aquaporin-4 antibody-positive NMOSD: A multinational pediatric study. *Neurol Neuroimmunol Neuroinflamm* 2020;7(05):e837. Doi: 10.1212/NXI.0000000000000837
- 93 Umeton RP, Waltz M, Aaen GS, Benson L, Gorman M, Goyal M, et al; US Network of Pediatric MS Centers. Therapeutic Response in Pediatric Neuromyelitis Optica Spectrum Disorder. *Neurology* 2023;100(09):e985–e994. Doi: 10.1212/WNL.0000000000201625
- 94 Zhou J, Zhang Y, Ji TY, Jin YW, Bao XH, Zhang YH, et al. [Clinical analysis of neuromyelitis optica spectrum disorders in childhood]. *Zhonghua Er Ke Za Zhi* 2019;57(02):118–124. Doi: 10.3760/cma.j.issn.0578-1310.2019.02.011
- 95 McKeon A, Lennon VA, Lotze T, Tenenbaum S, Ness JM, Rensel M, et al. CNS aquaporin-4 autoimmunity in children. *Neurology* 2008;71(02):93–100. Doi: 10.1212/01.wnl.0000314832.24682.c6
- 96 Fragomeni MO, Bichuetti DB, Oliveira EML. Pediatric-onset multiple sclerosis in Brazilian patients: Clinical features, treatment response and comparison to pediatric neuromyelitis optica spectrum disorders. *Mult Scler Relat Disord* 2018;25:138–142. Doi: 10.1016/j.msard.2018.07.036
- 97 Tenenbaum S, Yeh EAGuthy-Jackson Foundation International Clinical Consortium (GJCF-ICC) Pediatric NMOSD: A Review and Position Statement on Approach to Work-Up and Diagnosis. *Front Pediatr* 2020;8:339. Doi: 10.3389/fped.2020.00339
- 98 Ashtari F, Mehdipour R, Shaygannejad V, Asgari N. Pre-pregnancy, obstetric and delivery status in women with neuromyelitis optica spectrum disorder. *Mult Scler Relat Disord* 2020;44:102252. Doi: 10.1016/j.msard.2020.102252
- 99 Collongues N, Alves Do Rego C, Bourre B, Biotti D, Marignier R, Silva AMd, et al. Pregnancy in Patients With AQP4-Ab, MOG-Ab, or Double-Negative Neuromyelitis Optica Disorder. *Neurology* 2021;96(15):e2006–e2015. Doi: 10.1212/WNL.00000000000011744
- 100 Hadinejad M, Masoudi M, Sahraian MA, Mozdabadi RSK, Aliabadi HR, Shahmohammadi S, et al. Exploring the association of reproductive history with the development and course of neuromyelitis optica spectrum disorder. *Clin Neurol Neurosurg* 2022;219:107342. Doi: 10.1016/j.clineuro.2022.107342
- 101 Huang Y, Wang Y, Zhou Y, Huang Q, Sun X, Chen C, et al. Pregnancy in neuromyelitis optica spectrum disorder: A multicenter study from South China. *J Neurol Sci* 2017;372:152–156. Doi: 10.1016/j.jns.2016.11.054
- 102 Kim SH, Huh SY, Jang H, Park NY, Kim Y, Jung JY, et al. Outcome of pregnancies after onset of the neuromyelitis optica spectrum disorder. *Eur J Neurol* 2020;27(08):1546–1555. Doi: 10.1111/ene.14274
- 103 Kim W, Kim SH, Nakashima I, Takai Y, Fujihara K, Leite MI, et al. Influence of pregnancy on neuromyelitis optica spectrum disorder. *Neurology* 2012;78(16):1264–1267. Doi: 10.1212/WNL.0b013e318250d812

- 104 Shi B, Zhao M, Geng T, Qiao L, Zhao Y, Zhao X. Effectiveness and safety of immunosuppressive therapy in neuromyelitis optica spectrum disorder during pregnancy. *J Neurol Sci* 2017; 377:72–76. Doi: 10.1016/j.jns.2017.03.051
- 105 Shimizu Y, Fujihara K, Ohashi T, Nakashima I, Yokoyama K, Ikeguchi R, et al. Pregnancy-related relapse risk factors in women with anti-AQP4 antibody positivity and neuromyelitis optica spectrum disorder. *Mult Scler* 2016;22(11):1413–1420. Doi: 10.1177/1352458515583376
- 106 Natekar A, Pupco A, Bozzo P, Koren G. Safety of azathioprine use during pregnancy. *Can Fam Physician* 2011;57(12):1401–1402
- 107 Moretti ME, Verjee Z, Ito S, Koren G. Breast-feeding during maternal use of azathioprine. *Ann Pharmacother* 2006;40(12): 2269–2272. Doi: 10.1345/aph.1H152
- 108 Verberne EA, De Haan E, Van Tintelen JP, Lindhout D, Van Haelst MM. Fetal methotrexate syndrome: A systematic review of case reports. *Reprod Toxicol* 2019;87:125–139. Doi: 10.1016/j.reprotox.2019.05.066
- 109 Munoz SR, Lupattelli A, De Vries ST, Mol PGM, Nordeng H. Differences in medication beliefs between pregnant women using medication, or not, for chronic diseases: a cross-sectional, multinational, web-based study. *BMJ Open* 2020;10(02): e034529. Doi: 10.1136/bmjopen-2019-034529
- 110 Kim J-W, Suh C-H. The Safety of Medications During Pregnancy and Lactation in Patients with Inflammatory Rheumatic Diseases. *EMJ Rheumatology* 2021. Doi: 10.33590/emj/21-00017
- 111 Tektonidou MG, Andreoli L, Limper M, Amoura Z, Cervera R, Costedoat-Chalumeau N, et al. EULAR recommendations for the management of antiphospholipid syndrome in adults. *Ann Rheum Dis* 2019;78(10):1296–1304. Doi: 10.1136/annrheum-dis-2019-215213
- 112 Silva GD, Apóstolos-Pereira SL, Callegaro D. Estimated prevalence of AQP4 positive neuromyelitis optica spectrum disorder and MOG antibody associated disease in São Paulo, Brazil. *Mult Scler Relat Disord* 2023;70:104488. Doi: 10.1016/j.msard.2022.104488
- 113 Kümpfel T, Giehlhuber K, Aktas O, Ayzenberg I, Bellmann-Strobl J, Häußler V, et al; Neuromyelitis Optica Study Group (NEMOS) Update on the diagnosis and treatment of neuromyelitis optica spectrum disorders (NMOSD) - revised recommendations of the Neuromyelitis Optica Study Group (NEMOS). Part II: Attack therapy and long-term management. *J Neurol* 2024;271(01): 141–176. Doi: 10.1007/s00415-023-11910-z
- 114 Rojas JL, López PA, Criniti J, Pettinicchi JP, Caride A, Díaz EPC et al. Therapeutic strategies in NMOSD and MOGAD patients: A multicenter cohort study in Latin America. *Mult Scler Relat Disord* 2023;71:104508. Doi: 10.1016/j.msard.2023.104508
- 115 Compston A. Aids to the investigation of peripheral nerve injuries. Medical Research Council: Nerve Injuries Research Committee. His Majesty's Stationery Office: 1942; pp. 48 (iii) and 74 figures and 7 diagrams; with aids to the examination of the peripheral nervous system. By Michael O'Brien for the Guarantors of Brain. Saunders Elsevier: 2010; pp. [8] 64 and 94 Figures. *Brain* 2010;133(10):2838–2844. Doi: 10.1093/brain/awq270