

Best Practice Guidance for Adult Infusion Centres during the COVID-19 Pandemic: Report from the COVID-19 International Organization for the Study of IBD [IOIBD] Task Force.

Title: Best Practice Guidance for Adult Infusion Centres during the COVID-19 Pandemic: Report from the COVID-19 International Organization for the Study of IBD [IOIBD] Task Force.

Publication year: 2020

Journal: Journal of Crohn's & colitis

Authors: Division of Gastroenterology, Rabin Medical Center, Petah Tikva, Israel and the Sackler Faculty of Medicine, Tel Aviv University, Israel.

Society: European Respiratory Society

Origin country: International (Europe)

Target countries: Europe

DOI: 10.1093/ecco-jcc/jjaa147

PMID: 32959882

Source: <https://pubmed.ncbi.nlm.nih.gov/32959882/>

J Crohns Colitis J Crohns Colitis 3817 pheoup 101318676 eccojc Journal of Crohn's & Colitis 1873-9946 1876-4479 pmc-is-collection-domain yes pmc-collection-title Oxford University Press - PMC COVID-19 Collection PMC7543377 PMC7543377.1 7543377 7543377 32959882 10.1093/ecco-jcc/jjaa147 jjaa147 1 Original Article AcademicSubjects/MED00260 Best Practice Guidance for Adult Infusion Centres during the COVID-19 Pandemic: Report from the COVID-19 International Organization for the Study of IBD [IOIBD] Task Force Dotan Iris 1 1a Panaccione Remo 2 Kaplan Gilaad G 3 O'Morain Colm 4 Lindsay James O 5 Abreu Maria T 6 IOIBD 1 1 Division of Gastroenterology, Rabin Medical Center , Petah Tikva, Israel 1a Sackler Faculty of Medicine, Tel Aviv University , Israel 2 Inflammatory Bowel Disease Unit, Division of Gastroenterology and Hepatology, University of Calgary , Calgary, Canada 3 Research and Wellness Building, University of Calgary , Calgary, Canada 4 National Lead in Gastroenterology and Hepatology, Beacon Hospital , Sandymount, Dublin, Ireland 5 Department of Gastroenterology, The Royal London Hospital, Barts Health NHS Trust , Whitechapel, London, UK 6 Crohn's and Colitis Center, Professor of Medicine, Professor of Microbiology and Immunology, Division of Gastroenterology and Hepatology, University of Miami Miller School of Medicine , Miami, FL, USA 22 9 2020 366574 jjaa147 22 9 2020 08 10 2020 08 10 2020 30 03 2026 Copyright © 2020 European Crohn's and Colitis Organisation (ECCO). Published by Oxford University Press. All rights reserved. For permissions, please email: journals.permissions@oup.com 2020 This article is published and distributed under the terms of the Oxford University Press, Standard Journals Publication Model (https://academic.oup.com/journals/pages/open_access/funder_policies/chorus/standard_publication_model) This article is made available via the PMC Open Access Subset for unrestricted re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for the duration of the COVID-19 pandemic or until permissions are revoked in writing. Upon expiration of these permissions, PMC is granted a perpetual license to make this article available via PMC and Europe PMC, consistent with existing copyright protections. Abstract Infusion centres are a central part in the management of patients with inflammatory bowel disease [IBD] and could be a source of transmission of SARS-COV-2. Here we aimed to develop global guidance for best practices of infusion centres for IBD patients and to determine the impact of the COVID-19 pandemic on these centres. Under the auspices of the International Organization for the Study of Inflammatory Bowel Disease [IOIBD], a task force [TF] was formed, an online survey was developed to query infusion centre protocols during COVID-19, and recommendations were made, based on TF experience and opinion. Recommendations focus mainly on patients screening, infusion centres re-organization, personnel protection, and protocol modifications such as shortening infusion duration or replacing it with

subcutaneous alternatives. Implementing these recommendations will hopefully reduce exposure of both IBD patients and care givers to SARS-COV-2 and improve the function and safety of infusion centres during the COVID-19 pandemic as well as potential future threats.