

Clinical care of pregnant and postpartum women with COVID-19: Living recommendations from the National COVID-19 Clinical Evidence Taskforce.

Title: Clinical care of pregnant and postpartum women with COVID-19: Living recommendations from the National COVID-19 Clinical Evidence Taskforce.

Publication year: 2020

Journal: The Australian & New Zealand journal of obstetrics & gynaecology

Authors: Maternal, Child and Adolescent Health Program, Burnet Institute, Melbourne, Victoria, Australia.

Society: European Society of Cardiology

Origin country: International (Europe)

Target countries: Europe

DOI: 10.1111/ajo.13270

PMID: 33119139

Source: <https://pubmed.ncbi.nlm.nih.gov/33119139/>

Aust N Z J Obstet Gynaecol Aust N Z J Obstet Gynaecol 379 blackwellopen 0001027 AJO The Australian & New Zealand Journal of Obstetrics & Gynaecology 0004-8666 1479-828X pmc-is-collection-domain yes pmc-collection-title Wiley Open Access Collection PMC7820999 PMC7820999.1 7820999 7820999 33119139 10.1111/ajo.13270 AJO13270 1 Position Paper Position Papers Clinical care of pregnant and postpartum women with COVID-19: Living recommendations from the National COVID-19 Clinical Evidence Taskforce Clinical care of pregnant and postpartum women with COVID-19 J. P. Vogel et al . Vogel Joshua P. <https://orcid.org/0000-0002-3214-7096> 1 2 joshua.vogel@burnet.edu.au Tendal Britta <https://orcid.org/0000-0001-8354-2603> 2 Giles Michelle <https://orcid.org/0000-0003-3007-1822> 3 4 5 6 7 Whitehead Clare <https://orcid.org/0000-0001-8267-1630> 5 8 Burton Wendy 9 Chakraborty Samantha <https://orcid.org/0000-0002-9708-4532> 10 Cheyne Saskia <https://orcid.org/0000-0001-5061-2185> 11 Downton Teena 12 Fraile Navarro David <https://orcid.org/0000-0002-1108-7071> 13 Gleeson Glenda 14 Gordon Adrienne <https://orcid.org/0000-0002-2328-1066> 15 16 17 Hunt Jenny 18 Kitschke Jackie 19 McDonald Steven <https://orcid.org/0000-0003-2832-5205> 2 McDonnell Nolan 20 Middleton Philippa <https://orcid.org/0000-0002-8573-338X> 21 22 Millard Tanya <https://orcid.org/0000-0001-7135-3231> 2 Murano Melissa <https://orcid.org/0000-0001-6354-2729> 2 Oats Jeremy <https://orcid.org/0000-0003-3808-9808> 23 Tate Rhiannon 2 White Heath <https://orcid.org/0000-0002-4147-5278> 2 Elliott Julian <https://orcid.org/0000-0002-9165-5875> 2 3 Roach Vijay 24 Homer Caroline S.E. <https://orcid.org/0000-0002-7454-3011> 1 25 the National COVID-19 Clinical Evidence Taskforce McGowan Sharon Ballenden Nicola Barrett Terri Lee Beavis Vanessa Saunders James Beckford Buchanan Tanya Buchanan Grey Marina Casey Dawn Cowie Marita Doyle Joseph Frydenberg Mark Gnjidic Danijela Green Sally Greenland Rohan Griffin Ken Groombridge Stephan Hardy Louise Hodak Alison Holley Anthony Jovanovska Vase Knight Sabina Michaels Kristin Morley Peter Morphet Julia Nou Suzi Russo Phillip Sarson Megan Young Alan Norris Sarah Morris Donovan Bronwyn Gurry Sharon Hudson Eloise Hurley Shauna Primmer Declan Timms Samantha Whicker Susan Mukherjee Sutapa Agostino Jason Booth Karen Burr Lucy Byers Lyn Cameron Peter Cooper Megan Cheng Allen Fowler Peter Frydenberg Mark Glanville Alan Leder Karin McGloughlin Steve McMullan Brendan McPhee Ewen Mitchell Brett Morgan Mark Myles Paul O'Donnell Chris Parr Michael Phillips Jane Randall Rebecca Varndell Wayne Whyte Ian William Leeroy Randall Rebecca Brightwell Richard Condon Lynda Deshpande Amrita Ehm Adam Ferrie Monica Muller Joanne Pullin Lara Robinson Elizabeth Witt Adele Larkins Sarah Morgan Mark Taylor Georgina Agostino Jason Burgess Paul Burns Penny Byers Lyn Douglas Kirsty Ewald Ben Ewald Dan Fornasier Dianna Knight Sabina Nelson Carmel Peachey Louis Peiris David Driel Mieke Walters Lucie Weaver Ineke Burr Lucy Hendel Simon Shekar Kiran Avard Bronwyn Cairns Kelly Glanville Allan Gilroy Nicky Myles Paul O'Sullivan Robert Robinson

Owen Sharland Chantal McCarthy Sally Wark Peter McGoughlin Steve Nair Priya Hodgson Carol Ankravs Melissa French Craig Hansen Kim Huckson Sue Iredell Jon Janerka Carrie Jaspers Rose Litton Ed Macdonald Stephen Peake Sandra Seppelt Ian Bowen Asha McMullan Brendan Tingay David Vasilunas Nan Anderson Lorraine Best James Burns Penny Craig Simon Erickson Simon Fancourt Nick Goff Zoy Kapuya Vimbai Keyte Catherine Malyon Lorelle Wurzel Danielle Agar Meera Lindley Richard Smallwood Natasha Callary Mandy Chapman Michael Good Philip Jenkin Peter Morgan Deidre Naganathan Vasi Srikanth Velandai Tuffin Penny Whiting Elizabeth William Leroy Yates Patsy Barber Bridget Davies Jane Davis Josh Ewald Dan Gwee Amanda Leder Karin Matthews Gail McMahon James Peel Trisha Raftery Chris Rees Megan Roberts Jason Seppelt Ian Snelling Tom Wibrow Brad Baker Ross Curnow Jennifer Cutts Briony Enjeti Anoop Forbes Andrew Ho Prahlad Holyoak Adam Liley Helen McFadyen James McQuilten Zoe Merriman Eileen Savoia Helen Tan Chee Wee Tran Huyen Ward Chris Williams Katrina Ballard Neil Bendall Samantha Bhandari Neel Byers Lyn Craig Simon Ellis Dan Ewald Dan Fairley Craig Hoggard Brett Cong Minh Le Morley Peter Nair Priya Pearce Andrew Turner Tari Callesen Henriette Campbell Sue Ring Jenny Wilson Agnes Henry David Pearson Sallie Boyle Douglas Chidwick Kendal Chapman Wendy French Craig Pearce Chris Snelling Tom Bero Lisa Grundy Quinn Lexchin Joel Mintzes Barbara * 1 Maternal, Child and Adolescent Health Program Burnet Institute Melbourne Victoria Australia 2 Cochrane Australia School of Public Health and Preventive Medicine Monash University Melbourne Victoria Australia 3 Alfred Hospital Melbourne Victoria Australia 4 Monash Health Melbourne Victoria Australia 5 Royal Women's Hospital Melbourne Victoria Australia 6 Sunshine Hospital Melbourne Victoria Australia 7 Department of Obstetrics and Gynaecology Monash University Melbourne Victoria Australia 8 Department of Obstetrics and Gynaecology University of Melbourne Melbourne Victoria Australia 9 Morningside General Practice Clinic Brisbane Queensland Australia 10 Department of General Practice School of Primary and Allied Health Care Monash University Melbourne Victoria Australia 11 NHMRC Clinical Trials Centre University of Sydney Sydney New South Wales Australia 12 Australian College of Rural and Remote Medicine Brisbane Queensland Australia 13 Australian Institute of Health Innovation Macquarie University Sydney New South Wales Australia 14 Central Australia Primary and Public Health ■ Midwifery and Women's Health Alice Springs Northern Territory Australia 15 RPA Newborn Care Sydney Local Health District Discipline of Obstetrics, Gynaecology and Neonatology Central Clinical School Faculty of Medicine and Health University of Sydney Sydney New South Wales Australia 16 Charles Perkins Centre University of Sydney Sydney New South Wales Australia 17 Sydney Institute for Women, Children and their Families Sydney Local Health District Sydney New South Wales Australia 18 Victorian Aboriginal Health Service Melbourne Victoria Australia 19 Australian College of Midwives representative, Midwifery Group Practice Women's and Children's Hospital Adelaide South Australia Australia 20 Faculty of Health and Medical Sciences Obstetrics and Gynaecology University of Western Australia Perth Western Australia Australia 21 SAHMRI, Women and Children's Hospital Adelaide South Australia Australia 22 Faculty of Medical and Health Sciences The University of Adelaide Adelaide South Australia Australia 23 Melbourne School of Population and Global Health University of Melbourne Melbourne Victoria Australia 24 North Shore Private Hospital Sydney New South Wales Australia 25 Centre for Midwifery, Child and Family Health in the Faculty of Health University of Technology Sydney Sydney New South Wales Australia * Correspondence: A/Prof Joshua P. Vogel, Maternal, Child and Adolescent Health Program, Burnet Institute, Melbourne, Australia. Email: joshua.vogel@burnet.edu.au * See Acknowledgements section.

29 10 2020 12 2020 60 6 373552 10.1111/ajo.v60.6 840 851 27 9 2020 29 9 2020 22 01 2021 26 01 2021 24 09 2025 © 2020 The Authors. Australian and New Zealand Journal of Obstetrics and Gynaecology published by John Wiley & Sons Australia, Ltd on behalf of Royal Australian and New Zealand College of Obstetricians and Gynaecologists
<https://creativecommons.org/licenses/by-nc-nd/4.0/> This is an open access article under the terms of the <http://creativecommons.org/licenses/by-nc-nd/4.0/> License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made. To date, 18 living recommendations for the clinical care of pregnant and postpartum women with COVID-19 have been issued by the National COVID-19 Clinical Evidence Taskforce. This includes recommendations on mode of birth, delayed

umbilical cord clamping, skin-to-skin contact, breastfeeding, rooming-in, antenatal corticosteroids, angiotensin-converting enzyme inhibitors, disease-modifying treatments (including dexamethasone, remdesivir and hydroxychloroquine), venous thromboembolism prophylaxis and advanced respiratory support interventions (prone positioning and extracorporeal membrane oxygenation). Through continuous evidence surveillance, these living recommendations are updated in near real-time to ensure clinicians in Australia have reliable, evidence-based guidelines for clinical decision-making. Please visit <https://covid19evidence.net.au/> for the latest recommendation updates.

pmc-status-qastatus 0 pmc-status-live yes pmc-status-embargo no pmc-status-released yes
pmc-prop-open-access yes pmc-prop-olp no pmc-prop-manuscript no pmc-prop-legally-suppressed
no pmc-prop-has-pdf yes pmc-prop-has-supplement no pmc-prop-pdf-only no
pmc-prop-suppress-copyright no pmc-prop-is-real-version no pmc-prop-is-scanned-article no
pmc-prop-preprint no pmc-prop-in-epmc yes pmc-license-ref CC BY-NC-ND
source-schema-version-number 2.0 cover-date December 2020 details-of-publishers-converter
Converter:WILEY_ML3GV2_TO_JATSPMC version:5.9.6 mode:remove_FC converted:22.01.2021
Conflict of interest : Caroline Homer and Adrienne Gordon are investigators on NHMRC's Centre
for Research Excellence Grant for Stillbirth. Michelle Giles ■ Gilead Sciences (Remdesivir). The
other named authors have no conflict of interest to declare.

The first cases of novel coronavirus SARS-CoV-2 infection were reported in December 2019 in Wuhan, China. 1 Novel coronavirus disease 2019 (COVID-19) was declared a global pandemic by the World Health Organization on 11 March 2020, and as of 17 September 2020, there have been over 3068 million confirmed cases and more than 954 000 deaths globally. 2 In Australia, there have been 26 942 confirmed cases and 854 deaths. 3 COVID-19 commonly presents with fever and respiratory symptoms including cough and shortness of breath; other symptoms (such as myalgia, sore throat, fatigue, diarrhoea and anosmia) are less common. 4 An estimated 15% of infected individuals will develop severe disease (requiring oxygen support) and 5% will progress to critical disease. 4 Critical COVID-19 disease is characterised by respiratory failure, acute respiratory distress syndrome (ARDS), thromboembolism, sepsis, shock or multiorgan failure. After recovery from acute illness, some individuals may experience longer-term sequelae, particularly respiratory complications. 5

Experience from previous pandemics (H1N1) and coronavirus outbreaks (SARS and MERS) raised concerns that pregnant women may be particularly vulnerable to more severe COVID-19 disease. 6 , 7 While current data are limited, pregnant and recently pregnant women with COVID-19 are less likely to experience fever and myalgia, compared with non-pregnant women of reproductive age. 8 However, they have higher rates of intensive care unit admission or invasive ventilation, and their babies are more likely to be born preterm or require neonatal unit admission. 8 Epidemiological data from the Latin American region suggest maternal deaths may be increasing due to COVID-19. 9 Vertical transmission appears possible although probably rare. 10 , 11

In response to the COVID-19 pandemic, state, national and international organisations rapidly developed clinical guidelines on how best to care for pregnant and postpartum women with COVID-19. 12 Ideally, clinical guidelines are developed using systematic reviews of available evidence to address pre-specified questions, with expert panels formulating recommendations based on this evidence. However, many such recommendations become outdated as new evidence emerges. 13 , 14 This is a particular challenge in the COVID-19 context, where the volume of COVID-19-related literature is increasing dramatically. 15 Living guideline methodologies were developed to meet such challenges. 16 In living guidelines, digital technologies and continuous evidence surveillance are used to continually update systematic reviews and recommendations, so as to reflect the latest available evidence. 17 This approach has been used successfully in stroke, diabetes, maternal health and other topics. 18 , 19 , 20

The National COVID-19 Clinical Evidence Taskforce was established in March 2020 to produce living recommendations on the clinical care of people with COVID-19 in Australia. 21 As of 17

September 2020, the Taskforce has published 72 recommendations on disease severity, disease-modifying treatments, chemoprophylaxis, respiratory support, venous thromboembolism (VTE) prophylaxis, and use of therapies for pre-existing conditions, as well as specific recommendations on the care of women, neonates, children, adolescents, older people and people in palliative care with COVID-19. The aim of this article is to describe the Taskforce recommendations for the care of pregnant and postpartum women with COVID-19.

The living guidelines are being developed according to Australian National Health and Medical Research Council (NHMRC) standards, including the use of GRADE (Grading of Recommendations Assessment, Development and Evaluation) methodology. 22 The Taskforce brings together over 220 clinical experts from across Australia to assess the latest evidence and keep recommendations up-to-date on a weekly basis. All outputs are overseen by a Steering Committee composed of representatives from 31 peak health professional bodies, including the Australian College of Midwives (ACM) and the Royal Australian and New Zealand College of Obstetricians and Gynaecologists (RANZCOG). 23 The Taskforce's remit is to focus on the clinical care of people with COVID-19, with other national agencies responsible for producing guidance on infection prevention, control and diagnosis. The Taskforce also produces clinical flowcharts to facilitate translation of living recommendations into clinical practice. 24 As these living recommendations are regularly updated with new evidence, we encourage clinicians to visit <https://covid19evidence.net.au/> and freely subscribe to the weekly updates.

The Guidelines Leadership Group (GLG) oversees seven expert panels developing recommendations for primary, hospital and critical care settings. An evidence review team composed of experienced systematic reviewers and guideline methodologists identifies, appraises and synthesises evidence to inform the development of these recommendations. The Taskforce uses the MAGIC (Making GRADE the Irresistible Choice) digital platform for formulating and publishing recommendations. 25 The multi-disciplinary Pregnancy and Perinatal Care panel (one of seven panels) is composed to ensure diverse representation of setting, culture, geography and gender. It includes 13 experts from Aboriginal and Torres Strait Islander health, general practice, infectious diseases, midwifery, neonatology, obstetrics and gynaecology, obstetric anaesthesia, perinatal epidemiology, public health, and rural and remote medicine. Recommendations formulated by any panel are reviewed and approved by the GLG and the Steering Committee, prior to their publication.

We conducted a systematic mapping of available guidelines on the antenatal, intrapartum and postpartum care of women during the COVID-19 pandemic, identifying 81 guidelines from 48 organisations in Australia and internationally. 12 This mapping informed a prioritisation by panel members, who ranked a shortlist of clinical questions on the basis of: (1) interventions that are likely to impact on patient outcomes; (2) proportion of clinical population impacted; (3) extent of variation in current practice; and (4) likelihood of new evidence emerging. The panel also reviewed existing Taskforce recommendations on the clinical care of adults with COVID-19 to identify those clinical questions where adaptation was needed.

An information specialist oversees the evidence surveillance process, which combines daily searches of COVID-19-related articles and preprints across broad topic areas (such as pregnancy and newborn care) with targeted searches for specific clinical questions as they are proposed by the expert panels. 26 The evidence review team uses Covidence, 27 a collaborative online software for managing and streamlining systematic reviews, for screening literature. Titles and abstracts are initially screened by two reviewers independently, with full texts of potentially eligible studies retrieved and also reviewed in duplicate. Disagreements are resolved through discussion or consultation with a third member of the evidence review team. Eligible articles are primary research (typically trials) or systematic reviews relating to the clinical questions of interest. Systematic reviews (particularly living systematic reviews that are kept up-to-date) of reasonable quality are the preferred evidence source. If required, a limited update may be used to integrate findings from more recent primary studies into an existing systematic review. Where no such systematic review exists, the evidence review team initiates a new systematic review. For some

clinical questions where direct trial evidence is unlikely to be available for pregnant and postpartum women, lower quality evidence such as observational studies or studies from related topics (eg, studies of women with ARDS due to other causes) are considered.

Data on study characteristics and priority outcomes are extracted in duplicate using standardised data collection forms, with disagreements resolved through discussion or consulting a third reviewer. 27 Included studies are assessed using an appropriate quality assessment tool, such as AMSTAR 2 for systematic reviews and the Cochrane Risk of Bias 2.0 assessment tool for randomised trials. 28 , 29 , 30

Since 14 May 2020, the panel convened on a weekly basis to review the evidence, develop and update recommendations. Where necessary, assistance is sought from additional expert advisory groups (eg, haematology, critical care). The panel reviews evidence profiles on the benefits, harms and certainty of the evidence for priority questions, as well as considering the resource requirements, cost-effectiveness, feasibility, acceptability and equity of each intervention. This is done systematically using GRADE evidence to decision frameworks. 22 Recommendations are formulated in line with established Australian maternity care principles, including the need for evidence-based, person-centred care that is responsive to women's needs and preferences. 31 Recommendations are also reviewed by a separate Consumer Panel, co-convened by the Consumers Health Forum of Australia and the Taskforce. Ongoing literature surveillance is maintained for published recommendations. If relevant new studies become available, evidence profiles are updated and the recommendations may be revised.

The following recommendations are current as of 17 September 2020, with any subsequent updates freely available at <https://covid19evidence.net.au/> . Classifications of disease severity used by the Taskforce are described in Table 1 .

Classification of COVID-19 disease severity †

Adults not presenting any clinical features suggestive of moderate or severe disease or a complicated course of illness.

or mild upper respiratory tract symptoms

or cough, new myalgia or asthenia without new shortness of breath or a reduction in oxygen saturation

Stable adult patient presenting with respiratory and/or systemic symptoms or signs. Able to maintain oxygen saturation above 92% (or above 90% for patients with chronic lung disease) with up to 4 L/min oxygen via nasal prongs.

prostration, severe asthenia, fever > 38°C or persistent cough

clinical or radiological signs of lung involvement

no clinical or laboratory indicators of clinical severity or respiratory impairment

Adult patients meeting any of the following criteria:

arterial partial pressure of oxygen (PaO₂)/ inspired oxygen fraction (FiO₂) ≤300

Adult patient meeting any of the following criteria:

Occurrence of severe respiratory failure ($\text{PaO}_2/\text{FiO}_2 < 200$), respiratory distress or acute respiratory distress syndrome (ARDS). This includes patients deteriorating despite advanced forms of respiratory support (noninvasive ventilation, high-flow nasal oxygen) OR patients requiring mechanical ventilation.

OR other signs of significant deterioration

Recommendations were current at the time of writing. Please visit The National COVID-19 Clinical Evidence Taskforce (<https://covid19evidence.net.au/>) for the latest updates to the recommendations.

John Wiley & Sons, Ltd Pregnancy and perinatal care of women with COVID-19 Seven recommendations on mode of birth, delayed umbilical cord clamping, skin-to-skin contact, breastfeeding, rooming-in, antenatal corticosteroids and angiotensin-converting enzyme (ACE) inhibitors support continuing with the provision of usual care, although the need for masks and hand hygiene when mothers are caring and feeding their babies, is emphasised (Table 2).

Pregnancy and perinatal care recommendations for women with COVID-19 †

For pregnant women with COVID-19, mode of birth should remain as per usual care (conditional recommendation, very low certainty evidence)

There is currently no evidence to indicate that caesarean section for women with COVID-19 reduces the risk of vertical transmission to the newborn. Mode of birth should continue as per usual care. Respiratory deterioration due to COVID-19 may prompt urgent delivery on an individual basis.

Delayed umbilical cord clamping is supported as part of standard care, independent of the presence of COVID-19 (consensus recommendation)

There is currently no evidence that delayed umbilical cord clamping affects the risk of vertical transmission of COVID-19.

Early skin-to-skin contact after birth and during the postnatal period is supported, irrespective of the presence of COVID-19. However, parents with COVID-19 should use infection prevention and control measures (mask and hand hygiene) (consensus recommendation)

Early skin-to-skin contact refers to placing the naked baby prone on the parent's bare chest immediately after birth.

Skin-to-skin contact should be encouraged and continue as per usual practice in other postnatal and neonatal settings, such as neonatal intensive care unit and postnatal wards, providing infection prevention and control measures are maintained.

Breastfeeding is supported irrespective of the presence of COVID-19. However, women with COVID-19 who are breastfeeding should use infection prevention and control measures (mask and hand hygiene) while infectious (conditional recommendation, very low certainty evidence)

There is currently no evidence to indicate that breastfeeding increases the risk of vertical transmission to the newborn. As there are substantial known benefits for breastfeeding, women should be supported to initiate or continue breastfeeding. If the baby is being fed with expressed breastmilk or formula, these same infection prevention and control measures should be used.

For women with COVID-19 who have given birth, support rooming-in of mother and newborn in the birth suite and on the postnatal ward when both mother and baby are well. However, women with COVID-19 should use infection prevention and control measures (mask and hand hygiene) (conditional recommendation, very low certainty evidence)

There is currently no evidence to indicate that a woman with a known COVID-19 infection should be separated from her newborn to prevent transmission. As there are substantial known benefits for keeping mother and newborn together postpartum, women should be supported to be with their newborn as per usual care.

Women with COVID-19 should be encouraged and supported in using good hand hygiene before and after handling their baby, and using a mask while in close contact with their baby. To the extent possible, these women should practice physical distancing when not feeding or caring for the baby.

The use of antenatal corticosteroids for women at risk of preterm birth is supported as part of standard care, independent of the presence of COVID-19 (consensus recommendation)

There are clear benefits to using antenatal corticosteroids for women at risk of preterm birth at less than 34 weeks gestation. There is currently no evidence to suggest that antenatal corticosteroids cause additional maternal or fetal harm in the setting of COVID-19 when used for this indication. They should therefore be given where indicated.

The Taskforce has separate recommendations regarding the use of dexamethasone as a disease-modifying treatment in pregnant or breastfeeding women for COVID-19. Women with COVID-19 who are on oxygen and receiving dexamethasone do not require additional doses of corticosteroids for fetal lung maturation.

In postpartum women with COVID-19 who have hypertension requiring treatment with ACE inhibitors, there is currently no evidence to deviate from usual care. These medications should be initiated or continued unless otherwise contraindicated (consensus recommendation)

ACE inhibitors are contraindicated in the antenatal period due to risk of fetal and neonatal harm.

Recommendations were current at the time of writing. Please visit The National COVID-19 Clinical Evidence Taskforce (<https://covid19evidence.net.au/>) for the latest updates to the recommendations. ACE, angiotensin-converting enzyme

John Wiley & Sons, Ltd While systematic reviews on the risks of vertical transmission by mode of birth, rooming-in status and different infant feeding approaches have been published, only very low certainty evidence (predominantly case reports and case series) is currently available. 32 , 33 Additional evidence is available from a US multi-centre cohort study of 106 neonates born to mothers who were positive for COVID-19. 34 In this study, mode of birth, skin-to-skin contact and breastfeeding were managed as per usual care, although maternal use of surgical masks, hand hygiene and breast cleansing were encouraged. Newborns roomed-in with mothers, except for 17 newborns who were separated at parental request or due to medical complications. Newborns were tested by nasopharyngeal polymerase chain reaction at 12–24 h (n = 106), 5–7 days (n = 79) and 14 days (n = 72) of life, with all newborns testing negative at all timepoints. While the possibility of vertical transmission cannot be excluded based on this evidence, this must be weighed against the known, substantial benefits of these interventions. 35 , 36 , 37 , 38 , 39 , 40 , 41 , 42

No direct evidence was identified on delayed cord clamping in women with COVID-19, or on antenatal corticosteroids or use of ACE inhibitors for postpartum women with hypertension requiring drug treatment. Hence, these recommendations are aligned with current Australian pregnancy and perinatal care guidance on use of these interventions. 43 , 44 , 45

Evidence on corticosteroids compared with standard care for COVID-19 in adults comes from eight randomised trials (over 5700 patients), including the multi-centre RECOVERY trial that evaluated dexamethasone against usual care in 2104 patients. 46 , 47 Pregnant women were specifically excluded from all trials, except for RECOVERY (six pregnant women enrolled) and REMAP-CAP (the number of pregnant women enrolled has not been reported). 47 , 48 Corticosteroids reduce mortality for adult patients with severe or critical COVID-19 (relative risk (RR) 0.86 95% CI 0.77–0.97), and probably also reduce the need for mechanical ventilation or death, and time to discharge from hospital. However, in patients who do not require oxygen, corticosteroids may increase mortality (RR 1.27 95% CI 1.00–1.61). Considering these benefits and the likely acceptability of the intervention, using dexamethasone 6 mg daily (IV or oral) for up to ten days is recommended in pregnant or breastfeeding women with COVID-19 who are receiving oxygen (Table 3). While the RECOVERY Trial specified that enrolled pregnant women receive oral prednisolone or intravenous hydrocortisone, 49 dexamethasone is preferred as it is used for other clinical indications in pregnancy. 50 Corticosteroids should not be used in women with COVID-19 who do not require oxygen, due to the lack of benefit and possible harms in this subgroup.

Recommendations on use of disease-modifying treatments for pregnant or postpartum women with COVID-19 †

Use dexamethasone 6 mg daily intravenously or orally for up to ten days in pregnant or breastfeeding women with COVID-19 who are receiving oxygen (including mechanically ventilated patients) (strong recommendation, low certainty evidence) .

Do not routinely use dexamethasone to treat COVID-19 in pregnant or breastfeeding women who do not require oxygen (strong recommendation, low certainty evidence) .

Antenatal corticosteroids should still be used for fetal lung maturation in pregnant women at risk of preterm birth who also have COVID-19. Dexamethasone should still be used for other evidence-based indications in pregnant and breastfeeding women who have COVID-19.

Use of remdesivir for pregnant or breastfeeding women with COVID-19 outside of a trial setting should not be considered routinely (conditional recommendation, very low certainty evidence) .

As pregnant and breastfeeding women are often excluded from clinical trials, use of remdesivir in this population would be outside a clinical trial setting. Pregnant and breastfeeding women receiving remdesivir should nonetheless be enrolled in national COVID-19 registries. Currently, there is no direct evidence of the effects of remdesivir in pregnant and breastfeeding women. Information about the patients and the intervention (dosages, duration) in the trials used for this recommendation can be found in the Practical info tab.

Due to antagonism observed in vitro , concomitant use of remdesivir with chloroquine or hydroxychloroquine is not recommended. 90

Do not use hydroxychloroquine for the treatment of COVID-19 (strong recommendation, moderate certainty evidence) .

This recommendation applies to adults, children and adolescents, pregnant and breastfeeding women, older people living with frailty and those receiving palliative care.

Use of hydroxychloroquine may still be considered in the context of randomised trials with appropriate ethical approval, such as combination therapies that include hydroxychloroquine.

For people exposed to individuals with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection, do not use hydroxychloroquine for post-exposure prophylaxis outside of randomised trials with appropriate ethical approval (strong recommendation, low certainty

evidence) .

Trials are needed in special populations, including children and adolescents, pregnant and breastfeeding women, older people living with frailty and those receiving palliative care. Until further evidence is available, do not use hydroxychloroquine for post-exposure prophylaxis in these populations unless they are eligible to be enrolled in trials.

For people with COVID-19, do not use the other disease-modifying treatments ‡ outside of randomised trials with appropriate ethical approval (strong recommendation, very low certainty evidence) .

Trials are needed in special populations, including children and adolescents, pregnant and breastfeeding women, older people living with frailty and those receiving palliative care. Until further evidence is available, do not use in these populations unless they are eligible to be enrolled in trials.

Recommendations were current at the time of writing. Please visit The National COVID-19 Clinical Evidence Taskforce (<https://covid19evidence.net.au/>) for the latest updates to the recommendations.

The treatments include: aprepitant, baloxavir marboxil, calcifediol, chloroquine, colchicine, convalescent plasma, darunavir-cobicistat, favipiravir, human mesenchymal stem cells, immunoglobulin plus methylprednisolone, interferon β -1a, interferon β -1b, interferon gamma, lopinavir-ritonavir, ruxolitinib, sofosbuvir-daclatasvir, telmisartan and umifenovir.

John Wiley & Sons, Ltd Remdesivir Four trials have compared remdesivir with standard care in hospitalised, non-pregnant adults with COVID-19. 51 , 52 , 53 , 54 As pregnant and breastfeeding women were excluded from these trials, there is no direct evidence on the impact of remdesivir for COVID-19 in these women, as well as a lack of safety data on its use in pregnancy. 55 In non-pregnant adults, remdesivir may decrease time to recovery by a few days (hazards ratio (HR) 1.24, 95% CI 1.08–1.42) and time to improvement only slightly (HR 1.17, 95% CI 1.00–1.38). Its effect on mortality is uncertain, and it is currently unclear which regimen (five days or ten days) provides the most benefit. However, the panel noted that severity of disease is an important factor – for pregnant women with severe or critical COVID-19, the harm-to-benefit ratio may differ compared to pregnant women with mild or moderate disease. The Taskforce therefore made a conditional recommendation against the routine use of remdesivir in pregnant and breastfeeding women outside of a trial setting (Table 3).

There are currently 11 trials (over 6300 patients) comparing hydroxychloroquine with standard care in the treatment of patients with COVID-19. 56 , 57 , 58 , 59 , 60 , 61 , 62 , 63 , 64 , 65 , 66 In all trials, pregnant women were either excluded or their eligibility was not specified. Meta-analyses indicate that hydroxychloroquine is potentially harmful and no more effective than standard care, leading to a strong recommendation against its use for COVID-19 treatment (Table 3). Separately, hydroxychloroquine for post-exposure prophylaxis has been evaluated in two trials (3135 participants), both of which excluded pregnant and breastfeeding women; however, there is currently insufficient evidence to assess benefits. 67 , 68 The Taskforce recommends that hydroxychloroquine for post-exposure prophylaxis should not be used outside of randomised trials with appropriate ethical approval, emphasising that further trials (including in pregnant and breastfeeding women) are needed.

The Taskforce has issued a further 19 recommendations on a wide range of disease-modifying treatments (Table 3), including: aprepitant, azithromycin, baloxavir marboxil, calcifediol, chloroquine, colchicine, convalescent plasma, darunavir-cobicistat, favipiravir, human mesenchymal stem cells, immunoglobulin plus methylprednisolone, interferon β -1a, interferon β -1b, interferon gamma, lopinavir-ritonavir, ruxolitinib, sofosbuvir-daclatasvir, telmisartan and umifenovir. These treatments have typically been evaluated in one or two trials with a small number

of nonpregnant participants, resulting in low or very low certainty evidence from which no conclusions can be drawn. For some of these treatments (such as baloxavir marboxil or favipiravir), no pregnancy safety data are available, whereas others (such as azithromycin or lopinavir/ritonavir) have a well-described safety profile. 55 The Taskforce therefore recommends that these treatments should not be used outside of randomised trials, with emphasis that trials enrolling pregnant and breastfeeding women are needed.

While COVID-19 infection appears to increase the risk of VTE, there is currently no direct evidence on the exact duration of this increased risk, or which prophylactic anticoagulant regimen is optimal. 69 The consensus recommendations (Table 4) were formulated based on a review of available international guidelines 69 , 70 and consultation with expert haematologists. Recognising that pregnancy and the postpartum period are inherently hypercoagulable states, the statement emphasises that all pregnant and postpartum women should be assessed for additional risk factors for VTE on admission to hospital, when COVID-19 is diagnosed, if COVID-19 severity changes, and postpartum.

Venous thromboembolism (VTE) prophylaxis in pregnant and postpartum women †

For pregnant or postpartum women who are admitted to hospital (for any indication) and who have COVID-19, use prophylactic doses of anticoagulants, preferably low-molecular weight heparin (LMWH: eg, enoxaparin 40 mg once daily or dalteparin 5000 IU once daily) unless there is a contraindication, such as risk for major bleeding or imminent birth.

Prophylactic anticoagulants should be continued for at least 14 days after discharge or until COVID-19-related morbidity (including immobility, dehydration and/or shortness of breath) has resolved (consensus recommendation) .

Dosing of LMWH is dependent on pre-pregnancy body weight and current renal function.

For women with early pregnancy body weight outside of 50-90 kg, consider adjusted LMWH dosing.

There is limited evidence to guide the most appropriate dose in obese patients but standard dosing may be inadequate.

For pregnant women with severe or critical COVID-19, or where there are additional risk factors for VTE, consider using increased prophylactic dosing of anticoagulants, preferably LMWH (eg, enoxaparin 40 mg twice daily or dalteparin 5000 IU twice daily) unless there is a contraindication, such as risk for major bleeding or platelet count < 30 × 10⁹/L. Prophylactic anticoagulants should be continued for at least four weeks after discharge or until COVID-19-related morbidity (including immobility, dehydration and/or shortness of breath) has resolved (consensus recommendation) .

Dosing is dependent on pre-pregnancy body weight and current renal function. For women with early pregnancy body weight outside of 50-90 kg, consider adjusted LMWH dosing.

There is limited evidence to guide the most appropriate dose in obese patients but standard dosing may be inadequate.

Clinicians should refer to their local or jurisdictional guidance on additional VTE risk factors.

In some situations, continuation of LMWH throughout the rest of pregnancy and postpartum may be required. Involvement of specialist obstetricians, obstetric medicine physicians, haematologists or other physicians with expertise in VTE in pregnant women would be warranted.

For pregnant or postpartum women who are self-isolating at home with mild COVID-19 and where additional risk factors for VTE are present, consider using prophylactic doses of anticoagulants, preferably LMWH (eg, enoxaparin 40 mg once daily or dalteparin 5000 IU once daily) unless there is a contraindication, such as risk for major bleeding or imminent birth. Prophylactic anticoagulants should be continued for at least 14 days or until COVID-19-related morbidity (including immobility, dehydration and/or shortness of breath) has resolved. For pregnant or postpartum women who are self-isolating at home with mild COVID-19 and who have no additional risk factors for VTE, routine pharmacological prophylaxis is not recommended (consensus recommendation)

Dosing of LMWH is dependent on pre-pregnancy body weight and current renal function. For women with early pregnancy body weight outside of 50–90 kg, consider adjusted LMWH dosing.

There is limited evidence to guide the most appropriate dose in obese patients but standard dosing may be inadequate.

Clinicians should refer to their local or jurisdictional guidance on additional VTE risk factors.

For postpartum women who have had COVID-19 during pregnancy, consider using at least 14 days of prophylactic dosing of anticoagulants, preferably LMWH (eg, enoxaparin 40 mg once daily or dalteparin 5000 IU once daily) unless there is a contraindication, such as risk for major bleeding. Increased duration of six weeks should be considered if severe or critical COVID-19 and/or additional risk factors for VTE are present (consensus recommendation)

Dosing of LMWH is dependent on pre-pregnancy body weight and current renal function. For women with early pregnancy body weight outside of 50–90 kg, consider adjusted LMWH dosing.

There is limited evidence to guide the most appropriate dose in obese patients but standard dosing may be inadequate.

Clinicians should refer to their local or jurisdictional guidance on additional VTE risk factors.

Recommendations were current at the time of writing. Please visit The National COVID-19 Clinical Evidence Taskforce (<https://covid19evidence.net.au/>) for the latest updates to the recommendations.

John Wiley & Sons, Ltd Prone positioning and extracorporeal membrane oxygenation (ECMO) Internationally, there are few (and somewhat conflicting) guidelines on the use of prone positioning or ECMO in pregnant and postpartum women with COVID-19. 4 , 71 , 72 , 73 , 74 There is currently no primary evidence on their benefits and harms in these women, although it is known to be effective in some groups of adult patients with critical disease. 75 , 76 Recent reports describe their use in small numbers of pregnant women with critical COVID-19 disease. 77 , 78 , 79 , 80 The Taskforce recommends that prone positioning be considered in women receiving any form of supplemental oxygen therapy or mechanically ventilated women, although care should be taken to support the gravid uterus and reduce aorta-caval compression (Table 5). Referral to an ECMO centre for veno-venous ECMO can be considered for mechanically ventilated pregnant women, although this decision needs to consider gestational age, fetal viability, fetal well-being, and the risks and benefits to woman and baby. The Taskforce has issued separate recommendations on other respiratory support interventions that apply to adults (whether pregnant or non-pregnant), including the use of high-flow nasal oxygen therapy, noninvasive ventilation, respiratory management of the deteriorating patient, videolaryngoscopy, neuromuscular blockers, positive end-expiratory pressure and recruitment manoeuvres. These can be found on the Taskforce website. 21

Respiratory support for pregnant and postpartum women with severe or critical COVID-19 †

For mechanically ventilated pregnant women with COVID-19 and hypoxaemia despite optimising ventilation, consider prone positioning for more than 12 h a day (consensus recommendation) .

For pregnant and postpartum women with COVID-19 and respiratory symptoms who are receiving any form of supplemental oxygen therapy and have not yet been intubated, consider prone positioning. When positioning a pregnant woman in prone, care should be taken to support the gravid uterus to reduce aorta-caval compression. Women who are deteriorating should be considered for early endotracheal intubation and invasive mechanical ventilation. Birth of the baby should be considered when it may enhance maternal resuscitation or be beneficial to the fetus (consensus recommendation) .

Current reports suggest prone ventilation in adult patients is effective in improving hypoxia associated with COVID-19. This should be done in the context of a hospital guideline that includes suitable personal protective equipment for staff, and that minimises the risk of adverse events, e.g. accidental extubation.

Proning of a pregnant woman should avoid abdominal compression and ensure a woman's hips and chest are supported. In the absence of specialised equipment, proning can be performed using pillows and blankets.

Proning can be challenging in late gestation and delivery of the baby may be warranted.

Consider referral to an ECMO centre for venovenous ECMO in mechanically ventilated pregnant women with COVID-19 and refractory respiratory failure (despite optimising ventilation, including proning). Delivery of the baby prior to ECMO to enhance maternal resuscitation should be considered on a case-by-case basis (consensus recommendation) .

Due to the resource-intensive nature of ECMO and the need for experienced centres, healthcare workers and infrastructure, ECMO should only be considered in selected pregnant women with COVID-19 and severe ARDS.

The decision on whether to use ECMO should be taken in consultation with the woman's family, as well as obstetric and intensive care specialists. Key considerations include gestational age, fetal viability, fetal wellbeing and the risks and benefits to mother and baby.

Early referral to an ECMO centre is preferred.

As pregnant and postpartum women may have haemostatic alterations, anticoagulation regimens may need to be modified appropriately.

Recommendations were current at the time of writing. Please visit The National COVID-19 Clinical Evidence Taskforce (<https://covid19evidence.net.au/>) for the latest updates to the recommendations.

John Wiley & Sons, Ltd Discussion The treatment of COVID-19 is a rapidly expanding area of research, with an unprecedented global effort underway. However, with more than 53 000 COVID-19 articles in PubMed, as well as 2100 systematic review protocols and 1600 trial protocols registered online in less than eight months, it is increasingly difficult for clinicians to remain up-to-date on important developments. 15 , 21 , 81 There is also a growing recognition that a portion of the COVID-19 literature is of poor quality, further complicating its interpretation and application in clinical care. 82 The living guideline approach ensures that new, important evidence is rapidly translated into clinical recommendations, without compromising quality or rigour of guideline development. While living guidelines methods have been successfully applied to Australian guidelines on stroke and diabetes, 18 , 83 the approach of weekly updating used by this Taskforce is the most frequent of which we are aware. While resource-intensive, the collaborative

effort of over 220 clinical experts nationwide, as well as support from commonwealth, state and territory governments and other funders, has shown this model to be highly feasible in the context of a global public health emergency.

A key finding from these recommendations is the lack of direct evidence specific to pregnant and breastfeeding women with COVID-19. Historically, women (and particularly pregnant and breastfeeding women) have been excluded from adult clinical trials, despite repeated calls from professional associations, regulatory agencies and researchers for their inclusion.^{84, 85, 86} This trend has continued in the era of COVID-19, with pregnant and breastfeeding women excluded from large multicentre trials of COVID-19 therapeutics.⁸⁷ Pastick et al. reported that in 1282 ongoing COVID-19 therapeutic clinical trials enrolling people of reproductive age, only 48 explicitly confirmed that pregnant and breastfeeding women could be enrolled.⁸⁸ Exclusion of pregnant women from COVID-19 trials creates uncertainty regarding safety and effectiveness of promising management strategies, and ultimately leads to reduced access to important treatments. Consequently, the Taskforce recommendations have considered whether available evidence from trials of non-pregnant adults can be generalised to pregnant and breastfeeding women.

These recommendations are intended to be used in conjunction with existing principles and guidelines on routine maternity care. However, living recommendations can require more dynamic approaches to implementation. Clinicians and other stakeholders need to be continually updated regarding new or revised recommendations, and clinical tools (such as flowcharts and algorithms) need to be regularly revised to align with updated recommendations. Since their launch, the Taskforce website has been visited over 180 000 times. We encourage hospital and clinical networks to sign up to the free weekly email updates, and update their jurisdictional guidance or local clinical protocols as necessary. While the scope of the living guidelines has thus far focused on clinical care of people with COVID-19, on 9 September 2020 the Commonwealth Government announced a new partnership between the Taskforce and the Infection Control Expert Group (ICEG) to apply the living guidelines model to selected infection prevention and control issues in Australian healthcare settings.⁸⁹

Living guidelines are an innovative model that can accelerate the translation of new evidence into clinical practice. The National COVID-19 Clinical Evidence Taskforce has published 18 pregnancy and perinatal living recommendations, which will be updated as necessary in response to new, important evidence. We encourage clinicians in Australia's maternity services to freely subscribe to Taskforce updates (covid19evidence.net.au) to ensure they remain abreast of guideline updates.

All recommendations and evidence profiles are publicly available at <https://covid19evidence.net.au/>

We would like to acknowledge all individual members of the COVID-19 Taskforce. Steering Committee: Sharon McGowan (Chair), Nicola Ballenden, Terri Lee Barrett, Vanessa Beavis, James Beckford Saunders, Tanya Buchanan, Marina Buchanan Grey, Dawn Casey, Marita Cowie, Joseph Doyle, Mark Frydenberg, Danijela Gnjidic, Sally Green, Rohan Greenland, Ken Griffin, Stephan Groombridge, Louise Hardy, Alison Hodak, Anthony Holley, Vase Jovanovska, Sabina Knight, Kristin Michaels, Peter Morley, Julia Morphet, Suzi Nou, Phillip Russo, Megan Sarson, Alan Young. Executive Team: Julian Elliott, Rhiannon Tate, Britta Tendal, Sarah Norris, Bronwyn Morris Donovan, Joshua Vogel, Sharon Gurry, Eloise Hudson, Shauna Hurley, Declan Primmer, Samantha Timms, Susan Whicker. National Guidelines Leadership Group: Julian Elliott (Co-Chair), Sutapa Mukherjee (Co-Chair), Joshua Vogel (Deputy Chair), Jason Agostino, Karen Booth, Lucy Burr, Lyn Byers, Peter Cameron, Megan Cooper, Allen Cheng, Peter Fowler, Mark Frydenberg, Alan Glanville, Caroline Homer, Karin Leder, Steve McGloughlin, Brendan McMullan, Ewen McPhee, Brett Mitchell, Mark Morgan, Paul Myles, Chris O'Donnell, Michael Parr, Jane Phillips, Rebecca Randall, Wayne Varndell, Ian Whyte, Leeroy William. Consumer Panel: Rebecca Randall (Chair), Richard Brightwell, Lynda Condon, Amrita Deshpande, Adam Ehm, Monica Ferrie, Joanne Muller, Lara Pullin, Elizabeth Robinson, Adele Witt. Primary and Chronic Care Panel: Sarah Larkins (Co-Chair), Mark Morgan (Co-Chair), Georgina Taylor (Deputy Chair), Jason Agostino, Paul Burgess, Penny Burns, Lyn Byers, Kirsty Douglas, Ben Ewald, Dan Ewald, Dianna Fornasier,

Sabina Knight, Carmel Nelson, Louis Peachey, David Peiris, Mieke van Driel, Lucie Walters, Ineke Weaver. Hospital and Acute Care Panel: Lucy Burr (Chair), Simon Hendel (Deputy Co-Chair), Kiran Shekar (Deputy Co-Chair), Bronwyn Avar, Kelly Cairns, Allan Glanville, Nicky Gilroy, Paul Myles, Robert O'Sullivan, Owen Robinson, Chantal Sharland, Sally McCarthy, Peter Wark. Critical Care Panel: Steve McGoughlin (Co-Chair), Priya Nair (Co-Chair), Carol Hodgson, (Deputy Chair), Melissa Ankravs, Craig French, Kim Hansen, Sue Huckson, Jon Iredell, Carrie Janerka, Rose Jaspers, Ed Litton, Stephen Macdonald, Sandra Peake, Ian Seppelt. Pregnancy and Perinatal Care Panel: Caroline Homer (Co-Chair), Vijay Roach (Co-Chair), Michelle Giles (Deputy Co-Chair), Clare Whitehead (Deputy Co-Chair), Wendy Burton, Teena Downton, Glenda Gleeson, Adrienne Gordon, Jenny Hunt, Jackie Kitschke, Nolan McDonnell, Philippa Middleton, Jeremy Oats. Paediatric and Adolescent Care Panel: Asha Bowen (Co-Chair), Brendan McMullan (Co-Chair), David Tingay (Deputy Co-Chair), Nan Vasilunas (Deputy Co-Chair), Lorraine Anderson, James Best, Penny Burns, Simon Craig, Simon Erickson, Nick Fancourt, Zoy Goff, Vimbai Kapuya, Catherine Keyte, Lorelle Malyon, Danielle Wurzel. Palliative and Aged Care Panel: Meera Agar (Co-Chair), Richard Lindley (Co-Chair), Natasha Smallwood (Deputy Chair), Mandy Callary, Michael Chapman, Philip Good, Peter Jenkin, Deidre Morgan, Vasi Naganathan, Velandai Srikanth, Penny Tuffin, Elizabeth Whiting, Leeroy William, Patsy Yates. Disease-Modifying Treatment and Chemoprophylaxis Panel: Bridget Barber, Jane Davies, Josh Davis, Dan Ewald, Michelle Giles, Amanda Gwee, Karin Leder, Gail Matthews, James McMahon, Trisha Peel, Chris Raftery, Megan Rees, Jason Roberts, Ian Seppelt, Tom Snelling, Brad Wibrow. Expert Advisory Group: Ross Baker, Jennifer Curnow, Briony Cutts, Anoop Enjeti, Andrew Forbes, Prahlad Ho, Adam Holyoak, Helen Liley, James McFadyen, Zoe McQuilten, Eileen Merriman, Helen Savoia, Chee Wee Tan, Huyen Tran, Chris Ward, Katrina Williams. Cardiac Arrest Working Group: Neil Ballard, Samantha Bendall, Neel Bhanderi, Lyn Byers, Simon Craig, Dan Ellis, Dan Ewald, Craig Fairley, Brett Hoggard, Minh Le Cong, Peter Morley, Priya Nair, Andrew Pearce. Evidence Team: Britta Tendal, Steve McDonald, Tari Turner, Joshua Vogel, David Fraile Navarro, Heath White, Samantha Chakraborty, Saskia Cheyne, Henriette Callesen, Sue Campbell, Jenny Ring, Agnes Wilson, Tanya Millard, Melissa Murano. Observational Data Working Group: David Henry (Co-Chair), Sallie Pearson (Co-Chair), Douglas Boyle, Kendal Chidwick, Wendy Chapman, Craig French, Chris Pearce, Tom Snelling. Independent Conflicts of Interest Committee: Lisa Bero (Chair), Quinn Grundy, Joel Lexchin, Barbara Mintzes.

We would like to acknowledge the Member Organisations: Australian Living Evidence Consortium (Convenor), Cochrane Australia (Secretariat), Australasian Association of Academic Primary Care, Australian Association of Gerontology, Australasia College for Emergency Medicine, Australasian College for Infection Prevention and Control, Australasian Society for Infectious Diseases, Australasian College of Paramedicine, Australasian Society of Clinical and Experimental Pharmacologists and Toxicologists, Australian and New Zealand College of Anaesthetists, Australian and New Zealand Intensive Care Society, Australian & New Zealand Society for Geriatric Medicine, Australian College of Critical Care Nurses, Australian College of Midwives, Australian College of Nursing, Australian College of Rural and Remote Medicine, Australian COVID-19 Palliative Care Working Group, Australian Primary Health Care Nurses Association, Australian Resuscitation Council, Australasian Sleep Association, Australian Society of Anaesthetists, Australasian College of Paramedicine, College of Emergency Nurses Australasia, CRANaplus, National Aboriginal Community Controlled Health Organisation, Royal Australasian College of Physicians, Royal Australian College of Surgeons, Royal Australian and New Zealand College of Obstetricians and Gynaecologists, Royal Australian College of General Practitioners, Society of Hospital Pharmacists of Australia, Thoracic Society of Australia and New Zealand, Thrombosis and Haemostasis Society of Australia and New Zealand.

We would like to acknowledge our Partners: Australian Commission on Safety and Quality in Health Care, Cochrane, Consumers Health Forum of Australia, Covidence, Hereco, MAGIC, NPS MedicineWise, School of Public Health and Preventive Medicine, Monash University.

We would like to acknowledge our Funders: Australian Government Department of Health, Victorian Government Department of Health and Human Services, The Ian Potter Foundation, Walter

Cottman Endowment Fund, managed by Equity Trustees, Lord Mayors' Charitable Foundation.