

Clinical practice guidelines for visualized percutaneous breast tissue clips: Chinese Society of Breast Surgery (CSBrS) practice guideline 2021.

Title: Clinical practice guidelines for visualized percutaneous breast tissue clips: Chinese Society of Breast Surgery (CSBrS) practice guideline 2021.

Publication year: 2021

Journal: Chinese medical journal

Authors: Department of Breast Surgery, Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang 310009, China.

Society: European Respiratory Society

Origin country: International (Europe)

Target countries: Europe

DOI: 10.1097/CM9.0000000000001585

PMID: 34091526

Source: <https://pubmed.ncbi.nlm.nih.gov/34091526/>

Chin Med J (Engl) Chin Med J (Engl) 2337 chinesemj 7513795 CM9 Chinese Medical Journal 0366-6999 2542-5641 Wolters Kluwer Health PMC8367061 PMC8367061.1 8367061 8367061 34091526 10.1097/CM9.0000000000001585 CMJ-2021-190 1 Consensus Statements Clinical practice guidelines for visualized percutaneous breast tissue clips: Chinese Society of Breast Surgery (CSBrS) practice guideline 2021 Chen Wu-Zhen Jiang Jing-Xin Sun Shan-Shan Yu Xiu-Yan Huang Jian Chinese Society of Breast Surgery Yin Yan-Jie Hao Xiu-Yuan Department of Breast Surgery, Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang 310009, China. Correspondence to: Dr. Jian Huang, Department of Breast Surgery, Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang 310009, China E-Mail: drhuangjian@zju.edu.cn 5 8 2021 04 6 2021 134 15 388084 1768 1770 24 1 2021 05 08 2021 18 08 2021 05 05 2024 Copyright © 2021 The Chinese Medical Association, produced by Wolters Kluwer, Inc. under the CC-BY-NC-ND license. 2021 <https://creativecommons.org/licenses/by-nc-nd/4.0/> This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal. <http://creativecommons.org/licenses/by-nc-nd/4.0> pmc-status-qastatus 0 pmc-status-live yes pmc-status-embargo no pmc-status-released yes pmc-prop-open-access yes pmc-prop-olf no pmc-prop-manuscript no pmc-prop-legally-suppressed no pmc-prop-has-pdf yes pmc-prop-has-supplement yes pmc-prop-pdf-only no pmc-prop-suppress-copyright no pmc-prop-is-real-version no pmc-prop-is-scanned-article no pmc-prop-preprint no pmc-prop-in-epmc yes pmc-license-ref CC BY-NC-ND OPEN-ACCESS TRUE Breast tissue clip is a kind of tissue marker visible under ultrasound (US), mammogram (MMG), and magnetic resonance imaging (MRI) and could be used to precisely mark the breast lesions and axillary lymph nodes (LN) under US or MMG guidance. [1] In order to standardize the clinical application of tissue clip, the Chinese Society of Breast Surgery (CSBrS) identified key clinical issues and developed this clinical practice guideline after evaluating the evidence with reference to the GRADE (ie , Grading of Recommendations Assessment, Development, and Evaluation) system and the accessibility in clinical practice.

There were 78 voting committee members for the guideline: 65 breast surgeons (83.3%), four medical oncologists (5.1%), three radiologists (3.9%), two pathologists (2.6%), one radiation therapist (1.3%), and two epidemiologists (2.6%).

Clinicians specializing in breast diseases in China.

Recommendation 1: Indications of tissue clips

Recommendation 2: Contraindications of clip placement

Recommendation 3: Clinical issues for positioning tissue clip

With the widespread screening for breast cancer, detected nonpalpable breast lesions have been increased. [16] Clip placement in nonpalpable suspicious breast lesions could provide help in follow-up and surgical biopsy. [3] The expert panel recommended the clip placement in nonpalpable breast suspicious lesions as one indication.

Accurate localization of nonpalpable breast cancer is a prerequisite for breast-conserving surgery, whereas clip placement and localization could increase negative margin rate to 90%, [11] and reduce the secondary resection. [4 , 5] The expert panel recommended the clip placement and perioperative poisoning in nonpalpable breast cancer intended for breast-conserving surgery as one indication.

In breast cancer neoadjuvant therapy, the clinical complete remission rate could reach 20% to 57%. [6 , 9] Therefore, residual lesion re-localization after neoadjuvant therapy is important. Retrospective studies showed that clip placement at primary breast cancer before neoadjuvant therapy helped localize tumor intraoperatively and improve the negative surgical margin rate. [6] Besides, the 5-year local recurrence rate was lower in the clipped group than in the control. [6] The expert panel recommends the clip placement for primary breast cancer in neoadjuvant patients, especially planned for breast-conserving surgery.

Functional axillary LN dissection has received increasing attention recently. The ACOSOG Z1071, MARI, and ILINA trials have shown the feasibility of the targeted axillary dissection (TAD) technique, suggesting that the removal of sentinel and clipped LNs could reflect the axillary LNs status (false negative rate/FNR 2%–7%) and reduce the incidence of postoperative upper limb lymphedema. [7 – 9 , 17] The expert panel recommends clip placement to label the pathologically confirmed metastatic axillary LN (pN1) before neoadjuvant therapy, providing an opportunity for TAD.

The contraindications to US or MMG-guided clip placement for breast lesion and axillary LN should be based on the preoperative evaluation principles of breast lesion and regional LN biopsy, with reference to published exclusion criteria in clinical trials. [7 , 8] The expert panel recommends the contraindications for clip placement at Recommendation 2.

US-guided clip placement has been reported to be more accurate than X-ray-guided placement. [3 , 12] The expert panel preferred US-guided clip placement, with X-ray-guided placement as an optional method. Regarding the number of clips to be placed, which should balance localization accuracy and medical cost, [10 , 11] the expert panel recommended one clip for each primary breast lesion or metastatic axillary LN.

Due to the small size of clips, intraoperative position failure and loss of clips have been reported in 5% to 20% of cases. [7] Considering the clinical operability and accessibility in China, the expert panel recommended the use of guide wires or dyes for preoperative assisted clip localization, and isotopic tracers could be used when available. Intraoperative resected specimens need to be re-examined, and the expert panel recommends the use of radiographic imaging for intraoperative clip confirmation.

Regarding the timing of clip placement, delayed clip placement in primary breast lesions after 2-cycle of effective neoadjuvant therapy may achieve some health-economics advantages, [18] but further confirmation is needed from prospective studies. For metastatic axillary LN, published studies indicated that clip placement needed before neoadjuvant therapy. [7 – 9] The expert panel

recommends clip placement in pathologically confirmed primary breast cancer and metastatic axillary LN before neoadjuvant therapy.

Appendix 1: Technical guidelines for visualized percutaneous breast tissue clip placement.

The expert committee for these guidelines declares no conflict of interest.

These guidelines are a reference for breast disease specialists in clinical practice. However, the guidelines are not to be used as the basis for medical evaluation, and do not play an arbitrating role in the handling of any medical disputes. The guidelines are not a reference for patients or nonbreast specialists. The Chinese Society of Breast Surgery assumes no responsibility for results involving the inappropriate application of these guidelines and reserves the right to interpret and revise the guidelines.

List of Committee members (In alphabetical order by surname): Zhong-Wei Cao, Yuan-Jia Cheng, De-Tian Chen, Wu-Zhen Chen, Xue-Ning Duan, Zhi-Min Fan, Pei-Fen Fu, Jian Huang, Jun Jiang, Hong-Chuan Jiang, Feng Jin, Jing-Xin Jiang, Hua Kang, Rui Ling, Jin-Ping Liu, Ke Liu, Qian Liu, Yin-Hua Liu, Yun-Jiang Liu, Zhen-Zhen Liu, Yong-Hui Luo, Rong Ma, Da-Hua Mao, Jiang-Hua Ou, Xiang Qu, Guo-Sheng Ren, Ai-Lin Song, Er-Wei Song, Shan-Shan Sun, Li-Li Tang, Xing-Song Tian, Chuan Wang, Fei Wang, Jian-Dong Wang, Shu Wang, Shui Wang, Xiang Wang, Jiong Wu, Ke-Jin Wu, Ling Xin, Zhi-Gang Yu, Xiu-Yan Yu, Jian-Guo Zhang, Jin Zhang, Jing-Hua Zhang, Yi Zhao, Zuo-Wei Zhao, Wei Zhu, Ang Zheng, Qiang Zou.

How to cite this article: Chen WZ, Jiang JX, Sun SS, Yu XY, Huang J, Chinese Society of Breast Surgery. Clinical practice guidelines for visualized percutaneous breast tissue clips: Chinese Society of Breast Surgery (CSBrS) practice guideline 2021. Chin Med J 2021;134:1768–1770. doi: 10.1097/CM9.0000000000001585

Supplemental digital content is available for this article.