

Management of Functional Seizures Practice Guideline Executive Summary

Report of the AAN Guidelines Subcommittee

Benjamin Tolchin,¹ Laura H. Goldstein,² Markus Reuber,³ Jon Stone,⁴ David L. Perez,⁵ W. Curt LaFrance, Jr.,⁶ Aaron D. Fobian,⁷ James Dorman,⁸ Le H. Hua,⁹ Z Paige L'Erario,¹⁰ Sara J. Swanson,¹¹ Peter Gilli,¹² Bridget Mildon,¹³ Courtney Takahashi,¹⁴ Kylie A. Botchway-Doe,¹⁵ Katie Puskala Hamel,¹⁵ Heather M. Silsbee,¹⁵ and Maryam Oskoui¹⁶

Neurology® 2026;106:e214466. doi:10.1212/WNL.00000000000214466

Correspondence

American Academy of
Neurology
guidelines@aan.com

Abstract

This guideline provides evidence-based recommendations for clinicians, patients, and other stakeholders on the management of functional seizures. Following a National Academy of Medicine–compliant process, a multidisciplinary panel conducted a systematic review and integrated the findings with the authors' clinical experience to develop recommendations. A systematic review of the available evidence from first published articles to February 25, 2025, identified 12 Class II–III studies. The review found that psychological interventions are possibly effective in increasing the likelihood of achieving freedom from functional seizures, decreasing the frequency of functional seizures, decreasing anxiety, and improving health-related quality of life and psychosocial functioning for individuals with functional seizures. Key recommendations state that, when evaluating patients with seizure-like episodes, clinicians should seek historical and semiological information (including smartphone videos) from both patients and witnesses and may obtain video-EEG of all typical seizure-like episodes where feasible. Clinicians should evaluate patients diagnosed with functional seizures for co-occurring psychiatric disorders and co-occurring epilepsy. Clinicians should adhere to universal standards of care for patients, including speaking respectfully, refraining from unnecessary harm, and avoiding stigmatizing behavior. Clinicians should provide a specific diagnostic label and rationale for the diagnosis, should engage in shared decision making regarding the treatment plan, and should provide continuity of care to individuals diagnosed with functional seizures. When psychological interventions for functional seizures are indicated, clinicians should counsel patients regarding the potential benefits and risks of these interventions and should refer interested and appropriate patients to these interventions for the treatment of functional seizures. Clinicians should involve family, caregivers, or others in the social support network in the psychological treatment of individuals with functional seizures. Clinicians should not prescribe benzodiazepines or antiseizure medications for patients with functional seizures without co-occurring epilepsy or another indication for these medications and should counsel patients regarding the potential risks and lack of evidence of benefit for using these medications for functional seizures. Clinicians should taper off antiseizure medications for patients with functional seizures and without another indication for these medications. The guideline also identifies gaps in the available evidence and outlines potentially clinically impactful avenues for future research.

Introduction

Functional seizures are transient episodes of altered consciousness or involuntary movements resembling epileptic seizures or syncope but conceptualized to be driven by episodic dissociation or other cognitive-affective mechanisms.¹ Functional seizures have alternatively or previously been

MORE ONLINE

Supplementary Material



¹Department of Neurology, Yale School of Medicine, New Haven, CT; ²Department of Psychology, Institute of Psychiatry, Psychology and Neuroscience, King's College London, United Kingdom; ³Division of Neuroscience, University of Sheffield, United Kingdom; ⁴Centre for Clinical Brain Sciences, University of Edinburgh, Scotland; ⁵Departments of Neurology and Psychiatry, Massachusetts General Hospital, Mass General Brigham, Harvard Medical School, Boston; ⁶Departments of Psychiatry and Neurology, Rhode Island Hospital and Alpert Medical School, Brown University, Providence; ⁷Department of Psychiatry and Behavioral Neurobiology, Heersink School of Medicine, University of Alabama at Birmingham; ⁸Department of Neurology, Rush Medical College and Cook County Health, Chicago, IL; ⁹Department of Neurology, Lou Ruvo Center for Brain Health, Cleveland Clinic, Las Vegas, NV; ¹⁰Department of Neurology, Thomas Jefferson University, Philadelphia, PA; ¹¹Department of Neurology, Medical College of Wisconsin, Milwaukee; ¹²FND Portal, Canada; ¹³FND Hope, Salmon, ID; ¹⁴Department of Neurology, Boston University School of Medicine, MA; ¹⁵American Academy of Neurology, Minneapolis, MN; and ¹⁶Departments of Pediatrics and Neurology and Neurosurgery, McGill University, Montreal, Quebec, Canada.

Approved by the Guidelines Subcommittee on June 13, 2025; by the AAN Quality Committee on July 20, 2025; and by the AAN Institute Board of Directors on October 27, 2025.

Glossary

AAN = American Academy of Neurology; CBT = cognitive behavioral therapy; COI = conflict of interest; NBT = neurobehavioral therapy; PTSD = post-traumatic stress disorder; ReACT = Retraining and Control Therapy; RMD = raw mean difference; RR = risk ratio; SMC = standardized medical care; SMD = standardized mean difference; VEEG = video-EEG; WSAS = Work and Social Adjustment Scale.

labeled as dissociative seizures, psychogenic nonepileptic seizures, psychogenic nonepileptic attacks, or conversion disorder with attacks or seizures.² Functional seizure disorders are often characterized by high biopsychosocial complexity, with a range of implicated predisposing vulnerabilities, acute precipitants, and perpetuating factors,³⁻⁵ including concurrent psychiatric conditions and adverse life experiences. Adverse life events can include past or present physical injury; illness; neglect; or physical, emotional, or sexual abuse.^{4,6,7} However, co-occurring psychiatric disorders and adverse life experiences are not universally present, are not required diagnostic criteria, and may be less frequent in pediatric populations.⁸ Functional seizures are a common and often disabling subtype of functional neurologic symptom disorder, as classified in the *Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition Text Revision*.⁹ Functional seizures, and functional neurologic disorder more generally, do not involve the intentional simulation of signs or symptoms and should not be conflated with malingering or factitious disorder, which are much less common.¹⁰

There is a growing body of evidence demonstrating that psychological treatments may be effective in treating

functional seizures,¹¹ improving quality of life, and reducing health care costs.¹² Nevertheless, most people with functional seizures do not receive targeted and evidence-based treatment, contributing to ongoing functional seizures and disability.^{13,14} The limited utilization of evidence-based treatment for functional seizures is due to several factors, including stigmatization of individuals with functional seizures, limited communication and collaboration between neurologists and mental health clinicians, limited access to psychotherapy, poor reimbursements for counseling services, and a lack of familiarity with the disorder among many clinicians and the general public. This practice guideline provides evidence-based recommendations on the use of psychological and pharmacologic interventions in the management of functional seizures to reduce treatment gaps and stigma and improve patient outcomes. The guideline also identifies areas where evidence is unavailable or inadequate to inform future research. The guideline focuses on the following clinical questions:

1. In individuals with functional seizures, what is the efficacy of psychological interventions on the frequency and

Figure 1 Study Selection Flowchart




Table 1 Study Participant Demographics

Study and location	N	Mean age, y	% Female sex	Race and/or ethnicity	Employment status	Includes participants with co-occurring epilepsy	Psychiatric comorbidities	% Diagnosed with VEEG
Aboukasm et al., 1998, United States ¹⁷	100	43.3	90	Not available	64% unemployed	Y	Not available	100
Ataoglu et al., 2003, Turkey ¹⁸	30	23 (range 16–30)	97	Not available	Not available	N	Not available	0
Chen et al., 2014, United States ¹⁹	64	50.7	25	Not available	82.8% unemployed	N	39% PTSD	100
Fobian et al., 2020, United States ²⁰	29	15.1 (SD 2.5)	57	57.1% White, 28.6% Black, 14.3% Other	Not available	Y if free of epileptic seizures >6 mo	52% (anxiety and/or depression)	100
Goldstein et al., 2010, United Kingdom ²¹	66	37.4 (SD 12.6) in the intervention arm, 35.9 (SD 15.1) in the control arm	76	87.9% White, 6.0% Black, 4.5% Asian, 1.5% Other	55% unemployed	N	47% (any psychiatric comorbidity)	92
Goldstein et al., 2020, United Kingdom ²²	368	37.5 (SD 14.3)	72	90% White, 2% Black, 2% Asian, 7% Other	66% unemployed	Y if free of epileptic seizures >12 mo	69% (any psychiatric comorbidity)	53
Goldstein et al., 2022, United Kingdom ²³	368	37.5 (SD 14.3)	72	90% White, 2% Black, 2% Asian, 7% Other	66% unemployed	Y if free of epileptic seizures >12 mo	69% (any psychiatric comorbidity)	53
Khattak et al., 2006, Pakistan ²⁴	100	24.3 (SD 8.76)	88	Not available	Not available	N	Not available	0
LaFrance et al., 2010, United States ²⁵	38	34.4 (SD 12.6) in the control arm and 38 (SD 13.9) in the intervention arm	76.3	Not available	65.8% unemployed	Y if could be distinguished from functional seizure	61% mood disorders, 87% anxiety disorders	100
LaFrance et al., 2014, United States ²⁶	38	37.9 (SD 11.5) in the neurobehavioral psychotherapy arm, 39.1 (SD 13.2) in the neurobehavioral psychotherapy + sertraline arm, 39.7 (SD 11.7) in the sertraline arm, and 41.6 (SD 8.3) in the treatment as usual arm	81.6	Not available	68.4% unemployed	N	68% mood disorders, 79% anxiety disorders	100
Senf-Beckenbach et al., 2022, Germany ²⁷	53	34.7 (SD 17.9)	70.5	Not available	Not available	N	Not available	Not available
Tolchin et al., 2019, United States ²⁸	55	39.6 (SD 16.8) in the intervention arm and 40.7 (SD 14.3) in the control arm	82.7	66.7% White non-Hispanic, 10% Black non-Hispanic, 21.7% Hispanic	Not available	N	42% depression, 58% anxiety	100

Abbreviations: PTSD = post-traumatic stress disorder; VEEG = video-EEG.

- bothersomeness of functional seizures in comparison with standardized medical care?
- In individuals with functional seizures, what is the efficacy of psychological interventions on health-related quality
- of life, psychosocial functioning, and psychiatric comorbidities in comparison with standardized medical care?
- In individuals with functional seizures, what is the efficacy of psychopharmacologic interventions on the frequency

Table 2 Study Interventions, Comparators, and Outcomes

Study	Class	Intervention and duration	Comparator	Outcome measures	Time points
Aboukasm et al., 1998 ¹⁷	III	5 or more psychotherapy sessions with protocolized review of videotaped functional seizures	No feedback or no intervention	Functional seizure frequency, quality of life	Not specified
Ataoglu et al., 2002 ¹⁸	III	3 wk of twice daily paradoxical therapy	Diazepam 5–15 mg	Freedom from functional seizures, anxiety (HRSA)	6 wk
Chen et al., 2014 ¹⁹	III	3 monthly 1.5-hour-long group sessions of group psychoeducation	Routine follow-up in seizure clinic	Functional seizure frequency, functional seizure severity (SSQ), psychosocial functioning (WSAS), functional seizure-related emergency department visits, new functional symptoms, measure of knowledge and perception	3 mo, 6 mo
Fobian et al., 2020 ²⁰	III	8 weekly sessions of Retraining and Control Therapy	8 weekly sessions of nondirective supportive therapy	Functional seizure frequency, freedom from functional seizures, health-related quality of life (PedsQL), anxiety and depression (BASC-2), coping skills (ACOPe), somatization (CSSI)	7 d after treatment
Goldstein et al., 2010 ²¹	III	12 weekly hour-long CBT sessions (minimum of 9 sessions) + supportive sessions with a neuropsychiatrist	SMC consisting of supportive sessions (no CBT) with neuropsychiatrist	Functional seizure frequency; freedom from functional seizures, anxiety, and depression (HADS); work and social adjustment (WSAS); health service use (modified CSRI)	4 mo, 6 mo
Goldstein et al., 2020 ²²	III	CBT + SMC, CBT = 12 sessions over 4–5 mo + 1 booster session at 9 mo after randomization	SMC consisting of meeting with a neurologist, including guidance on delivering the diagnosis and information booklets + regular outpatient care from a psychiatrist with expertise in functional seizures but not including psychotherapy	Functional seizure frequency, freedom from functional seizures, seizure severity and bothersomeness (SSQ), quality of life (SF12v2, EQ-5D-5L), psychosocial functioning (WSAS), anxiety (GAD-7), depression (PHQ-9), psychological distress (CORE-10), somatic symptom burden (modified PHQ-15), caregivers' ratings of functional seizures, clinical impression of improvement (CGI) rated by patient and physician, satisfaction with treatment	12 mo after randomization
Goldstein et al., 2022 ²³	III	CBT + SMC, CBT = 12 sessions over 4–5 mo + 1 booster session at 9 mo post randomization	SMC consisting of meeting with a neurologist, including guidance on delivering the diagnosis and information booklets + regular outpatient care from a psychiatrist with expertise in functional seizures, but not including psychotherapy	Functional seizure frequency, seizure severity and bothersomeness (SSQ), quality of life (SF12v2, EQ-5D-5L), psychosocial functioning (WSAS), anxiety (GAD-7), depression (PHQ-9), psychological distress (CORE-10), somatic symptom burden (modified PHQ-15), caregivers' ratings of functional seizures, clinical impression of improvement (CGI) rated by patient and physician, satisfaction with treatment	6 mo after randomization
Khattak et al., 2006 ²⁴	III	Daily behavioral therapy sessions while admitted, followed by 4 weekly behavioral therapy sessions + routine clinical care (drug treatment, reassurance, explanation for symptoms)	Routine clinical care	Functional seizure frequency, psychosocial functioning (CGI), anxiety and depression (HADS)	4 wk after discharge
LaFrance et al., 2010 ²⁵	II	Sertraline 25–200 mg flexible dosing, adjusted in 6 biweekly sessions over 12 wk	Placebo	Functional seizure frequency, freedom from functional seizures, quality of life (QOLIE-31), anxiety (DTS), depression (BDI-II, HRSD), psychosocial functioning (LIFE-RIFT)	12 wk
LaFrance et al., 2014 ²⁶	III	1) Sertraline 25–200 mg flexible dosing; 2) 12 weekly sessions of neurobehavioral psychotherapy; 3) combined neurobehavioral psychotherapy + sertraline	Treatment as usual (follow-up with neurologist without psychotherapy or antidepressant medication)	Functional seizure frequency, freedom from functional seizures, quality of life (QOLIE-31), anxiety (BAI), depression (BDI-II)	16 wk

Continued

Table 2 Study Interventions, Comparators, and Outcomes (continued)

Study	Class	Intervention and duration	Comparator	Outcome measures	Time points
Senf-Beckenbach et al., 2022 ²⁷	III	10 weekly sessions of body-focused group psychotherapy (CORDIS)	10 weekly guided self-help group sessions	Level of dissociation (DES-20), seizure severity (LSSS), depression (PHQ-9), somatic symptoms (PHQ-15)	6.5 mo
Tolchin et al., 2019 ²⁸	III	Motivational interviewing followed by 12 weekly sessions of psychotherapy	Informational interview followed by 12 weekly sessions of psychotherapy	Adherence to psychotherapy, change in functional seizure frequency, freedom from functional seizures, quality of life (QOLIE-10), monthly emergency department visits	16 wk

Abbreviations: ACOPE = Adolescent Coping Orientation for Problem Experiences; BAI = Beck Anxiety Index; BASC-2 = Behavior Assessment System for Children, Second Edition; BDI-II = Beck Depression Inventory-II; CBT = Cognitive Behavioral Therapy; CGI = Clinical Global Impressions Scale; CORDIS = a novel integrative body-focused psychotherapy regimen; CORE-10 = Clinical Outcomes in Routine Evaluation-10; CSRI = Client Service Receipt Inventory; CSSI = Children's Somatic Symptoms Inventory; DES-20 = Dissociation Experiences Scale-20; DTS = Davidson Trauma Scale; EQ-5D-5L = EuroQoL 5-Dimension 5-Level; GAD-7 = Generalized Anxiety Disorder-7; HADS = Hospital Anxiety and Depression Scale; HRSA = Hamilton Rating Scale for Anxiety; HRSD = Hamilton Rating Scale for Depression; LIFE-RIFT = Longitudinal Interval Follow-up Evaluation-Range of Impaired Functioning Tool; LSSS = Liverpool Seizure Severity Scale; PedsQL = Pediatric Quality of Life Inventory; PHQ-15 = Patient Health Questionnaire-15; PHQ-9 = Patient Health Questionnaire-9; QOLIE-10 = Quality of Life in Epilepsy-10; QOLIE-31 = Quality of Life in Epilepsy-31; SF-12v2 = Short-Form 12-Item Survey-version 2; SMC = standard medical care; SSQ = Seizure Severity Questionnaire; WSAS = Work and Social Adjustment Scale.

and bothersomeness of functional seizures in comparison with placebo or active treatment?

4. In individuals with functional seizures, what is the efficacy of psychopharmacologic interventions on health-related quality of life, psychosocial functioning, and psychiatric comorbidities, in comparison with placebo or active treatment?

This article is a summary of the complete guideline (eAppendix 1), which includes the full systematic review, additional clinical context, and suggestions for future research.

Description of the Analytic Process

The Guidelines Subcommittee of the American Academy of Neurology (AAN) convened a multidisciplinary panel of experts to develop a practice guideline on the management of functional seizures. As required by the AAN, most (at least 51 percent) of the guideline developers (B.T., L.H.G., M.R., J.S.,

D.L.P., J.D., L.H.H., Z.P.L., S.J.S., P.G., B.M., C.T., and M.O.) and the lead developer (B.T.) are free of conflict of interest (COI) relevant to the subject matter of this guideline. Two members were determined to have relevant COIs (W.C.L. and A.D.F.), but they were judged to be not significant enough to preclude them from authorship. The authors determined to have COIs did not review or rate the evidence. Because the panel majority was free of COIs, the entire panel voted on the guideline recommendations, with those panel members free of COIs leading recommendation development. This full author panel was solely responsible for decisions concerning the design, analysis, and reporting of the systematic review and subsequent guideline. The development of this guideline follows the methodologies described in the 2017 edition of the AAN's Guideline Development Process Manual.¹⁵

The panel searched the Medline, Embase, PsychINFO, and CINAHL databases for relevant peer-reviewed articles in any language that met inclusion criteria (Figure 1). Each of the 12



Figure 2 Psychological Interventions and Probability of Achieving Freedom From Functional Seizures

Figure 3 Psychological Interventions and Functional Seizure Frequency



selected articles was rated by 2 panel members using the AAN criteria for classification of therapeutic studies.¹⁵

A modified form of the Grading of Recommendations Assessment, Development and Evaluation process was used to develop conclusions of the systematic review.^{15,16} The panel formulated practice recommendations using a modified Delphi process.

Analysis of Evidence

Table 1 provides a summary of descriptive characteristics of the 12 included studies. Table 2 provides a summary of interventions, measures, and outcomes for each of the included studies. eTable 1 provides details on the risk of bias for each of the 12 included studies.

Psychological Interventions

The psychological interventions studied included 2 functional seizure-specific cognitive behavioral therapies (CBTs),²¹⁻²³ neurobehavioral therapy (NBT),²⁶ Retraining and Control Therapy (ReACT),²⁰ behavioral therapy,²⁴ motivational interviewing,²⁹ group psychoeducation,¹⁹ body-focused group therapy,²⁷ and psychotherapy with protocolized review of videotaped functional seizures.¹⁷ Psychological interventions typically ran for 3–12 sessions. Long-term outcome data are not available beyond 1 year of follow-up.

The applicable population for each of the following conclusions is people with functional seizures.

Conclusions

Psychological interventions, compared with standard medical care (SMC), routine clinical care, nondirective supportive therapy, informational interviews, diazepam, or no therapy, possibly increase the probability of achieving freedom from functional seizures during the follow-up period. Based on 7 Class III studies,^{17,18,20-22,26-28} the pooled estimate of the risk ratio (RR) is 1.87 for seizure freedom, favoring psychological interventions (95% CI 1.36–2.56, I² = 3%) (Figure 2). Our confidence in the estimate is low, anchored by the class of studies. Exclusion of the nonrandomized study¹⁷ did not change the magnitude or significance of the RR.

Psychological interventions, compared with SMC or nondirective supportive therapy, possibly decrease functional seizure frequency at follow-up. Based on 3 Class III studies,^{20,21,23} the pooled estimate of the standardized mean difference (SMD) is -0.81, favoring psychological interventions (95% CI -1.58 to -0.05, I² = 86%) (Figure 3). Our confidence in the estimate is low, anchored by the class of studies.

There is insufficient evidence to determine whether psychological interventions, compared with SMC or nondirective supportive therapy, are more or less likely to change severity or bothersomeness of functional seizures. Based on 2 Class III studies,^{21,27} the pooled estimate of the SMD is -0.64, favoring

Figure 4 Psychological Interventions and Health-Related Quality of Life





Figure 5 Psychological Interventions and Psychosocial Functioning



psychological interventions (95% CI -1.41 to 0.13, $I^2 = 69\%$). Our confidence in the estimate is very low, anchored by the class of studies and imprecision.

Because of the heterogeneity of psychological interventions evaluated in our primary analysis, we reran our analysis including only studies evaluating functional seizure-specific CBT,^{21,22} which is the only psychological intervention for which multiple studies exist.

Functional seizure-specific CBT, compared with SMC, possibly increases the probability of achieving freedom from functional seizures. Based on 2 Class III studies,^{21,22} the pooled estimate of the RR is 1.76, favoring CBT (95% CI 1.10–2.80, $I^2 = 0\%$). Our confidence in the estimate is low, anchored by the class of studies.

There is insufficient evidence to determine whether functional seizure-specific CBT is more or less likely than SMC to result in a change in functional seizure frequency at follow-up. Based on 2 Class III studies,^{21,22} the pooled estimate of the SMD is -0.38, favoring CBT. However, the 95% CI is -0.81 to 0.04 ($I^2 = 60\%$). Our confidence in the estimate is very low, anchored by the class of studies and imprecision.

Psychological interventions, compared with SMC, nondirective supportive therapy, or informational interviews,


possibly increase health-related quality of life. Based on 3 Class III studies,^{20,22,28} the pooled estimate of the SMD is 0.37, favoring psychological interventions (95% CI 0.16–0.57, $I^2 = 0\%$) (Figure 4). Our confidence in the estimate is low, anchored by the class of studies.

Psychological interventions, compared with SMC or no intervention, possibly improve psychosocial functioning. Based on 3 Class III studies,^{19,21,22} the pooled estimate of the SMD is -0.44 on the Work and Social Adjustment Scale (WSAS), favoring psychological interventions (95% CI -0.64 to -0.24, $I^2 = 0\%$) (Figure 5). Our confidence in the estimate is low, anchored by the class of studies.

Psychological interventions, compared with routine clinical care, nondirective supportive therapy, SMC, or diazepam, possibly decrease anxiety. Based on 5 Class III studies,^{18,20–22,24} the pooled estimate of the SMD is -0.68, favoring psychological interventions (95% CI -1.30 to -0.07, $I^2 = 87\%$) (Figure 6). Our confidence in the estimate is low, anchored by the class of studies.

There is insufficient evidence to determine whether psychological interventions, compared with SMC, routine clinical care, or nondirective supportive therapy, result in a change in depression. Based on 4 Class III studies,^{20,21,23,24} the pooled estimate of the SMD is -0.43, favoring psychological

Figure 6 Psychological Interventions and Anxiety



interventions (eFigure 1). However, the 95% CI is -0.98 to 0.12 ($I^2 = 84\%$). Our confidence in the estimate is very low, anchored by the class of studies and imprecision.

Because of the heterogeneity of psychological interventions evaluated in our primary analysis, we again reran this analysis including only studies evaluating functional seizure-specific CBT,²¹⁻²³ which is the only psychological intervention for which multiple studies exist.

There is insufficient evidence to determine whether functional seizure-specific CBT, compared with SMC, changes health-related quality of life. Based on a single Class III study,²² the estimated mean difference in health-related quality of life as measured on the EuroQol 5-Dimension 5-Level Visual Analogue Scale is 6.16 , favoring CBT (95% CI 1.48 – 10.84 , $p = 0.010$). Our confidence in the estimate is very low, anchored by the number and class of studies.

Functional seizure-specific CBT, compared with SMC, possibly improves psychosocial functioning (decreases WSAS scores). Based on 2 Class III studies,^{21,22} the pooled estimate of the SMD is -0.40 , favoring CBT (95% CI -0.61 to -0.19 , $I^2 = 0\%$). Our confidence in the estimate is low, anchored by the class of studies.

Functional seizure-specific CBT, compared with SMC, possibly causes no change in anxiety. Based on 2 Class III studies,^{21,22} the pooled estimate of the SMD is -0.19 , favoring CBT. However, the 95% CI is -0.40 to 0.02 ($I^2 = 0\%$). Our confidence in the estimate is low, anchored by the class of studies and imprecision.

Pharmacologic Interventions

Regarding pharmacologic interventions, 2 studies evaluated the use of sertraline^{25,26} and 1 study examined the use of diazepam.¹⁸ The applicable population for each of the following conclusions is people with functional seizures.

Conclusions

There is insufficient evidence to determine whether sertraline, compared with placebo or routine clinical care, changes the probability of achieving freedom from functional seizures. Based on 1 Class II study²⁵ and 1 Class III study,²⁶ the pooled estimate of the RR is 2.34 , favoring sertraline (95% CI 0.34 – 16.09 , $I^2 = 49\%$). Our confidence in the estimate is very low, anchored by imprecision.

There is insufficient evidence to determine whether there is a relative change in seizure rates between those taking sertraline and those taking placebo (1 Class II study, RR 0.51 , 95% CI 0.25 – 1.05 , $p = 0.29$).²⁵ Our confidence in the estimate is very low, anchored by the number and class of studies and downgraded for imprecision.

There is insufficient evidence to determine whether diazepam, compared with paradoxical therapy, changes the probability of

achieving freedom from functional seizures (1 Class III study, $t = 2.27$, $p = 0.034$).¹⁸ Our confidence in the estimate is very low, anchored by the number and class of studies.

There is insufficient evidence to determine whether sertraline is more or less likely than placebo to improve health-related quality of life. A single Class II study²⁵ found a nonsignificant raw mean difference (RMD) of 9.80 in health-related quality of life, as measured by the Quality of Life in Epilepsy-31, favoring sertraline (95% CI -9.17 to 28.77 , $p = 0.311$). Our confidence in the estimate is very low, anchored by the number and class of studies and downgraded for imprecision.

There is insufficient evidence to determine whether sertraline, compared with placebo, changes psychosocial functioning. A single Class II study²⁵ found a nonsignificant RMD of -2.00 in psychosocial functioning, as measured on the Longitudinal Interval Follow-up Evaluation–Range of Functioning Tool, favoring sertraline (95% CI -5.22 to 1.22 , $p = 0.223$). Our confidence in the estimate is very low, anchored by the number and class of studies and downgraded for imprecision.

There is insufficient evidence to determine whether sertraline is more or less likely than placebo to improve anxiety. A single Class II study²⁵ found a nonsignificant RMD of -3.10 in anxiety, as measured on the Davidson Trauma Scale, favoring sertraline (95% CI -32.68 to 26.48 , $p = 0.837$). Our confidence in the estimate is very low, anchored by the number and class of studies and downgraded for imprecision.

There is insufficient evidence to determine whether sertraline is more or less likely than placebo to improve depression. A single Class II study²⁵ found a nonsignificant RMD of -1.70 in depression, as measured on the Hamilton Rating Scale for Depression, favoring sertraline (95% CI -8.43 to 5.03 , $p = 0.621$). Our confidence in the estimate is very low, anchored by the number and class of studies and downgraded for imprecision.

There is insufficient evidence to determine whether diazepam, compared with paradoxical therapy, changes anxiety. A single Class III study¹⁸ found an RMD of -3.73 in anxiety, as measured by the Hamilton Rating Scale for Anxiety, favoring paradoxical therapy (95% CI of -6.96 to -0.50 , $p = 0.024$). Our confidence in the estimate is very low, anchored by the number and class of studies.

Practice Recommendations

Diagnosis of Functional Seizures

Recommendation 1 Rationale

Historically, there have been significant delays in the diagnosis of functional seizures, with the diagnosis being delayed for an average of 7–8 years after symptom onset.^{30,31} Prompt and accurate diagnosis of functional seizures is essential for

effective treatment and avoidance of iatrogenesis based on misdiagnoses. A detailed clinical history and careful analysis of the seizure semiology, including characteristics positively associated with functional seizures, contribute to the accurate diagnosis of functional seizures.³²⁻³⁷ Historical and semiological information obtained from both patients and witnesses can improve the accuracy of the diagnosis.³⁸ When feasible, a brief physical examination during the ictal event can contribute to the accurate diagnosis of functional seizures and epileptic seizures.^{32,39-42} Serum prolactin, lactate, and creatine kinase may be more elevated after a bilateral tonic-clonic epileptic seizure than after other episodes involving transient loss of consciousness, but each of these laboratory tests has been associated with significant rates of false-positive and false-negative results when used to differentiate between functional seizures, epileptic seizures, and syncope.⁴³⁻⁴⁷

In combination with clinical history and semiology, video-EEG (VEEG) assessment of all typical types of episodes is the gold standard for diagnosing definite or “documented” functional seizures and differentiating them from epileptic seizures or other physiologic events.⁴⁸ VEEG can be obtained in an epilepsy monitoring unit, during 30–60-minute EEGs in the inpatient or outpatient setting, through continuous inpatient EEG monitoring, or using home-video EEG.⁴⁹ When functional seizures present with a semiology resembling syncope, additional long-term ECG monitoring and tilt-table testing may help to differentiate functional seizures from syncope.⁵⁰ VEEG is not available in all practice settings and may not be feasible for patients with infrequent seizure-like events.¹⁴ When VEEG is not feasible, a diagnosis of probable functional seizures may be based on review of history, semiology, and interictal EEG.⁴⁸ Similarly, when VEEG capture of a typical event is not feasible, a diagnosis of clinically established functional seizures can be based on history, semiology, ambulatory EEG capture of a typical event (without video), and separate video recording of a typical event.⁴⁸ Videos of seizure-like episodes captured on smartphones and reviewed by a neurologist can facilitate the evaluation of seizure semiology and allow accurate diagnosis in most of the cases.⁵¹ Individuals with functional seizures frequently experience multiple functional neurologic symptoms.^{52,53} Clinicians with greater experience in the diagnosis of seizure disorders can more accurately diagnose functional seizures.⁵⁴⁻⁵⁷

Recommendation 1 Statements

1A. Clinicians *should* include functional seizures in the differential diagnosis and in the initial workup of patients presenting with seizure-like or syncope-like episodes to make a prompt and accurate diagnosis. (Level B)

1B. When evaluating patients with seizure-like or syncope-like episodes, clinicians *should* seek historical and semiological information from both patients and witnesses when available to support a prompt and accurate diagnosis. (Level B)

1C. When acutely evaluating patients with prolonged seizure-like episodes in emergency settings, clinicians *should*

perform a brief ictal physical examination to support a prompt and accurate diagnosis. (Level B)

1D. When diagnostic ambiguity exists between epileptic and functional seizures after review of historical and semiological features including available video and EEG data, clinicians *may* obtain VEEG of typical seizure-like episodes where feasible to confirm the diagnosis with the greatest possible level of clinical certainty. (Level C)

1E. When diagnostic ambiguity exists between syncope and functional seizures after review of historical and semiological features, clinicians *should* evaluate blood pressure and cardiac rhythm during episodes using ECG monitoring and/or tilt-table testing where available to confirm the diagnosis with the greatest possible level of clinical certainty. (Level B)

1F. Where VEEG, ECG, and/or tilt-table capture of typical seizure-like or syncope-like episodes are not feasible, clinicians *should* use ambulatory EEG, interictal EEG, interictal ECG if available, smartphone video of typical seizure-like episodes, and historical and semiological features to make a diagnosis with the greatest possible level of clinical certainty. (Level B)

1G. When diagnosing a patient with functional seizures, clinicians *should* screen and evaluate for other functional neurologic symptoms, to support prompt treatment. (Level B)

1H. When diagnosis or management of functional or epileptic seizures is beyond the clinician’s scope of practice or expertise, they *should* refer patients with seizures to an appropriate specialist to support a prompt and accurate diagnosis. (Level B)

Assessment of Psychiatric Comorbidities and Epilepsy

Recommendation 2 Rationale

A lifetime history of psychiatric disorders, including mood disorders, anxiety disorders, and posttraumatic stress disorder (PTSD), co-occurs at a high frequency with functional seizures.^{29,58} Substance use disorders can also co-occur at elevated rates among people with functional seizures.^{59,60,e1} Abuse, neglect, and other adverse life experiences are 3–5 times more common among individuals with functional neurologic disorders than in the general population and twice as common as among individuals with other psychiatric disorders.^{4,e2} Co-occurring psychiatric disorders can make it more difficult for some individuals to engage in psychotherapeutic treatment, with a greater number and severity of psychiatric symptoms associated with higher rates of treatment nonadherence.^{e3,e4} Adherence to psychotherapeutic treatment is associated with better outcomes for functional seizure frequency and quality of life.^{29,e4} Accordingly, treatment of co-occurring psychiatric disorders may also facilitate psychotherapeutic treatment of functional seizures.

Lifestyle modifications such as diet, exercise, and meditation can improve quality of life in individuals with common co-

occurring psychiatric disorders such as mood disorders, anxiety disorders, and PTSD.^{e5-e7} Appropriate psychotherapeutic treatments can improve quality of life and psychosocial functioning among individuals with co-occurring psychiatric disorders, including mood, anxiety, and personality disorders.^{e8-e10} Psychopharmaceuticals such as selective serotonin reuptake inhibitors and serotonin and norepinephrine reuptake inhibitors are also effective in treating psychiatric disorders, including mood disorders, anxiety disorders, and PTSD.^{e11,e12} Therapeutic neuromodulation is also effective in treating common co-occurring psychiatric disorders, including major depression.^{e13-e16}

Recommendation 2 Statements

2A. Clinicians *should* evaluate patients diagnosed with functional seizures for co-occurring psychiatric disorders (including affective, trauma-related, personality, and substance use disorders) to facilitate treatment of both co-occurring disorders and functional seizures. (Level B)

2B. Clinicians *should* offer patients diagnosed with functional seizures and active co-occurring psychiatric disorders, who do not already receive mental health care, a referral to a mental health specialist for appropriate evidence-based treatment of their co-occurring psychiatric disorders. (Level B)

Recommendation 3 Rationale

Functional seizures may co-occur in up to 12% of people with epilepsy, and epilepsy may co-occur in up to 20% of adults with functional seizures, 30%–40% of children with functional seizures, and up to 50% of individuals with intellectual disabilities and functional seizures.^{e17,e18} Antiseizure medications improve epileptic seizure control, mortality rates, and quality of life among individuals with epilepsy.^{e19-e21} Treating co-occurring epileptic and functional seizures requires clinicians and patients to accurately identify and differentiate the different seizure types so that appropriate treatments can be titrated accordingly. VEEG assessment of all typical types of episodes in combination with clinical history and seizure semiology is the gold standard for diagnosing functional seizures and differentiating them from epileptic seizures or other physiologic events.⁴⁸

Recommendation 3 Statements

3A. Clinicians *should* evaluate patients diagnosed with functional seizures for co-occurring epilepsy to deliver appropriate treatment. (Level B)

3B. Clinicians *should* use history, semiology, and—where feasible—VEEG to help patients with co-occurring functional and epileptic seizures to accurately identify and distinguish different seizure types to deliver appropriate treatment. (Level B)

3C. Clinicians *should* counsel patients with functional seizures and co-occurring epilepsy about the risks and benefits of antiseizure medications for the treatment of epileptic seizures, as well as their lack of efficacy in the treatment of functional seizures. (Level B)

3D. Clinicians *should* prescribe appropriate antiseizure treatments for epileptic seizures for patients with functional seizures and co-occurring epilepsy. (Level B)

General Principles of Management

Recommendation 4 Rationale

Functional seizures and other forms of functional neurologic disorder are diagnosed based on history, physical examination, and semiology, as well as evaluation of alternative diagnoses and comorbidities as clinically indicated.^{33,48,e22} Psychological interventions may be effective in improving seizure frequency and quality of life. Mental health specialists and other clinicians can be trained in providing these psychological interventions. Neurologists can also facilitate treatment of functional seizures through effective communication of the diagnosis and other approaches to promote engagement (e.g., use of motivational interviewing strategies).^{29,e23} Patients with functional seizures often have comorbidities and other neurologic symptoms, some of which can develop after the diagnosis of functional seizures.^{e24}

Some common clinical practices have been shown to harm patients with functional seizures through stigmatization, humiliation, or inflicting pain.^{e25} Patients with functional seizures are entitled to the same rights as other patients, including dignity, clear and respectful communication, and protection from harm. Patients with functional seizures may have different illness experiences or different understandings of the diagnosis depending on cultural context.^{e26-e28}

The provision of a clear explanation of the diagnosis and treatment plan is an essential platform for further treatment. Development of a seizure action plan can provide increased control and decreased vulnerability for patients and caregivers.^{e29} Many local and national driving restrictions relating to seizures and/or episodic alterations of awareness may apply to patients with functional seizures.^{e30} Functional seizures frequently interfere with employment, schooling, and other social functions.^{23,e31-e33} Effective patient self-management has been linked to improvements in health outcomes for a number of chronic conditions.^{e34-e37} Clinicians can support self-management and self-efficacy by directing patients to educational and support resources. For patients with chronic illnesses, continuity of care is associated with increased patient satisfaction and adherence with treatment plans.^{e38-e40}

Recommendation 4 Statements

4A. Neurologists and mental health clinicians *should* collaborate in the assessment and treatment of functional seizures to facilitate evidence-based treatment. (Level B)

4B. Clinicians *should* adhere to universal standards of care for patients, including speaking respectfully, refraining from unnecessary harm, and avoiding stigmatizing behavior to prevent harm to patients. (Level B)

4C. Clinicians *should* provide a specific diagnostic label and the rationale for the diagnosis to patients and caregivers in

a manner that is clear, empathetic, supportive, and takes into account the patient's cultural context to facilitate patient understanding of the diagnosis and treatment. (Level B)

4D. Clinicians *should* engage in shared decision making regarding the treatment plan, taking into account the patient's understanding of the diagnosis, to facilitate patient engagement with treatment. (Level B)

4E. Clinicians *should* counsel patients and caregivers on how to manage an acute functional seizure episode, to enhance patients' control and decrease vulnerability. (Level B)

4F. Clinicians *should* ask patients of driving age about driving and provide advice appropriate to the regulations of the region of practice to facilitate adherence to relevant regulations. (Level B)

4G. Clinicians *should* ask about the impact of functional seizures on occupational and social functioning to assess patients' psychosocial needs. (Level B)

4H. Clinicians *should* provide or direct the patient and caregivers to resources for learning and support, such as patient and professional advocacy organizations, to promote self-management and self-efficacy. (Level B)

4I. Clinicians *should* provide continuity of care to individuals diagnosed with functional seizures to facilitate treatment and increase patient satisfaction. (Level B)

Psychological Interventions

Recommendation 5 Rationale

Psychological interventions in general can be helpful in possibly achieving seizure freedom, reducing functional seizure frequency, reducing anxiety, and improving health-related quality of life and psychosocial functioning in individuals with functional seizures. NBT, paradoxical therapy, behavioral therapy, psychoeducation, ReACT, and motivational interviewing in combination with psychotherapy have been studied in the treatment of functional seizures, and several studies have examined functional seizure-specific CBT. Studies of NBT, paradoxical therapy, CBT, and motivational interviewing in combination with psychotherapy suggest that these psychological interventions possibly increase the probability of achieving freedom from functional seizures. Studies of ReACT and CBT suggest that these psychological interventions possibly decrease functional seizure frequency. Studies of ReACT, CBT, and motivational interviewing in combination with psychotherapy suggest that these psychological interventions possibly decrease anxiety and increase health-related quality of life. Studies of group psychoeducation and CBT suggest that these psychological interventions possibly improve psychosocial functioning. Studies of CBT suggest that it possibly increases the probability of achieving freedom from functional seizures and possibly improves psychosocial functioning.

Psychological interventions are generally safe and well tolerated.^{e41,e42} Offering available effective and safe treatments is critical to the process of shared decision making between clinicians and patients. The success of psychological

interventions depends, in part, on the support and participation of family, particularly in the treatment of children.^{e43,e44}

The involvement of caregivers or other members of the social support system may also contribute to psychological treatment success for adults.^{e45,e46} Therefore, psychological interventions may differently involve family, caregivers, or others in the patient's social support network, depending on the patient's circumstances.

Recommendation 5 Statements

5A. When psychological interventions for the treatment of functional seizures are indicated and accessible, clinicians *should* counsel patients regarding the potential benefits and risks of such interventions to facilitate shared decision making. (Level B)

5B. When caring for patients with functional seizures, clinicians whose scope of practice does not include counseling patients regarding the possible benefits and demands of psychological interventions *should* refer patients to a clinician knowledgeable about modalities of psychological treatment to facilitate evidence-based treatment. (Level B)

5C. Clinicians *should* refer interested and appropriate patients diagnosed with functional seizures to psychological interventions for the treatment of functional seizures to improve frequency of functional seizures, health-related quality of life, and psychosocial functioning. (Level B)

5D. Clinicians *should*, with patient permission, involve family, caregivers, or others in the social support network in the psychological treatment of adults with functional seizures to improve treatment outcomes. (Level B)

5E. Clinicians *should* involve family in the psychological treatment of children with functional seizures to improve treatment outcomes. (Level B)

Pharmacologic Interventions

Recommendation 6 Rationale

Many patients with functional seizures are initially misdiagnosed with epilepsy and treated with antiseizure medications before the correct diagnosis is made.³⁰ Even after the diagnosis of functional seizures, many patients are prescribed antiseizure medications and/or psychopharmacologic interventions, including benzodiazepines,^{e47} despite insufficient evidence of benefit. Although antiseizure medications can be effective in treating co-occurring epilepsy and psychopharmacologic medications can be effective in treating co-occurring psychiatric disorders, there is a lack of evidence of efficacy of either type of medication in the direct treatment of functional seizures.

While benzodiazepines can provide anxiolysis^{e48} and can sometimes abort prolonged functional seizures in the short term, they are habit forming,^{e49} can cause cognitive impairment,^{e50} and increase risk of motor vehicle accidents in the long term.^{e51} In the acute setting, administration of benzodiazepines for treatment of prolonged functional seizures can result in intubation and iatrogenic harm.^{e52,e53} While

antiseizure medications may provide reassurance to clinicians, patients, or caregivers who are uncertain of the diagnosis of functional seizures, these medications cause adverse effects ranging from more common but mild effects such as fatigue and dizziness, to severe and even life-threatening effects such as Stevens-Johnson syndrome, aplastic anemia, and hepatic failure.^{e54} In patients with functional seizures without co-occurring epilepsy who were previously treated with antiseizure medications, immediate tapering of antiseizure medications leads to improved outcomes compared with delayed withdrawal.^{e55} Besides epilepsy, other indications for antiseizure medications, including potentially benzodiazepines, may include but are not limited to mood disorders, anxiety disorders, migraine, or neuropathic pain.

Recommendation 6 Statements

6A. Clinicians *should* counsel patients with functional seizures without co-occurring epilepsy or another indication for benzodiazepines regarding the potential risks and lack of evidence of benefit for functional seizures associated with benzodiazepines. (Level B)

6B. Clinicians *should not* prescribe benzodiazepines for acute abortive treatment in patients with functional seizures without co-occurring epilepsy, anxiety disorders, or another indication, to reduce the risk of adverse effects. (Level B)

6C. Clinicians *should* counsel patients with functional seizures and without co-occurring epilepsy or another indication for an antiseizure medication about the lack of benefit and the potential risks of taking antiseizure medications for the treatment of functional seizures. (Level B)

6D. Clinicians *should not* prescribe antiseizure medications to patients with functional seizures without co-occurring epilepsy or another indication for antiseizure medications, to reduce the risk of adverse effects. (Level B)

6E. Clinicians *should* taper off antiseizure medications for patients with functional seizures and without another indication for antiseizure medications to reduce the risk of adverse effects. (Level B)

Suggestions for Future Research

The systematic review of evidence relating to the management of functional seizures identified several clinically important areas with limited or no evidence, where future research may be especially impactful. The full guideline (eAppendix 1) includes suggestions for future research relating to diagnosis, pathophysiology, psychological interventions, special populations, psychopharmacologic interventions, neurostimulation, and implementation science.

Disclaimer

Clinical practice guidelines, practice advisories, systematic reviews, Evidence in Focus articles, and other guidance published by the American Academy of Neurology and its affiliates are assessments of current scientific and clinical

information provided as an educational service. The information (1) should not be considered inclusive of all proper treatments or methods of care, or as a statement of the standard of care; (2) is not continually updated and may not reflect the most recent evidence (new evidence may emerge between the time information is developed and when it is published or read); (3) addresses only the question(s) specifically identified; (4) does not mandate any particular course of medical care; (5) represents the medical/scientific opinion of the author panel based on an analysis of then-current clinical data, literature, and public comment (where applicable); and (6) is not intended to substitute for the independent professional judgment of the treating provider, as the information does not account for individual variation among patients. In all cases, the selected course of action should be considered by the treating provider in the context of treating the individual patient. Use of the information is voluntary. The AAN provides this information on an “as is” basis, and makes no warranty, expressed or implied, regarding the information. The AAN specifically disclaims any warranties of merchantability or fitness for a particular use or purpose. The AAN assumes no responsibility for any injury or damage to persons or property arising out of or related to any use of this information or for any errors or omissions.

Conflict of Interest

The American Academy of Neurology (AAN) is committed to producing independent, critical, and trustworthy clinical practice guidelines and evidence-based documents. Significant efforts are made to minimize the potential for conflicts of interest to influence the recommendations of this evidence-based document. Management and disclosure of document developer relationships is conducted in compliance with the 2017 AAN process manual section titled, “Implementing the AAN Conflict of Interest Policy for Guidelines and Case Definitions,” which can be viewed at aan.com.

Author Contributions

B. Tolchin: drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; analysis or interpretation of data. L.H. Goldstein: drafting/revision of the manuscript for content, including medical writing for content; study concept or design; analysis or interpretation of data. M. Reuber: drafting/revision of the manuscript for content, including medical writing for content; study concept or design; analysis or interpretation of data. J. Stone: drafting/revision of the manuscript for content, including medical writing for content; study concept or design; analysis or interpretation of data. D.L. Perez: study concept or design; analysis or interpretation of data; drafting/revision of the manuscript for content, including medical writing for content. W.C. LaFrance: drafting/revision of the manuscript for content,

including medical writing for content; study concept or design; analysis or interpretation of data. A.D. Fobian: drafting/revision of the manuscript for content, including medical writing for content; analysis or interpretation of data. J. Dorman: drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; analysis or interpretation of data. L.H. Hua: drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; analysis or interpretation of data. Z.P. L'Erario: drafting/revision of the manuscript for content, including medical writing for content; analysis or interpretation of data. S.-J. Swanson: drafting/revision of the manuscript for content, including medical writing for content; analysis or interpretation of data. P. Gilli: drafting/revision of the manuscript for content, including medical writing for content; analysis or interpretation of data. B. Mildon: drafting/revision of the manuscript for content, including medical writing for content; analysis or interpretation of data. C. Takahashi: drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; analysis or interpretation of data. K.A. Botchway-Doe: study concept or design. K.P. Hamel: major role in the acquisition of data; study concept or design. H.M. Silsbee: drafting/revision of the manuscript for content, including medical writing for content. M. Oskoui: drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; analysis or interpretation of data.

Study Funding

This practice guideline was developed with financial support from the American Academy of Neurology Institute (AANI). The authors who have served as AAN subcommittee members or methodologists (B.T., J.D., L.H.H., C.T., M.O.), or who are AAN staff members (K.B.D., K.P.H., H.S.), may have been reimbursed by the AAN for expenses related to travel to subcommittee meetings where drafts of manuscripts were reviewed.

Disclosure

B. Tolchin's institution has received research support from the US Department of Veterans Affairs and the C.G. Swebilius Foundation. L.H. Goldstein has received personal compensation in the range of \$0–\$499 (free conference registration) for serving as an officer or member of the Board of Directors for the Functional Neurological Disorder Society and has received publishing royalties from publications relating to health care. The institution of L.H. Goldstein has received research support from the UK National Institute for Health and Care Research and the Motor Neurone Disease Association. M. Reuber has received personal compensation in the range of \$500–\$4,999 for serving as a consultant for Angelini Pharma, Eisai Pharma, and UCB Pharma; has received personal compensation in the range of \$10,000–\$49,999 for serving as an editor, associate editor, or editorial advisory board member for Elsevier; and has received publishing

royalties from Oxford University Press and Jessica Kingsley Publishers for publications relating to health care. The institution of M. Reuber has received compensation in the range of \$500–\$4,999 for M. Reuber's service on a scientific advisory or data safety monitoring board for IQVIA MedTech and has received research support from UCB Pharma. J. Stone has received personal compensation in the range of \$100,000–\$499,999 for serving as an expert witness for multiple UK legal firms; and has received publishing royalties from a publication relating to health care. The institution of J. Stone has received research support from the Scottish Government. D.L. Perez has received personal compensation in the range of \$500–\$4,999 for serving as an editor, associate editor, or editorial advisory board member for *The Journal of Neuropsychiatry and Clinical Neurosciences*; has received personal compensation in the range of \$5,000–\$9,999 for serving as an editor, associate editor, or editorial advisory board member for *Brain and Behavior*; and has received publishing royalties from a publication relating to health care. The institution of D.L. Perez has received research support from the US National Institutes of Health (NIH) and the Sidney R. Baer Jr. Foundation. W.C. LaFrance has received personal compensation in the range of \$10,000–\$49,999 for serving as an expert witness for medicolegal work; and has received author and editor publishing royalties from a publication relating to health care. The institution of W.C. LaFrance has received research support from the US Department of Defense. A.D. Fobian has stock in ReACT FND Health; has received personal compensation in the range of \$0–\$499 (free conference registration) for serving as a board member with the FND Society; and has received intellectual property interests from a discovery or technology relating to health care. A.D. Fobian's institution has received research support from the US National Institute of Mental Health, the National Center for Complementary and Integrative Health, and the National Institute of Diabetes and Digestive and Kidney Diseases. An immediate family member of A.D. Fobian has stock in Gilead and Lab Corp. J. Dorman has received personal compensation for serving as an employee of Cook County Health. J. Dorman's institution has received research support from the NIH. An immediate family member of J. Dorman has received personal compensation for serving as an employee of Rush University and CVS Caremark and has received personal compensation in the range of \$5,000–\$9,999 for serving as a consultant for Merck. L.H. Hua has received personal compensation in the range of \$500–\$4,999 for serving as a consultant for Veila Bio/Horizon, Greenwich, BMS, and Alexion; has received personal compensation in the range of \$5,000–\$9,999 for serving as a consultant for EMD Serono; has received personal compensation in the range of \$10,000–\$49,999 for serving as a consultant for Genzyme and Genentech; has received personal compensation in the range of \$500–\$4,999 for serving on a scientific advisory or data safety monitoring board for Novartis and EMD Serono; has received personal compensation in the range of \$500–\$4,999 for serving on a speakers bureau for Genzyme; and has received personal

compensation in the range of \$5,000–\$9,999 for serving on a speakers bureau for Bristol Myers Squibb and TG Therapeutics. The institution of L.H. Hua has received research support from Biogen and Genentech. Z.P. Lerario has received personal compensation in the range of \$500–\$4,999 for serving as a consultant for Ohio State University, University of Colorado, Sacred Heart University, The LOFT LGBT Community Center, Oregon Stroke Conference, and the Child Neurology Society; has received personal compensation in the range of \$5,000–\$9,999 for serving as an expert witness for Weiss Law; has received publishing royalties from a publication relating to health care; and serves as an editorial board member for *Neurology® Clinical Practice*. S.-J. Swanson has received personal compensation in the range of \$500–\$4,999 for serving as an expert witness for various law firms; and has received personal compensation in the range of \$5,000–\$9,999 for serving as a consultant for Northwestern Mutual. The institution of S.J. Swanson has received personal compensation in the range of \$10,000–\$49,999 for S.-J. Swanson's service as a consultant for Northwestern Mutual. P. Gilli has received personal compensation in the range of \$0–\$499 (free conference registration) for serving as co-chair of a session at the Functional Neurologic Disorder Society biannual meeting. B. Mildon has received personal compensation in the range of \$500–\$4,999 for serving as a speaker-faculty with the American Epilepsy Society; and has received personal compensation in the range of \$500–\$4,999 for serving as an Expert by Experience Partner with the University of College London Centre for Equality Research in Brain Sciences. C. Takahashi has received personal compensation in the range of \$0–\$499 for serving as a consultant for Pfizer; has received personal compensation in the range of \$500–\$4,999 for serving as an expert witness for Keches Law Group, RGLZ, and Meehan, Boyle, Black & Bogdanow; has received personal compensation in the range of \$5,000–\$9,999 for serving as an expert witness for McGowan, Hood, and Felder; has received personal compensation in the range of \$10,000–\$49,999 for serving as an expert witness for Brain IDX and McLeod Law Group; and has received personal compensation in the range of \$500–\$4,999 for serving as an Employee, Founder with Tora Connections. K. Botchway-Doe, K. Puskala Hamel, and H. Silsbee are employees of the American Academy of Neurology. M. Oskoui has received personal compensation in the range of \$500–\$4,999 for serving as an officer or member of the Board of Directors for the Association des Neurologues du Quebec, has a non-compensated relationship as a member of the Medical and Scientific Advisory Committee with Muscular Dystrophy Canada, and has received personal compensation in the range of \$10,000–\$49,999 for serving as a methodologist with the American Academy of Neurology. The institution of M. Oskoui has received research support from Hoffmann-La Roche Ltd., Muscular Dystrophy Canada, Canadian Institutes of Health Research, Santhera, Novartis, Biogen, and Roche Genentech. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

Publication History

Received by *Neurology®* September 3, 2025. Accepted in final form October 31, 2025.

References

- Drane DL, Fani N, Hallett M, Khalsa SS, Perez DL, Roberts NA. A framework for understanding the pathophysiology of functional neurological disorder. *CNS Spectr*. 2020;1-7. doi:10.1017/S1092852920001789
- Tolchin B, Perez DL, Szaflarski JP, et al. What's in a name? *Epilepsy Behav*. 2020;112:107364. doi:10.1016/j.yebeh.2020.107364
- LaFrance WC Jr, Devinsky O. Treatment of nonepileptic seizures. *Epilepsy Behav*. 2002;3(5 suppl):19-23. doi:10.1016/s1525-5069(02)00505-4
- Ludwig L, Pasman JA, Nicholson T, et al. Stressful life events and maltreatment in conversion (functional neurological) disorder: systematic review and meta-analysis of case-control studies. *Lancet Psychiatry*. 2018;5(4):307-320. doi:10.1016/S2215-0366(18)30051-8
- Diprose W, Sundram F, Menkes DB. Psychiatric comorbidity in psychogenic nonepileptic seizures compared with epilepsy. *Epilepsy Behav*. 2016;56:123-130. doi:10.1016/j.yebeh.2015.12.037
- Pareés I, Kojovic M, Pires C, et al. Physical precipitating factors in functional movement disorders. *J Neurol Sci*. 2014;338(1-2):174-177. doi:10.1016/j.jns.2013.12.046
- Stone J, Carson A, Aditya H, et al. The role of physical injury in motor and sensory conversion symptoms: a systematic and narrative review. *J Psychosom Res*. 2009;66(5):383-390. doi:10.1016/j.jpsychores.2008.07.010
- Stager L, Morris S, Szaflarski JP, Fobian AD. Psychiatric and personality factors in pediatric functional seizures: a prospective case-control study. *Seizure*. 2022;98:105-112. doi:10.1016/j.seizure.2022.04.006
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (DSM-5-TR), Fifth Edition, Text Revision ed*. American Psychiatric Association Publishing; 2022.
- Edwards MJ, Yogarajah M, Stone J. Why functional neurological disorder is not feigning or malingering. *Nat Rev Neurol*. 2023;19(4):246-256. doi:10.1038/s41582-022-00765-z
- Aybek S, Perez DL. Diagnosis and management of functional neurological disorder. *BMJ*. 2022;376:o64. doi:10.1136/bmj.o64
- Bean MR, Watson MM, Moore ML, Strom LA. Referral to a functional seizures clinic reduces inpatient and emergency department health care utilization and costs. *Neurol Clin Pract*. 2025;15(1):e200393. doi:10.1212/CPJ.0000000000000393
- Tolchin B, Dworetzky BA, Baslet G. Long-term adherence with psychiatric treatment among patients with psychogenic nonepileptic seizures. *Epilepsia*. 2018;59(1):e18-e22. doi:10.1111/epi.13969
- Hingray C, El-Hage W, Duncan R, et al. Access to diagnostic and therapeutic facilities for psychogenic nonepileptic seizures: an international survey by the ILAE PNES Task Force. *Epilepsia*. 2018;59(1):203-214. doi:10.1111/epi.13952
- Gronseth GS, Cox J, Gloss D, et al, on behalf of the Guideline Development, Dissemination, and Implementation Subcommittee of the American Academy of Neurology. *Clinical Practice Guideline Process Manual*. American Academy of Neurology; 2017.
- Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol*. 2011;64(4):383-394. doi:10.1016/j.jclinepi.2010.04.026
- Aboukasm A, Mahr G, Gahry BR, Thomas A, Barkley GL. Retrospective analysis of the effects of psychotherapeutic interventions on outcomes of psychogenic nonepileptic seizures. *Epilepsia*. 1998;39(5):470-473. doi:10.1111/j.1528-1157.1998.tb01407.x
- Ataoglu A, Ozcetin A, Icmeli C, Ozbulut O. Paradoxical therapy in conversion reaction. *J Korean Med Sci*. 2003;18(4):S81-S84. doi:10.3346/jkms.2003.18.4.S81
- Chen DK, Maheshwari A, Franks R, Trolley GC, Robinson JS, Hrachovy RA. Brief group psychoeducation for psychogenic nonepileptic seizures: a neurologist-initiated program in an epilepsy center. *Epilepsia*. 2014;55(1):156-166. doi:10.1111/epi.12481
- Fobian AD, Long DM, Szaflarski JP. Retraining and control therapy for pediatric psychogenic non-epileptic seizures. *Ann Clin Transl Neurol*. 2020;7(8):1410-1419. doi:10.1002/acn3.51138
- Goldstein LH, Chalder T, Chigwedere C, et al. Cognitive-behavioral therapy for psychogenic nonepileptic seizures: a pilot RCT. *Neurology*. 2010;74(24):1986-1994. doi:10.1212/WNL.0b013e3181e39658
- Goldstein LH, Robinson EJ, Mellers JDC, et al. Cognitive behavioural therapy for adults with dissociative seizures (CODES): a pragmatic, multicentre, randomised controlled trial. *Lancet Psychiatry*. 2020;7(6):491-505. doi:10.1016/S2215-0366(20)30128-0
- Goldstein LH, Robinson EJ, Chalder T, et al. Six-month outcomes of the CODES randomised controlled trial of cognitive behavioural therapy for dissociative seizures: a secondary analysis. *Seizure*. 2022;96:128-136. doi:10.1016/j.seizure.2022.01.016
- Khattak T, Farooq S, Jan B. Behavior therapy in dissociative convulsions disorder. *J Coll Physicians Surg Pak*. 2006;16(5):359-363.
- LaFrance WC Jr, Keitner GI, Papandonatos GD, et al. Pilot pharmacologic randomized controlled trial for psychogenic nonepileptic seizures. *Neurology*. 2010;75(13):1166-1173. doi:10.1212/WNL.0b013e3181f4d5a9

26. LaFrance WC Jr, Baird GL, Barry JJ, et al. Multicenter pilot treatment trial for psychogenic nonepileptic seizures: a randomized clinical trial. *JAMA Psychiatry*. 2014; 71(9):997-1005. doi:10.1001/jamapsychiatry.2014.817
27. Senf-Beckenbach P, Hoheisel M, Devine J, et al. Evaluation of a new body-focused group therapy versus a guided self-help group program for adults with psychogenic non-epileptic seizures (PNES): a pilot randomized controlled feasibility study. *J Neurol*. 2022;269(1):427-436. doi:10.1007/s00415-021-10652-0
28. Tolchin B, Baslet G, Suzuki J, et al. Randomized controlled trial of motivational interviewing for psychogenic nonepileptic seizures. *Epilepsia*. 2019;60(5):986-995. doi:10.1111/epi.14728
29. Tolchin B, Dworetzky BA, Martino S, Blumenfeld H, Hirsch LJ, Baslet G. Adherence with psychotherapy and treatment outcomes for psychogenic nonepileptic seizures. *Neurology*. 2019;92(7):e675-e679. doi:10.1212/WNL.0000000000006848
30. Reuber M, Fernández G, Bauer J, Helmstaedter C, Elger CE. Diagnostic delay in psychogenic nonepileptic seizures. *Neurology*. 2002;58(3):493-495. doi:10.1212/wnl.58.3.493
31. Kerr WT, Zhang X, Hill CE, et al. Factors associated with delay to video-EEG in dissociative seizures. *Seizure*. 2021;86:155-160. doi:10.1016/j.seizure.2021.02.018
32. Avbersek A, Sisodiya S. Does the primary literature provide support for clinical signs used to distinguish psychogenic nonepileptic seizures from epileptic seizures? *J Neurol Neurosurg Psychiatry*. 2010;81(7):719-725. doi:10.1136/jnnp.2009.197996
33. Gilmour GS, MacIsaac R, Subotic A, Wiebe S, Josephson CB. Diagnostic accuracy of clinical signs and symptoms for psychogenic nonepileptic attacks versus epileptic seizures: a systematic review and meta-analysis. *Epilepsy Behav*. 2021;121(pt A):108030. doi:10.1016/j.yebeh.2021.108030
34. Kerr WT, Janio EA, Chau AM, et al. Objective score from initial interview identifies patients with probable dissociative seizures. *Epilepsy Behav*. 2020;113:107525. doi:10.1016/j.yebeh.2020.107525
35. Reuber M, Chen M, Jamnadas-Khoda J, et al. Value of patient-reported symptoms in the diagnosis of transient loss of consciousness. *Neurology*. 2016;87(6):625-633. doi:10.1212/WNL.0000000000002948
36. Reuber M, Monzoni C, Sharrack B, Plug L. Using interactional and linguistic analysis to distinguish between epileptic and psychogenic nonepileptic seizures: a prospective, blinded multirater study. *Epilepsy Behav*. 2009;16(1):139-144. doi:10.1016/j.yebeh.2009.07.018
37. Jenkins L, Cosgrove J, Chappell P, Kheder A, Sokhi D, Reuber M. Neurologists can identify diagnostic linguistic features during routine seizure clinic interactions: results of a one-day teaching intervention. *Epilepsy Behav*. 2016;64(pt A):257-261. doi:10.1016/j.yebeh.2016.08.008
38. Chen M, Jamnadas-Khoda J, Broadhurst M, et al. Value of witness observations in the differential diagnosis of transient loss of consciousness. *Neurology*. 2019;92(9):e895-e904. doi:10.1212/WNL.0000000000007017
39. Bell WL, Park YD, Thompson EA, Radtke RA. Ictal cognitive assessment of partial seizures and pseudoseizures. *Arch Neurol*. 1998;55(11):1456-1459. doi:10.1001/archneur.55.11.1456
40. Muthusamy S, Seneviratne U, Ding C, Phan TG. Using semiology to classify epileptic seizures vs psychogenic nonepileptic seizures: a meta-analysis. *Neurol Clin Pract*. 2022;12(3):234-247. doi:10.1212/CPJ.0000000000001170
41. Seneviratne U, Rajendran D, Brusco M, Phan TG. How good are we at diagnosing seizures based on semiology? *Epilepsia*. 2012;53(4):e63-e66. doi:10.1111/j.1528-1167.2011.03382.x
42. Sen A, Scott C, Sisodiya SM. Stertorous breathing is a reliably identified sign that helps in the differentiation of epileptic from psychogenic non-epileptic convulsions: an audit. *Epilepsy Res*. 2007;77(1):62-64. doi:10.1016/j.eplepsyres.2007.07.009
43. Willert C, Spitzer C, Kussrow S, Runge U. Serum neuron-specific enolase, prolactin, and creatine kinase after epileptic and psychogenic non-epileptic seizures. *Acta Neurol Scand*. 2004;109(5):318-323. doi:10.1046/j.1600-0404.2003.00232.x
44. Laxer KD, Mullooly JP, Howell B. Prolactin changes after seizures classified by EEG monitoring. *Neurology*. 1985;35(1):31-35. doi:10.1212/wnl.35.1.31
45. Patel J, Tran QK, Martinez S, Wright H, Pourmand A. Utility of serum lactate on differential diagnosis of seizure-like activity: a systematic review and meta-analysis. *Seizure*. 2022;102:134-142. doi:10.1016/j.seizure.2022.10.007
46. Abubakar A, Wambacq I. Diagnostic value of serum prolactin levels in PNES in the epilepsy monitoring unit. *Neurol Clin Pract*. 2016;6(2):116-119. doi:10.1212/CJP.0000000000000232
47. Abdelnaby R, Elgenidy A, Heckelmann J, et al. The role of creatine kinase in distinguishing generalized tonic-clonic seizures from psychogenic non-epileptic seizures (PNES) and syncope: a retrospective study and meta-analysis of 1300 patients. *Neurol Res Pract*. 2023;5(1):56. doi:10.1186/s42466-023-00286-0
48. LaFrance WC Jr, Baker GA, Duncan R, Goldstein LH, Reuber M. Minimum requirements for the diagnosis of psychogenic nonepileptic seizures: a staged approach: a report from the International League Against Epilepsy Non-epileptic Seizures Task Force. *Epilepsia*. 2013;54(11):2005-2018. doi:10.1111/epi.12356
49. Nurse ES, Perera T, Hannon T, Wong V, Fernandes KM, Cook MJ. Rates of event capture of home video EEG. *Clin Neurophysiol*. 2023;149:12-17. doi:10.1016/j.clinph.2023.02.165
50. Blad H, Lamberts RJ, van Dijk GJ, Thijs RD. Tilt-induced vasovagal syncope and psychogenic pseudosyncope: overlapping clinical entities. *Neurology*. 2015;85(23):2006-2010. doi:10.1212/WNL.0000000000002184
51. Tatum WO, Hirsch LJ, Gelfand MA, et al. Assessment of the predictive value of outpatient smartphone videos for diagnosis of epileptic seizures. *JAMA Neurol*. 2020; 77(5):593-600. doi:10.1001/jamaneurol.2019.4785
52. Butler M, Shipston-Sharman O, Seynaeve M, et al. International online survey of 1048 individuals with functional neurological disorder. *Eur J Neurol*. 2021;28(11):3591-3602. doi:10.1111/ene.15018
53. Ducroizat A, Zimanti I, Golder D, et al. Functional neurological disorder: clinical manifestations and comorbidities; an online survey. *J Clin Neurosci*. 2023;110:116-125. doi:10.1016/j.jocn.2023.02.014
54. Wasserman D, Herskovitz M. Epileptic vs psychogenic nonepileptic seizures: a video-based survey. *Epilepsy Behav*. 2017;73:42-45. doi:10.1016/j.yebeh.2017.04.020
55. Birca V, Keezer MR, Chamelian L, Lortie A, Nguyen DK. Recognition of psychogenic versus epileptic seizures based on videos. *Can J Neurol Sci*. 2022;49(3):352-360. doi:10.1017/cjn.2021.143
56. Seneviratne U, Ding C, Bower S, Craig S, Leech M, Phan TG. Video-based training improves the accuracy of seizure diagnosis. *J Neurol Neurosurg Psychiatry*. 2014;85(4):466-470. doi:10.1136/jnnp-2013-306618
57. Syed TU, LaFrance WC Jr, Kahrman ES, et al. Can semiology predict psychogenic nonepileptic seizures? A prospective study. *Ann Neurol*. 2011;69(6):997-1004. doi:10.1002/ana.22345
58. Rady A, Elfatry A, Molokhia T, Radwan A. Psychiatric comorbidities in patients with psychogenic nonepileptic seizures. *Epilepsy Behav*. 2021;118:107918. doi:10.1016/j.yebeh.2021.107918
59. Merikangas KR, McClair VL. Epidemiology of substance use disorders. *Hum Genet*. 2012;131(6):779-789. doi:10.1007/s00439-012-1168-0
60. Gorenflo R, Ho R, Carrazana E, et al. Identification of risk factors and distinguishing psychogenic nonepileptic seizures from epilepsy: a retrospective case-control study. *Clin Neurol Neurosurg*. 2022;217:107221. doi:10.1016/j.clineuro.2022.107221