

Multinational Association of Supportive Care in Cancer (MASCC) 2020 clinical practice recommendations for the management of severe gastrointestinal and hepatic toxicities from checkpoint inhibitors.

Title: Multinational Association of Supportive Care in Cancer (MASCC) 2020 clinical practice recommendations for the management of severe gastrointestinal and hepatic toxicities from checkpoint inhibitors.

Publication year: 2020

Journal: Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer

Authors: Massachusetts General Hospital, Boston, MA, USA.

Society: Multinational Association of Supportive Care in Cancer

Origin country: International

Target countries: International

DOI: 10.1007/s00520-020-05707-3

PMID: 32856210

Source: <https://pubmed.ncbi.nlm.nih.gov/32856210/>

pmc Support Care Cancer Support Care Cancer 319 nihpa 9302957 Supportive care in cancer : official journal of the Multinational Association of Supportive Care in Cancer 0941-4355 1433-7339 pmc-is-collection-domain yes pmc-collection-title NIHPA Author Manuscripts PMC8507388 PMC8507388.1 8507388 8507388 NIHMS1743141 32856210 10.1007/s00520-020-05707-3 NIHMS1743141 NIHPA1743141 1 Article MULTINATIONAL ASSOCIATION OF SUPPORTIVE CARE IN CANCER (MASCC) 2020 CLINICAL PRACTICE RECOMMENDATIONS FOR THE MANAGEMENT OF SEVERE GASTROINTESTINAL AND HEPATIC TOXICITIES FROM CHECKPOINT INHIBITORS Dougan Michael 1 Blidner Ada G. 2 Choi Jennifer 3 Cooksley Tim 4 Glezerman Ilya 5 Ginex Pamela 6 Girotra Monica 7 Gupta Dipti 8 Johnson Douglas 9 Shannon Vickie R. 10 Suarez-Almazor Maria 11 Anderson Ronald 12 Rapoport Bernardo L. 12 13 1 Massachusetts General Hospital, Boston, Massachusetts, United States of America. Harvard Medical School, Boston, Massachusetts, United States of America. 2 Laboratory of Immunopathology, Institute of Biology and Experimental Medicine-CONICET, Buenos Aires, Argentina. 3 Division of Oncodermatology, Robert H. Lurie Comprehensive Cancer Center, Northwestern University Feinberg School of Medicine, Chicago, Illinois, United States of America. 4 Tim Cooksley, Manchester University Foundation Trust, Manchester, United Kingdom. The Christie, University of Manchester, Manchester, United Kingdom. 5 Renal Service, Department of Medicine, Memorial Sloan-Kettering Cancer Center, New York, New York, United States of America. 6 Oncology Nursing Society, Pittsburgh, Pennsylvania, United States of America. 7 Endocrine Division, Department of Medicine, Weill Cornell Medical College, and Department of Medicine, Memorial Sloan-Kettering Cancer Center, New York, New York, United States of America. 8 Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, New York, United States of America. 9 Department of Medicine, Vanderbilt University Medical Center and Vanderbilt Ingram Cancer Center, Nashville, Tennessee, United States of America. 10 Department of Pulmonary Medicine, The University of Texas MD Anderson Cancer Center, Houston, United States of America. 11 Section of Rheumatology and Clinical Immunology, University of Texas MD Anderson Cancer Center, Houston, United States of America. 12 Department of Immunology, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa. 13 The Medical Oncology Centre of Rosebank, 129 Oxford Road, Saxonwold 2196, Johannesburg, South Africa. Authors' contributions

All of the authors contributed equally to the conceptualization of the manuscript; MD drafted the manuscript, while BLR and DBJ provided clinical input and BLR, DBJ and RA editorial oversight. All of the authors provided critical appraisal of the manuscript and approve of its submission.

27 8 2020 12 2020 28 12 391689 6129 6143 06 10 2021 01 12 2021 01 12 2021 13 10 2021 11 06 2024 This file is available for text mining. It may also be used consistent with the principles of fair use under the copyright law. Immune-related adverse events (IrAEs) affecting the gastrointestinal (GI) tract and liver are among the most frequent and most severe inflammatory toxicities from contemporary immunotherapy. Inflammation of the colon and or small intestines (entero)colitis is the single most common GI IrAE and is an important cause of delay of discontinuation of immunotherapy. The severity of these GI IrAEs can range from manageable with symptomatic treatment alone to life-threatening complications, including perforation and liver failure. The frequency and severity of GI IrAEs is dependent on the specific immunotherapy given, with cytotoxic T lymphocyte antigen (CTLA)-4 blockade more likely to induce severe GI IrAEs than blockade of either programmed cell death protein 1 (PD-1) or PD-1 ligand (PD-L1), and combination therapy showing the highest rate of GI IrAEs, particularly in the liver. To date, we have minimal prospective data on the appropriate diagnosis and management of GI IrAEs, and recommendations are based largely on retrospective data and expert opinion. Although clinical diagnoses of GI IrAEs are common, biopsy is the gold standard for diagnosis of both immunotherapy-induced enterocolitis and hepatitis and can play an important role in excluding competing, though less common, diagnoses and ensuring optimal management. GI IrAEs typically respond to high dose corticosteroids, though a significant fraction of patients requires secondary immune suppression. For colitis, both TNF- α blockade with infliximab and integrin inhibition with vedolizumab have proved highly effective in corticosteroid-refractory cases. Detailed guidelines have been published for the management of low-grade GI IrAEs. In the setting of more severe toxicities, involvement of a GI specialist is generally recommended. The purpose of this review is to survey the available literature and provide management recommendations focused on the GI specialist.

Inflammation in the gastrointestinal (GI) tract and liver are among the most common immune-related adverse events (IrAEs) resulting from inhibitors of the immune checkpoints cytotoxic T lymphocyte antigen (CTLA)-4, programmed cell death protein 1 (PD-1), and PD-1 ligand (PD-L1) [1]. These toxicities can occur throughout the GI tract from stomach to rectum, and can, less commonly, involve the hepatic parenchyma, biliary tree, pancreas, and potentially the gallbladder [1 – 5]. The mechanisms underlying these inflammatory toxicities are poorly understood, but they are presumed to arise from inhibition of the regulatory mechanisms responsible for preventing T cell immunity to self-antigens and/or commensal microbes [6]. This is consistent with the predominantly lymphocytic inflammation seen on histopathology [7 , 8].

The purpose of these practice recommendations is to provide an overview of the gastrointestinal and hepatic complications of immune checkpoint blockade, including recommendations for diagnosis and for treatment. These practice recommendations will focus specifically on the management of advanced complications that typically require involvement of a gastroenterologist or hepatologist, may benefit from endoscopic evaluation, and may not respond to initial treatment strategies.

Inflammation in the colon (colitis), with or without accompanying inflammation in the small intestine (enterocolitis), is the most common gastrointestinal IrAE associated with current immunotherapies [1 , 9 – 12]. Although isolated colitis is more common than enteritis, the precise frequencies are not yet established [1 , 9 – 12]. Patients can also develop enteritis without colitis, as well as gastroenteritis [1 , 13]. Isolated, symptomatic gastritis is less common [1 , 14 – 16]. In addition to enteritis, a small portion of patients present with newly symptomatic celiac disease, though whether this represents de novo disease or worsening of underlying, undiagnosed celiac disease is unknown [1].

The severity of GI IrAEs can range from readily manageable with lifestyle modification and symptom-directed treatment to life-threatening complications such as perforation [1 , 9 – 12 , 17]. Strictureing disease is rare, and fistulizing disease, a common variant of the inflammatory bowel disease, Crohn's disease, does not appear to occur [1 , 9 – 12 , 18]. The most common presenting symptom of checkpoint inhibitor (entero)colitis is watery, non-bloody diarrhea [1 , 9 – 12 , 19]. In clinical trials and most retrospective studies, diarrhea severity is rated using the Common Terminology Criteria for Adverse Events (CTCAE) using a grade 1 (mild) to grade 5 (death) scale [1 , 20 – 22]. In its mildest form (grade 1), diarrhea from checkpoint inhibitor (entero)colitis presents as <4 extra bowel movements a day, rarely occurring overnight and often associated with eating [1 , 20 – 22]. As symptoms escalate to 4 to 6 extra bowel movements a day, they are defined as grade 2 symptoms [1 , 20 – 22]. These patients often have cramping and urgency, but bleeding is rare [1 , 9 – 12 , 19 – 22]. Grade 3 diarrhea (> 7 extra bowel movements a day) typically presents with cramping, urgency, and can include nocturnal bowel movements and incontinence [1 , 9 – 12 , 19 – 22]. Fevers, severe abdominal pain, and hemodynamic instability are uncommon, but are more often components of grade 4 diarrhea. Deaths from enterocolitis have been reported, but are rare [1 , 9 – 12 , 19 – 22].

Because upper intestinal inflammation happens in a portion of patients with colitis, nausea, vomiting, and decreased appetite can also occur [1 , 20 – 22]. Patients who have isolated gastritis or duodenitis can present primarily with upper GI symptoms, such as nausea and vomiting, epigastric pain, decreased appetite or weight loss [1 , 13]. Although many patients with isolated upper GI inflammation have some diarrhea, some will have normal bowel movements [1 , 13]. Bleeding is less frequent in immunotherapy-induced (entero)colitis than in other forms of colitis, and is often indirectly related to the severity of the colitis, representing hemorrhoids, dermal irritation, or metastatic disease [1 , 20 – 22]. Constipation can occur in patients on immunotherapy, although the degree to which these cases are inflammatory in etiology is unclear, and constipation that rises to the level of medical intervention involving a subspecialist is rare [1].

Blockade of CTLA-4 leads to more frequent IrAEs, often severe IrAEs, in the GI tract than does blockade of PD-1 or PD-L1, though mild GI IrAEs are seen for both [1 , 20 – 22]. Ipilimumab, the only currently approved antibody that targets CTLA-4, causes gastrointestinal symptoms in about 40% of patients when given at the 3 mg/kg standard dose for melanoma [1 , 20 – 23]. The frequency of these symptoms is directly dependent on ipilimumab dose, with the adjuvant melanoma dose of 10 mg/kg having a substantially higher incidence of GI adverse events than occurs with the 3 mg/kg or 1 mg/kg doses [23 , 24]. Severe inflammation that requires urgent management is less common than milder GI inflammation, but still occurs in 10–15% of patients on ipilimumab [1 , 20 – 24].

Patients on PD-L1 inhibitors also have a fairly high incidence of mild GI symptoms with about 20% of patients developing diarrhea of any grade [1 , 20 – 23]. Severe GI toxicities are much less frequent, however, affecting 2–5% of patients [1 , 20 – 23]. At this point, no direct comparisons among the various PD-1 and PD-L1 inhibitors are available, but comparisons across trials suggest that GI IrAEs are reasonably uniform among the available drugs, and are not substantially affected by whether PD-1 or PD-L1 is the target [1 , 20 – 23].

Toxicity from combination checkpoint inhibitor therapy is at least additive, and there may be some synergistic effects on the frequency and severity of GI IrAEs [1]. Diarrhea occurs in about half of patients who receive ipilimumab and nivolumab in combination at the standard melanoma dosing (3 mg/kg ipilimumab and 1 mg/kg nivolumab), with severe GI toxicities occurring in 15–20% of patients [1 , 25]. At the lower doses of ipilimumab (1 mg/kg) used in the more recent regimens (renal cell carcinoma, microsatellite instability high colorectal cancer, lung cancer, etc), severe GI toxicities occur in closer to 5% of combination treated patients [26 – 29].

For patients with mild symptoms (grade 1 and some grade 2), diagnosis is often made clinically based on suggestive symptoms (e.g. new onset diarrhea) in the setting of immunotherapy treatment without another obvious etiology [1 , 20 – 22]. Infectious causes are important to

exclude, as this population is at increased risk for hospital/medical setting acquired infections [1 , 20 – 22]. Depending on the context and pre-test probability as assessed by the treating clinician, stool cultures, stool ova and parasite testing, and testing for *Clostridium difficile* should all be considered as part of the initial diagnostic evaluation [1 , 20 – 22]. In many cases, suggestive symptoms and the exclusion of infections are sufficient to diagnose a patient with a grade 1–2 GI toxicity, which will then typically be managed by the oncology treatment team [1 , 20 – 22].

Cross-sectional imaging can have a role in the diagnosis of colitis from checkpoint inhibitors [30 – 32]. Although the standard appearance is indistinguishable from other forms of colitis (i.e. infectious, ischemic), in a retrospective review of patients on ipilimumab with new onset diarrhea referred for imaging by computed tomography (CT) of the abdomen and a colonoscopy, the CT scan was found to be 85.2% sensitive and 75% specific for the presence of colitis [30 – 32]. This corresponded to a positive predictive value of 95.8% and a negative predictive value of 42.9% [30]. Whether these test characteristics are applicable to PD-1/PD-L1 colitis is unclear at present. Overall, these findings suggest that cross-sectional imaging is valuable for confirming the diagnosis of checkpoint inhibitor enterocolitis in patients with a high clinical likelihood of having the disease, but that CT has limited utility in excluding enterocolitis.

Endoscopic biopsies are the gold standard for diagnosis of GI IrAEs from checkpoint inhibitors, though the importance of endoscopy and tissue biopsies in the evaluation and treatment of these patients has not been rigorously examined [1 , 20 – 22 , 33]. The differential diagnosis in patients with suspected GI toxicities from immunotherapy includes checkpoint inhibitor (entero)colitis (the most likely diagnosis), as well as isolated checkpoint inhibitor (gastro)enteritis [1 , 9 – 13 , 33]. Complications of the malignancy itself can present similarly to inflammatory toxicities [34]. Infections are rare, but important causes of GI symptoms in patients on immunotherapy, as are side-effects from concurrent medications, and other sporadic GI illnesses such as diverticulitis and ischemic colitis [1 , 35 , 36]. Celiac disease can rarely present as new diarrhea in this population, as can pancreatic insufficiency [1 , 2 , 37 , 38].

Most patients with checkpoint inhibitor (entero)colitis, enteritis, or gastritis will respond to first line treatment with high dose corticosteroids and will achieve remission of inflammation after a 4–6 week corticosteroid taper [1 , 9 – 12]. Approximately 30–40% of patients with GI IrAEs from checkpoint inhibitors will require secondary immune suppression with a biologic agent [9]. The best studied biologic therapies are infliximab and vedolizumab, both commonly used for the treatment of inflammatory bowel disease (IBD) [1 , 9 – 12 , 39 – 44].

Recent retrospective analyses have demonstrated that endoscopic findings are the most important clinical factors for predicting resistance to first-line treatment with corticosteroids [10 , 12]. Specifically, the presence of colonic ulcerations on endoscopy suggests a more difficult treatment course [10 , 12]. Importantly, diarrhea grading as determined by the CTCAE was not predictive. Whether having this information prospectively can improve patient outcomes is unclear. However, a recent retrospective analysis found that patients who did receive endoscopic evaluation early in their treatment course tended to have faster symptom resolution and shorter duration exposure to immune suppression [33].

Hepatitis is much less common than (entero)colitis in patients treated with either ipilimumab or PD-L1 monotherapy, but when combined with other agents, hepatitis is considerably more frequent [1 , 3]. Because of the lower incidence compared to (entero)colitis, at this time, much less is known about checkpoint inhibitor hepatitis diagnosis and management than is known for (entero)colitis. The incidence of hepatitis that is detectable on laboratory testing is <5% in clinical trials of checkpoint inhibitor monotherapy, and severe hepatitis is rare [1 , 3 , 45 , 46]. When ipilimumab is combined with nivolumab, the incidence of hepatitis rises to nearly 25% with severe hepatitis in 2–5% of cases [1 , 3 , 45 , 46]. This level of synergy in a toxicity is uncommon with current immunotherapies, where many of the toxicities are usually additive in combination treatments [1 , 3 , 45 , 46]. For this reason, the synergistic toxicity seen in patients for hepatitis suggests that, in the liver, the PD-L1 and CTLA-4 pathways have important functional redundancy in maintaining

immune homeostasis. When immunotherapies are combined with conventional chemotherapies or with targeted agents, liver injury also becomes significantly more common, suggesting that checkpoint inhibitors increase the sensitivity of the liver to other toxic insults [47 – 50].

Liver inflammation induced by immunotherapy is typically detected on routine monitoring and is rarely symptomatic [3 , 45 , 46]. However, patients with cancer are at increased risk for a wide variety of injuries to the liver, including metastatic spread, thromboembolic disease, biliary disease including obstruction, and infection, all of which can be symptomatic [3 , 45 , 46]. For this reason, careful evaluation of abnormal liver tests is essential to making a correct diagnosis and providing appropriate management [3]. In severe liver inflammation (grade 3–4), biopsies have an important role in confirming the diagnosis [3 , 20 – 22]. Typical histologic patterns of checkpoint inhibitor hepatitis have been described, though whether these patterns predict response to treatment or other outcomes is presently unknown [51 – 55]. Interestingly, although spontaneous autoimmune hepatitis (AIH) is a well-described syndrome that we might expect to resemble checkpoint inhibitor hepatitis histologically, the two diseases appear quite distinct [51 – 55]. AIM is characterized by an influx of plasma cells; checkpoint inhibitor hepatitis, on the other hand, is most typically a lymphocytic hepatitis, but can also appear as granulomatous inflammation, while fibrin ring granulomas are a described subtype of PD-1 hepatitis specifically [51 – 55]. Most patients with checkpoint inhibitor hepatitis will respond to corticosteroids, though a small fraction requires secondary immune suppression [3 , 20 – 22]. Several agents have been reported to be effective in these circumstances, including mycophenolate mofetil, tacrolimus, and azathioprine [3 , 20 – 22]. At present, we have no data to recommend one treatment over another, and all 3 are likely to have some deleterious effects on T cell immunity. Anti-thymocyte globulin (ATG) has been reported to be effective in a case of severe, fulminant hepatitis related to checkpoint inhibitors [56]. The role of infliximab in treating checkpoint inhibitor hepatitis is presently unknown, though it is generally considered to be risky given the rare association between infliximab and acute liver injury [57].

Current evaluation and management guidelines for GI adverse events from checkpoint inhibitors are based almost entirely on retrospective analysis and expert opinion [20 – 22]. One prospective study has been published evaluating enteric budesonide compared to placebo for the prevention of diarrhea and colitis associated with ipilimumab treatment [58]. This trial was negative and has generally led to the conclusion that budesonide is ineffective as treatment for immunotherapy-associated enterocolitis, although this trial did not assess therapeutic (i.e. not prophylactic) use [58].

Several comprehensive guidelines have been published recently that focus on evaluation and management of IrAEs directed toward the primary oncology team [20 – 22]. The CTCAE provides definitions for adverse events both based on symptoms (e.g. diarrhea) and clinicopathologic diagnoses (e.g. colitis). Generally, gastroenterologists will become involved when patients develop severe symptoms (CTCAE grade 2–4), and at the point where a pathologic diagnosis is required. Here we provide recommendations targeted toward the consulting gastroenterologist to assist in management of these more severe IrAEs in the GI tract. These recommendations focus on patients with (entero)colitis. Although enteritis and gastroenteritis can also occur in patients on immunotherapy, these other luminal inflammatory syndromes are less common, and are currently managed identically to enterocolitis due to paucity of clinical evidence comparing management strategies in each setting [1 , 13 , 20 – 22 , 59].

Exclusion of infectious causes of diarrhea is important in all patients presenting with Grade 2–4 diarrhea or colitis on immunotherapy. This includes stool culture. Ova and parasite testing should be considered based on risk factors and local prevalence. C. difficile toxin testing is also reasonable in this population, as all of these patients have hospital exposure, and many have had prior exposure to antibiotics.

Fecal calprotectin and lactoferrin are highly sensitive tests for colonic inflammation that are often used in inflammatory bowel disease to rule out luminal inflammation when triaging patients for colonoscopic evaluation [60]. Whether fecal calprotectin or lactoferrin can help stratify patients for

endoscopic evaluation remains unclear at the moment, but is a reasonable strategy if rapid testing is available. Calprotectin may also be useful in monitoring colitis response to treatment in addition to following symptom resolution [33]. Fecal elastase may also be appropriate as an early test in patients presenting with diarrhea on immunotherapy to exclude pancreatic insufficiency, particularly in patients who have not responded adequately to initial management with corticosteroids, or who are presenting with steatorrhea [2]. Serological testing for celiac disease (tissue transglutaminase (TTG)-Immunoglobulin (Ig)A and total IgA) is reasonable, but we advocate confirmatory biopsies by EGD in patients with positive test results [1 , 37 , 38]. Importantly, both newly symptomatic celiac disease and pancreatic insufficiency occurring in the setting of checkpoint inhibitors may be steroid unresponsive [1 , 2 , 37 , 38].

Because of the high risk that patients with immunotherapy-associated enterocolitis will require secondary immune suppression (between 30–40%), we advocate testing for hepatitis B (surface antigen, surface antibody, and core antibody) and for latent tuberculosis (either purified protein derivative (PPD), or tuberculosis enzyme-linked immunospot (ELISpot)) at the time of initial evaluation for enterocolitis [9].

Inflammation in the GI mucosa induced by immunotherapy targeting CTLA-4 and/or PD-L1 can involve any part of the GI tract from the stomach to the rectum [1 , 10 , 12]. Colitis or enterocolitis are the most common types of inflammation [1 , 10 , 12].

Although clinically important, symptoms and CTCAE grade correlate poorly with the extent and severity of mucosal injury found on endoscopy in patients with suspected checkpoint inhibitor colitis [61 , 62]. Perhaps more importantly, symptoms also do not predict response to treatment [61 , 62]. In contrast, the degree of mucosal injury found on endoscopy is the most predictive factor of treatment responsiveness [61 , 62]. Endoscopic evaluation can be useful in identifying patients with milder symptoms who have significant mucosal injury [61]. These patients are less likely to respond to corticosteroids and may require secondary immune suppression [61]. They may also be at higher risk of developing complications should immunotherapy resume, though data addressing this question are currently lacking. At the same time, endoscopy can identify patients with severe symptoms and no visible mucosal injury (checkpoint microscopic colitis); patients with microscopic colitis may respond well to colonic formulations of budesonide alone [62].

In the great majority of patients, colonic inflammation involves the left colon, and while regional variability does occur, most patients can be diagnosed by directed biopsies of the left colon [1 , 7 , 10 , 12]. Thus, flexible sigmoidoscopy, which is both easier on the patient and less expensive than colonoscopy, is often sufficient for making a diagnosis. A portion of patients with luminal inflammation from immunotherapy will have findings isolated to the upper GI tract, although the precise frequency has not been fully defined [1 , 13 , 59]. Esophagogastroduodenoscopy (EGD) is reasonable in patients who have had negative flexible sigmoidoscopies, but who have a sufficiently high likelihood of having luminal inflammation based on symptoms and history, potentially in conjunction with a full colonoscopy.

No guidelines have yet been developed describing appropriate biopsy numbers/coverage. We typically take four gastric biopsies, and four in the duodenum beyond the duodenal bulb during upper endoscopies. Esophagitis from immunotherapy is rare, and we do not biopsy the esophagus in the absence of mucosal changes or symptoms suggestive of esophagitis. In the colon, we typically collect 2–4 biopsies from the descending colon, sigmoid colon, and rectum and pool these unless regional variation is observed on endoscopic evaluation. In the event of localized inflammation, we guide biopsies toward endoscopically abnormal tissue.

Currently we have no rigorous evidence supporting any specific management strategy for the GI toxicities of immunotherapy [9 , 20 – 22 , 39 , 43 , 44]. However, current guidelines universally recommend high-dose corticosteroids as initial management based on extensive experience from patients on immunotherapy clinical trials, as well as subsequent retrospective analyses [20 – 22] and are summarized in Table 1 . Corticosteroids are often initiated by the primary oncology team

using doses between 0.5 and 2 mg/kg of methylprednisolone or equivalent daily [20 – 22]. These are delivered intravenously to patients who are hospitalized. For outpatients, doses closer to those used for IBD (40–60 mg of prednisone) are generally effective [1 , 20 – 22]. Local treatment with colonic budesonide preparations were shown to be ineffective in prophylaxis of CTLA-4 blockade colitis, but may still have a role in the treatment of checkpoint colitis without macroscopic mucosal injury (i.e. microscopic checkpoint colitis) [62]. Steroid tapers are typically performed over 4–6 weeks, depending on the severity of the initial inflammation and the rapidity of the initial response [1 , 20 – 22].

Approximately two thirds of patients will respond to initial management with corticosteroids and will not require any further treatment [1 , 9 , 20 – 22]. The remaining patients require escalation of immune suppression, which is generally done in consultation with a gastroenterologist [1 , 9 , 20 – 22].

For patients who do not respond adequately to corticosteroids, both infliximab and vedolizumab appear to have substantial efficacy in the treatment of enterocolitis associated with CTLA-4 or PD-L1 inhibitors individually or in combination [9 , 20 – 22 , 39 , 43 , 44 , 63]. Alternative antibodies targeting tumor necrosis factor (TNF) α are likely to be effective, though clinical experience is minimal. Both agents are used at standard IBD doses, although infliximab can be dose-escalated in partial responders [9 , 20 – 22 , 39 , 43 , 44 , 63].

The effectiveness of infliximab and vedolizumab has not been directly compared, but both appear to act rapidly, often within a week [9 , 20 – 22 , 39 , 43 , 44 , 63]. Onset of symptom control is generally more rapid than would be expected when treating IBD [9 , 20 – 22 , 39 , 43 , 44 , 63]. Optimal dose and schedule of secondary immune suppression is not established for either agent, though standard IBD dosing is typically sufficient [9 , 20 – 22 , 39 , 43 , 44 , 63]. Between 1 and 3 infusions is typically sufficient for enterocolitis control for both vedolizumab and infliximab, and very few patients ever require maintenance therapy [9 , 20 – 22 , 39 , 43 , 44 , 63]. Recommendations in respect of dose and duration are based on symptom control, with treatment continued only in those patients who have residual symptoms. The utility of calprotectin monitoring or follow-up endoscopy as strategies to guide this decision remains unclear, neither do we know how guided treatment schedules with variable doses compare to fixed doses with 3 infusions in patients irrespective of response.

Early initiation of secondary immune suppression is associated with faster symptom resolution, decreased hospitalization, and decreased risk of enterocolitis recurrence in retrospective analyses [44]. In general, the decision to start secondary immune suppression is made for two reasons: 1) inadequate response to initial treatment with corticosteroids, and 2) recurrence of enterocolitis symptoms during corticosteroid taper. We recommend using the grade of mucosal severity to guide the decision on how quickly to use secondary immune suppression. Patients who have ulcerating disease are treated within 72 hours of corticosteroid initiation if they do not have resolution back to grade 1 symptoms. For less severe enterocolitis a longer duration of steroid treatment may be appropriate before determining that a patient is unresponsive [10 , 12]. Secondary immune suppression is initiated after reemergence of symptoms during a steroid taper. Following initiation of secondary immune suppression, the steroid dose is increased being equivalent to the last effective dose, with tapering commenced on resolution of symptoms.

Several factors should be considered in choosing between infliximab and vedolizumab as treatment for immune-related enterocolitis. Based on currently available evidence, both drugs can be considered as equivalent first-line treatments [9 , 20 – 22 , 39 , 43 , 44 , 63]. Infliximab is a systemically active immune suppressive agent that modulates immune responses, while vedolizumab blocks $\alpha 4\beta 7$ integrin, which primarily affects the trafficking of lymphocytes into the gut. Consequently, patients with active infections of anatomical sites outside the GI tract may be more appropriately treated by vedolizumab. Additionally, vedolizumab should be avoided in patients with GI malignancies or metastases (approximately 5% of patients with melanoma) [34 , 64]. The impact of these two drugs on anti-tumor immunity in humans is unknown, though vedolizumab is

unlikely to have deleterious effects on anti-tumor immune responses outside of the gut. Blockade of TNF α with agents such as infliximab, has been linked to improved anti-tumor responses in murine models of anti-tumor immunity [65 , 66]. Infliximab has been associated with an increased incidence of melanoma in a meta-analysis of patients with IBD, but not in larger analyses of patients treated for rheumatologic conditions [67 , 68]. Retrospective analyses of patients with melanoma treated with immunotherapy that developed colitis and received infliximab have shown no negative impact on tumor outcomes [42]. Irrespective of the aforementioned issues, drug availability and coverage are important considerations and may ultimately be the principal deciding factors for determining selection between infliximab and vedolizumab in most clinical scenarios.

A small fraction of patients with immune-related enterocolitis will fail to respond to corticosteroids as well as infliximab or vedolizumab. In these patients, confirmation of ongoing inflammation and exclusion of opportunistic infections is essential. Based on patient risk factors, recommended investigations should include repeat stool cultures, *C. difficile* testing, as well as ova and parasite testing.

Although measurement of calprotectin or lactoferrin may be useful in suggesting ongoing luminal inflammation, endoscopy should be considered in all of these patients. These investigations are necessary to provide evidence of mucosal healing and to confirm ongoing inflammation histopathologically. Additional differential diagnoses include cytomegalovirus (CMV) infection, other opportunistic viral infections, and fungal colitis. Provided that these infections are excluded, and ongoing significant inflammation is confirmed, escalation of immune suppression is appropriate.

Patients with immunotherapy-induced enterocolitis often do not have the time to wait for a full washout period before initiation of alternative immune suppression. Initiation of alternative immune suppression often begins within 2–4 weeks of treatment with the prior biologic therapy [39].

Current evidence, albeit limited, suggest that switching from infliximab to vedolizumab, or vice versa, is the most appropriate management strategy after failure of the initial choice of biologic therapy [39]. The response rates to third-line immune suppression are generally lower than would be expected for the same agent as second-line treatment. In patients who have failed both biologic therapies, fecal microbiota transplant (FMT) can be considered depending on accessibility [69]. Clinical trials are ongoing to determine whether FMT is appropriate as earlier management and to define response rates. FMT has shown some efficacy in IBD, but is known to be highly effective in *C. difficile* colitis, suggesting that colonic dysbiosis or occult infections may play an important role in driving ongoing inflammation in some patients with immune-related enterocolitis from immunotherapy [70 , 71].

Whether other forms of immune suppression that are effective in IBD are also effective in immune-related enterocolitis is not clear. Mesalamine has been used in some cases of mild enterocolitis. Immunomodulatory drugs such as 6-mercaptopurine or methotrexate typically has a slow onset of therapeutic efficacy in IBD and may not act fast enough to be clinically useful in severe cases of immune-related enterocolitis. Although unproven, these drugs may also have a deleterious effect on the anti-tumor response.

In very refractory cases, alternative biologic agents such as ustekinumab would be reasonable to try, though currently we have no published evidence of efficacy [72]. Based on comprehensive analyses of the correlates of effective anti-tumor immunity, and the pathways of immunotherapy resistance, signaling through Janus kinase (JAK) kinases is likely to play an important role in anti-cancer treatment. Although JAK inhibitors (e.g. tofacitinib) present a potential therapeutic option, this remains to be proven and should be used with extreme caution in this patient population. Similarly, CTLA-4Ig (e.g. abatacept, belatacept) is likely to be effective in CTLA-4 blockade colitis, and potentially PD-1 blockade colitis, given its efficacy in colitis associated with CTLA-4 haploinsufficiency [73 – 80]. Yet the mechanism of action of CTLA-4Ig also means that it is likely to interfere with effective anti-tumor responses. All the above mention strategies are not proven and requires clinical studies to confirm the efficacy.

Although the following algorithm focuses on diarrhea as the presenting symptom, other GI symptoms such as nausea, vomiting, and decreased appetite can be managed similarly. For primarily upper GI symptoms, upper endoscopy is favored over lower endoscopy for evaluation. For patients with decreased appetite, weight loss, or abdominal pain, cross-sectional imaging has a critical role in management for any severe (grade 3–4) symptoms. The rationale for these grading-based recommendations is presented in detail in the preceding sections (Table 1).

Low-grade diarrhea - somewhere between 1 and 3 extra bowel movements a day – is frequent with current immunotherapy and is managed with symptomatic control. This can include motility slowing agents such as loperamide or diphenoxylate-atropine, dietary modification including bulking agents like psyllium, and lifestyle changes. Immunotherapy is generally continued for grade 1 symptoms. In addition to diarrhea, low-grade nausea, vomiting, and decreased appetite can all occur from immunotherapy, as can, albeit less frequently, pain and cramping.

The evaluation and management of grade 2 diarrhea (4 to 7 additional bowel movements a day) is less well established. This is similar for nausea, vomiting, and decreased appetite. Many of these patients are managed by their oncology teams through temporary holding of immunotherapy and empiric treatment courses with corticosteroids. However, some of these patients are referred to subspecialists for tissue diagnoses. In these instances, the endoscopy can be valuable for two reasons. The first is to identify those patients who do not have mucosal inflammation at all. In these patients, corticosteroids are not likely to be beneficial, and alternative treatments can be considered. Because symptoms correlate poorly with the extent and severity of mucosal disease, and do not predict response to therapy, endoscopic evaluation plays an important role in management decisions. The identification of mucosal ulcerations increases the likelihood that a patient will require secondary immune suppression to control checkpoint colitis, while identification of microscopic colitis may indicate responsiveness to colonic formulations for budesonide.

Corticosteroids are first-line therapy for biopsy proven grade 2 enterocolitis. No rigorous studies have examined steroid dose. Typically, patients will respond to doses in the range of 0.5–1 mg/kg oral prednisone or, for inpatients, intravenous methylprednisolone. Doses of 40–60 mg of oral prednisone may also be effective in many cases. Steroid tapers generally occur over a 4–6 week period and can be guided by the severity of mucosal inflammation observed on biopsy, with slower tapers for patients with colonic ulcers, and rapid tapers for patients who have histologic inflammation without endoscopic evidence of inflammation.

Grade 3 diarrhea is often managed in conjunction with a specialist in gastroenterology. Patients can have grade 3 diarrhea either because they failed to respond to initial management for grade 2 diarrhea, or because they initially present with severe diarrhea (> 7 extra bowel movements a day). Grade 4 diarrhea is associated with life-threatening complications such as end-organ injury, severe hypotension, anemia, etc. Many of these patients have significant mucosal inflammation, and endoscopic evaluation is recommended in all of them, ideally prior to initiation of corticosteroids.

The purpose of endoscopy in this group of patients is several-fold. First, a small, but significant subset of these patients will have an alternative cause for their diarrhea. This includes infectious and ischemic colitis, new onset celiac disease, pancreatic insufficiency (endoscopically normal), and other medication-induced diarrhea (which will often show minimal mucosal changes) [2 , 61 , 62 , 81]. This is particularly important in patients who are initially referred to a gastroenterologist after having failed first-line corticosteroids. The second reason for endoscopic evaluation is to determine mucosal severity. Similar to the rationale for grade 2 symptoms, determining which of these patients has severe mucosal injury can help guide the transition to more rapid initiation of secondary immune suppression [61] and identifying microscopic colitis can help avoid systemic corticosteroids altogether [62].

Corticosteroids remain first line therapy for biopsy proven grade 3/4 enterocolitis, although initial corticosteroid doses are typically 1–2 mg/kg methylprednisolone or equivalent. Some evidence suggests that earlier initiation of secondary immune suppression may be beneficial in patients with more severe GI adverse events, though this evidence is largely based on mucosal severity rather than the severity of the presenting symptoms.

ALT/AST upper limit of normal (ULN) to $3 \times \text{ULN}$, ALKP ULN to $2.5 \times \text{ULN}$, TBILI ULN to $1.5 \times \text{ULN}$

Frequent blood monitoring is indicated (1–2 $\times$ weekly). A patient history of concomitant drugs usage (including all over-the-counter medications), herbal supplements, as well as alcohol use should be obtained. Immunotherapy does not need to be delayed, and sub-specialists are typically not involved.

ALT/AST $>3 - 5 \times \text{ULN}$, ALKP $>2.5 - 5 \times \text{ULN}$, TBILI $>1.5 - 3 \times \text{ULN}$

Twice weekly monitoring, and withholding immunotherapy. Patients with grade 2 hepatitis should be investigated for potential non-immune mediated etiologies. These investigation includes a right upper quadrant ultrasound, hepatitis A, B, and C serology, Epstein-Barr virus (EBV) and CMV testing. Autoimmune disease serology, iron studies, and measurement of ceruloplasmin, and alpha-1-antitrypsin (A1AT) testing (in selected patients based on prior history). As for all patients with abnormal liver testing results, concomitant medications and alcohol use should be assessed. For patients with a predominantly cholestatic hepatitis, advanced imaging of the biliary tract should be considered such as magnetic resonance cholangiopancreatography (MRCP) or endoscopic ultrasound (EUS) if indicated. Patients with abdominal malignancies or known hepatic metastases should also undergo cross-sectional imaging. For patients without another clear etiology for their hepatitis and with a predominantly hepatocellular pattern of injury, checkpoint hepatitis is the most likely diagnosis and empiric treatment with 0.5 – 1 mg/kg prednisone or prednisone equivalent is a reasonable treatment option. Whether biopsy should be obtained prior to initiation of corticosteroids in these patients is presently unclear. For patients who do not respond within one week to corticosteroids with a least a 50% reduction in laboratory values, a biopsy is indicated to confirm the diagnosis. Secondary immune suppression should be considered if the diagnosis is confirmed. Azathioprine [1–2 mg/kg), mycophenolate mofetil (500 – 1000 mg BID), and tacrolimus [targeting blood levels of 8–10 nanograms (ng)/ml or lower if a response is detected early] have all been used as secondary immune suppression in these patients even though optimal doses and dosing schedules have not been determined.

ALT/AST $>5 \times \text{ULN}$, ALKP $>5 \times \text{ULN}$, TBILI $>3 \times \text{ULN}$

Patients with higher-grade hepatic injury will typically be referred to Hepatology. In addition to all of the considerations for grade 2 hepatic injury, we recommend performing biopsies on all of these patients to exclude a non-immune-mediated cause of the liver injury, given the severity of the detected laboratory changes, and the potential clinical impact of missing an alternative diagnosis. Treatment for confirmed Grade 3/4 checkpoint hepatitis is similar to that of Grade 2 hepatitis, though doses of corticosteroid as high as 2 mg/kg could be considered in severe cases. Failure to respond within one week to corticosteroids with at least a 50% reduction in laboratory values should prompt the addition of secondary immune suppression. Azathioprine (1–2 mg/kg), mycophenolate mofetil (500 – 1000 mg BID), and tacrolimus (targeting blood levels or 8–10 ng/ml or lower if a response is detected early) have all been used as secondary immune suppression in these patients even though, as mentioned above, optimal doses and dosing schedules remains to be determined. In severe, fulminant hepatitis, ATG is another treatment option. The role of infliximab in the management of immune-checkpoint hepatitis remains unclear.

Immune-mediated adverse events in the GI tract and liver are important limitations on current checkpoint inhibitor therapies. Although we are beginning to develop a robust understanding of the clinical presentations of these IrAEs, our understanding of the immune mechanisms that drive them

remains inconclusive. Several empiric treatments have been developed for these toxicities. These appear to have reasonable efficacy based on retrospective analyses, although optimal diagnostic and treatment strategies have not been established in prospective clinical trials. For this reason, we do not yet know if any specific treatment strategy is more efficacious than another. Moreover, there is a limited understanding of how these GI and hepatic irAEs affect cancer treatment outcomes, and how treatment of these irAEs may influence the anti-tumor immune response.

Future work should focus on developing a clear understanding of the immune mechanisms that drive GI and hepatic irAEs. Clinical trials should prospectively compare various treatment strategies using both organ-specific outcomes (e.g. resolution of colitis), and tumor-specific outcomes (e.g. progression-free and overall survival). These trials will be necessary for the establishment of truly evidence-based recommendations to optimize outcomes for these patients.

Professor BL Rapoport is supported by supported by the Cancer Association of South Africa (CANSA) and the National Research Foundation (NRF) of South Africa.

Dr. I. Glezerman is supported by the NIH/NCI (Cancer Center Support Grant P30 CA008748)

AB, RA, JC, TC, PG, DG and VRS have no conflict of interest to declare. MD reports grants from Novartis, other (SAB) from Neoleukin Therapeutics, personal fees from Partner Therapeutics, personal fees from Tillotts Pharma, grants from Genentech, outside the submitted work. MG reports consultant work with Bristol Myers Squibb (BMS), and AstraZeneca, outside the submitted work. IG reports other (Stock Ownership) from Pfizer Inc., personal fees from CytomX Inc, outside the submitted work. DBJ reports other (advisory board) from Array Biopharma, grants and other (advisory board) from BMS, other (advisory board) from Jansen, grants from Incyte, other (advisory board) from Merck, other (advisory board) from Novartis, outside the submitted work. In addition, DBJ has a patent Co-inventor on use of CTLA-4 agonist for IAEs pending. BLR reports personal fees and other (advisory board) from Merck and Co, grants, personal fees and other (advisory board) from BMS, grants, personal fees and other (advisory board) from Roche South Africa, personal fees and other (advisory board) from AstraZeneca, during the conduct of the study. MSA reports personal fees from Gilead, grants from Pfizer, personal fees from Abbvie, outside the submitted work.

10.3389/fimmu.2017.01547 . 29230210 PMC5715331 2. Eshet Y , Baruch EN , Shapira-Frommer R , Steinberg-Silman Y , Kuznetsov T , Ben-Betzalel G , Daher S , Gluck I , Asher N , Apter S , Schachter J , Bar J , Boursi B , Markel G (2018) Clinical significance of pancreatic atrophy induced by immune-checkpoint inhibitors: a case-control study . *Cancer Immunol Res* 6 (12): 1453 – 1458 . doi: 10.1158/2326-6066.CIR-17-0659 . 30275274 3. Reynolds K , Thomas M , Dougan M (2018) Diagnosis and management of hepatitis in patients on checkpoint blockade . *Oncologist* 23 (9): 991 – 997 . doi: 10.1634/theoncologist.2018-0174 . 29853659 PMC6192661 4. Raschi E , Mazzearella A , Antonazzo IC , Bendinelli N , Forcesi E , Tuccori M , Moretti U , Poluzzi E , De Ponti F (2019) Toxicities with immune checkpoint inhibitors: emerging priorities from disproportionality Analysis of the FDA Adverse Event Reporting System . *Target Oncol* 14 (2): 205 – 221 . doi: 10.1007/s11523-019-00632-w . 30927173 5. Abu-Sbeih H , Tran CN , Ge PS , Bhutani MS , Alasadi M , Naing A , Jazaeri AA , Wang Y (2019) Case series of cancer patients who developed cholecystitis related to immune checkpoint inhibitor treatment . *J Immunother Cancer* 7 (1): 118 . doi: 10.1186/s40425-019-0604-2 . 31053161 PMC6499962 6. Pauken KE , Dougan M , Rose NR , Lichtman AH , Sharpe AH (2019) Adverse events following cancer immunotherapy: obstacles and opportunities . *Trends Immunol* 40 (6): 511 – 523 . doi: 10.1016/j.it.2019.04.002 . 31053497 PMC6527345 7. Marthey L , Mateus C , Mussini C , Nachury M , Nancey S , Grange F , Zallot C , Peyrin-Biroulet L , Rahier JF , Bourdier de Beauregard M , Mortier L , Coutzac C , Soularue E , Lanoy E , Kapel N , Planchard D , Chaput N , Robert C , Carbonnel F (2016) Cancer immunotherapy with anti-CTLA-4 monoclonal antibodies induces an inflammatory bowel disease . *J Crohns Colitis* 10 (4): 395 – 401 . doi: 10.1093/ecco-jcc/jjv227 . 26783344 PMC4946758 8. Chen JH , Pezhouh MK , Lauwers GY , Masia R (2017) Histopathologic features of colitis due to immunotherapy with anti-PD-1 antibodies . *Am J Surg Pathol* 41 (5): 643 – 654 . doi:

10.1097/PAS.0000000000000829 . 28296676 9. Beck KE , Blansfield JA , Tran KQ , Feldman AL , Hughes MS , Royal RE , Kammula US , Topalian SL , Sherry RM , Kleiner D , Quezado M , Lowy I , Yellin M , Rosenberg SA , Yang JC (2006) Enterocolitis in patients with cancer after antibody blockade of cytotoxic T-lymphocyte-associated antigen 4 . *J Clin Oncol* 24 (15) : 2283 – 2289 . doi: 10.1200/JCO.2005.04.5716 . 16710025 PMC2140223 10. Geukes Foppen MH , Rozeman EA , van Wilpe S , Postma C , Snaebjornsson P , van Thienen JV , van Leerdam ME , van den Heuvel M , Blank CU , van Dieren J , Haanen JBAG (2018) Immune checkpoint inhibition-related colitis: symptoms, endoscopic features, histology and response to management . *ESMO Open* 3 (1) : e000278 . doi: 10.1136/esmoopen-2017-000278 . 29387476 PMC5786923 11. Wang DY , Mooradian MJ , Kim D , Shah NJ , Fenton SE , Conry RM , Mehta R , Silk AW , Zhou A , Compton ML , Al-Rohil RN , Lee S , Voorhees AL , Ha L , McKee S , Norrell JT , Mehnert J , Puzanov I , Sosman JA , Chandra S , Gibney GT , Rapisuwon S , Eroglu Z , Sullivan R , Johnson DB (2018) Clinical characterization of colitis arising from anti-PD-1 based therapy . *Oncoimmunology* 8 (1) : e1524695 . doi: 10.1080/2162402X.2018.1524695 . 30546965 PMC6287774 12. Wang Y , Abu-Sbeih H , Mao E , Ali N , Qiao W , Trinh VA , Zobniw C , Johnson DH , Samdani R , Lum P , Shuttlesworth G , Blechacz B , Bresalier R , Miller E , Thirumurthi S , Richards D , Raju G , Stroehlein J , Diab A (2018) Endoscopic and histologic features of immune checkpoint inhibitor-related colitis . *Inflamm Bowel Dis* 24 (8) : 1695 – 1705 . 29718308 10.1093/ibd/izy104 13. Sokal A , de Chou CS , Delyon J , Roche B , Lourenco N , Lebbe C , Baroudjian B ; PATIO group (2018) Enteritis without colitis in patients treated with immune checkpoint inhibitors: a tricky diagnosis . *Melanoma Res* 28 (5) : 483 – 484 . doi: 10.1097/CMR.0000000000000484 . 30148767 14. Yip RHL , Lee LH , Schaeffer DF , Horst BA , and Yang HM (2018) Lymphocytic gastritis induced by pembrolizumab in a patient with metastatic melanoma . *Melanoma Res* 28 (6) : 645 – 647 . doi: 10.1097/CMR.0000000000000502 . 30256271 15. Nishimura Y , Yasuda M , Ocho K , Iwamuro M , Yamasaki O , Tanaka T , Otsuka F (2018) Severe gastritis after administration of nivolumab and ipilimumab . *Case Rep Oncol* 11 (2) : 549 – 556 . doi: 10.1159/000491862 . 30186138 PMC6120399 16. Boike J , Dejulio T (2017) Severe esophagitis and gastritis from nivolumab therapy . *ACG Case Rep J* 4 : e57 . doi: 10.14309/crj.2017.57 . 28459081 PMC5404341 17. Celli R , Kluger HM , Zhang X (2018) Anti-PD-1 therapy-associated perforating colitis . *Case Rep Gastrointest Med* 2018 : 3406437 . doi: 10.1155/2018/3406437 29955400 PMC6000840 18. Abraham C , Cho JH (2009) Inflammatory bowel disease . *N Engl J Med* 361 (21) : 2066 – 2078 . doi: 10.1056/NEJMr0804647 . 19923578 PMC3491806 19. Wang Y , Abu-Sbeih H , Mao E , Ali N , Ali FS , Qiao W , Lum P , Raju G , Shuttlesworth G , Stroehlein J , Diab A (2018) Immune-checkpoint inhibitor-induced diarrhea and colitis in patients with advanced malignancies: retrospective review at MD Anderson . *J Immunother Cancer* 6 (1) : 37 . doi: 10.1186/s40425-018-0346-6 . 29747688 PMC5946546 20. Brahmer JR , Lacchetti C , Schneider BJ , Atkins MB , Brassil KJ , Caterino JM , Chau I , Ernstoff MS , Gardner JM , Ginex P , Hallmeyer S , Holter Chakrabarty J , Leighl NB , Mammen JS , McDermott DF , Naing A , Nastoupil LJ , Phillips T , Porter LD , Puzanov I , Reichner CA , Santomaso BD , Seigel C , Spira A , Suarez-Almazor ME , Wang Y , Weber JS , Wolchok JD , Thompson JA ; National Comprehensive Cancer Network (2018) Management of immune-related adverse events in patients treated with immune checkpoint inhibitor therapy: American Society of Clinical Oncology Clinical Practice Guideline . *J Clin Oncol* 36 (17) : 1714 – 1768 . doi: 10.1200/JCO.2017.77 . 29442540 PMC6481621 21. Haanen J , Carbone F , Robert C , Kerr KM , Peters S , Larkin J , Jordan K ; ESMO Guidelines Committee (2017) Management of toxicities from immunotherapy: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up . *Ann Oncol* 28 (suppl_4) : iv119 – iv142 . doi: 10.1093/annonc/mdx225 . 28881921 22. Puzanov I , Diab A , Abdallah K , Bingham CO 3rd , Brogdon C , Dadu R , Hamad L , Kim S , Lacouture ME , LeBoeuf NR , Lenihan D , Onofrei C , Shannon V , Sharma R , Silk AW , Skondra D , Suarez-Almazor ME , Wang Y , Wiley K , Kaufman HL , Ernstoff MS (2017) Managing toxicities associated with immune checkpoint inhibitors: consensus recommendations from the Society for Immunotherapy of Cancer (SITC) Toxicity Management Working Group . *J Immunother Cancer* 5 (1) : 95 . doi: 10.1186/s40425-017-0300-z . 29162153 PMC5697162 23. Hodi FS , O'Day SJ , McDermott DF , Weber RW , Sosman JA , Haanen JB , Gonzalez R , Robert C , Schadendorf D , Hassel JC , Akerley W , van den Eertwegh AJ , Lutzky J , Lorigan P , Vaubel JM , Linette GP , Hogg D , Ottensmeier CH , Lebbe C , Peschel C , Quirt I , Clark JI , Wolchok JD , Weber JS , Tian J , Yellin MJ , Nichol GM , Hoos A , Urba WJ (2010) Improved survival with ipilimumab in patients with

metastatic melanoma . *N Engl J Med* 363 (8): 711 – 723 . doi: 10.1056/NEJMoa1003466 . 20525992 PMC3549297

24. Eggermont AM , Chiarion-Sileni V , Grob JJ , Dummer R , Wolchok JD , Schmidt H , Hamid O , Robert C , Ascierto PA , Richards JM , Lebbé C , Ferraresi V , Smylie M , Weber JS , Maio M , Bastholt L , Mortier L , Thomas L , Tahir S , Hauschild A , Hassel JC , Hodi FS , Taitt C , de Pril V , de Schaetzen G , Suci S , Testori A (2016) Prolonged survival in stage III melanoma with ipilimumab adjuvant therapy . *N Engl J Med* 375 (19): 1845 – 1855 . doi: 10.1056/NEJMoa1611299 27717298 PMC5648545

25. Wolchok JD , Chiarion-Sileni V , Gonzalez R , Rutkowski P , Grob JJ , Cowey CL , Lao CD , Wagstaff J , Schadendorf D , Ferrucci PF , Smylie M , Dummer R , Hill A , Hogg D , Haanen J , Carlino MS , Bechter O , Maio M , Marquez-Rodas I , Guidoboni M , McArthur G , Lebbé C , Ascierto PA , Long GV , Cebon J , Sosman J , Postow MA , Callahan MK , Walker D , Rollin L , Bhore R , Hodi FS , Larkin J (2017) Overall survival with combined nivolumab and ipilimumab in advanced melanoma . *N Engl J Med* 377 (14): 1345 – 1356 . doi: 10.1056/NEJMoa1709684 . 28889792 PMC5706778

26. Hammers HJ , Plimack ER , Infante JR , Rini BI , McDermott DF , Lewis LD , Voss MH , Sharma P , Pal SK , Razak ARA , Kollmannsberger C , Heng DY , Sprattlin J , McHenry MB , Amin A (2017) Safety and efficacy of nivolumab in combination with ipilimumab in metastatic renal cell carcinoma: The CheckMate 016 Study . *J Clin Oncol* 35 (34): 3851 – 3858 . doi: 10.1200/JCO.2016.72.1985 28678668 PMC7587408

27. Hellmann MD , Ciuleanu TE , Pluzanski A , Lee JS , Otterson GA , Audigier-Valette C , Minenza E , Linardou H , Burgers S , Salman P , Borghaei H , Ramalingam SS , Brahmer J , Reck M , O'Byrne KJ , Geese WJ , Green G , Chang H , Szustakowski J , Bhagavatheeswaran P , Healey D , Fu Y , Nathan F , Paz-Ares L (2018) Nivolumab plus ipilimumab in lung cancer with a high tumor mutational burden . *N Engl J Med* 378 (22): 2093 – 2104 . doi: 10.1056/NEJMoa1801946 . 29658845 PMC7193684

28. Motzer RJ , Tannir NM , McDermott DF , Aren Frontera O , Melichar B , Choueiri TK , Plimack ER , Barthélémy P , Porta C , George S , Powles T , Donskov F , Neiman V , Kollmannsberger CK , Salman P , Gurney H , Hawkins R , Ravaud A , Grimm MO , Bracarda S , Barrios CH , Tomita Y , Castellano D , Rini BI , Chen AC , Mekan S , McHenry MB , Wind-Rotolo M , Doan J , Sharma P , Hammers HJ , Escudier B ; CheckMate 214 Investigators (2018) Nivolumab plus ipilimumab versus sunitinib in advanced renal-cell carcinoma . *N Engl J Med* 378 (14): 1277 – 1290 . doi: 10.1056/NEJMoa1712126 . 29562145 PMC5972549

29. Overman MJ , Lonardi S , Wong KYM , Lenz HJ , Gelsomino F , Aglietta M , Morse MA , Van Cutsem E , McDermott R , Hill A , Sawyer MB , Hendlish A , Neyns B , Svrcek M , Moss RA , Ledezine JM , Cao ZA , Kamble S , Kopetz S , André T (2018) Durable clinical benefit with nivolumab plus ipilimumab in dna mismatch repair-deficient/microsatellite instability-high metastatic colorectal cancer . *J Clin Oncol* 36 (8): 773 – 779 . doi: 10.1200/JCO.2017.76.9901 . 29355075

30. Garcia-Neuer M , Marmarelis ME , Jangi SR , Luke JJ , Ibrahim N , Davis M , Weinberg J , Donahue H , Bailey N , Hodi FS , Buchbinder EL , Ott PA (2017) Diagnostic comparison of ct scans and colonoscopy for immune-related colitis in ipilimumab-treated advanced melanoma patients . *Cancer Immunol Res* 5 (4): 286 – 291 . doi: 10.1158/2326-6066.CIR-16-0302 . 28373217

31. Kim KW , Ramaiya NH , Krajewski KM , Shinagare AB , Howard SA , Jagannathan JP , Ibrahim N (2013) Ipilimumab-associated colitis: CT findings . *AJR Am J Roentgenol* 200 (5): W468 – W474 . doi: 10.2214/AJR.12.9751 . 23718569

32. Barina AR , Bashir MR , Howard BA , Hanks BA , Salama AK , and Jaffe TA (2016) Isolated recto-sigmoid colitis: a new imaging pattern of ipilimumab-associated colitis . *Abdom Radiol (NY)* 41 (2): 207 – 214 . doi: 10.1007/s00261-015-0560-3 . 26867901

33. Abu-Sbeih H , Ali FS , Luo W , Qiao W , Raju GS , and Wang Y (2018) Importance of endoscopic and histological evaluation in the management of immune checkpoint inhibitor-induced colitis . *J Immunother Cancer* 6 (1): 95 . doi: 10.1186/s40425-018-0411-1 . 30253811 PMC6156850

34. Bello E , Cohen JV , Mino-Kenudson M , Dougan M (2019) Antitumor response to microscopic melanoma in the gastric mucosa mimicking ipilimumab-induced gastritis . *J Immunother Cancer* 7 (1): 41 . doi: 10.1186/s40425-019-0524-1 . 30744698 PMC6371540

35. Lankes K , Hundorfean G , Harrer T , Pommer AJ , Agaimy A , Angelovska I , Tajmir-Riahi A , Göhl J , Schuler G , Neurath MF , Hohenberger W , Heinzerling L (2016) Anti-TNF-refractory colitis after checkpoint inhibitor therapy: Possible role of CMV-mediated immunopathogenesis . *Oncoimmunology* 5 (6): e1128611 . doi: 10.1080/2162402X.2015.1128611 . 27471608 PMC4938313

36. Zhou C , Klionsky Y , Treasure ME , Bruno DS (2019) Pembrolizumab-induced immune-mediated colitis in a patient with concurrent *Clostridium difficile* infection . *Case Rep Oncol* 12 (1): 164 – 170 . doi: 10.1159/000497155 . 31043955 PMC6477466

37. Gentile NM , D'Souza A , Fujii LL , Wu TT ,

Murray JA (2013) Association between ipilimumab and celiac disease . Mayo Clin Proc 88 (4): 414 – 417 . doi: 10.1016/j.mayocp.2013.01.015 . 23541015 38. Abdel-Wahab N , Shah M , Suarez-Almazor ME (2016) Adverse events associated with immune checkpoint blockade in patients with cancer: a systematic review of case reports . PLoS One 11 (7): e0160221 . doi: 10.1371/journal.pone.0160221 . 27472273 PMC4966895 39. Abu-Sbeih H , Ali FS , Alsaadi D , Jennings J , Luo W , Gong Z , Richards DM , Charabaty A , Wang Y (2018) Outcomes of vedolizumab therapy in patients with immune checkpoint inhibitor-induced colitis: a multi-center study . J Immunother Cancer 6 (1): 142 . doi: 10.1186/s40425-018-0461-4 . 30518410 PMC6280383 40. Bergqvist V , Hertervig E , Gedeon P , Kopljär M , Griph H , Kinhult S , Carneiro A , Marsal J (2017) Vedolizumab treatment for immune checkpoint inhibitor-induced enterocolitis . Cancer Immunol Immunother 66 (5): 581 – 592 . doi: 10.1007/s00262-017-1962-6 . 28204866 PMC5406433 41. Hsieh AH , Ferman M , Brown MP , Andrews JM (2016) Vedolizumab: a novel treatment for ipilimumab-induced colitis . BMJ Case Rep 2016 : 216641 . doi: 10.1136/bcr-2016-216641 . PMC5015180 27539137 42. Arriola E , Wheeler M , Karydis I , Thomas G , Ottensmeier C (2015) Infliximab for IPILIMUMAB-related colitis-letter . Clin Cancer Res 21 (24): 5642 – 5643 . doi: 10.1158/1078-0432.CCR-15-2471 . 26672088 43. Johnson DH , Zobniw CM , Trinh VA , Ma J , Bassett RL Jr. , Abdel-Wahab N , Anderson J , Davis JE , Joseph J , Uemura M , Noman A , Abu-Sbeih H , Yee C , Amaria R , Patel S , Tawbi H , Glitza IC , Davies MA , Wong MK , Woodman S , Hwu WJ , Hwu P , Wang Y , Diab A (2018) Infliximab associated with faster symptom resolution compared with corticosteroids alone for the management of immune-related enterocolitis . J Immunother Cancer 6 (1): 103 . doi: 10.1186/s40425-018-0412-0 . 30305177 PMC6180568 44. Abu-Sbeih H , Ali FS , Wang X , Mallepally N , Chen E , Altan M , Bresalier RS , Charabaty A , Dadu R , Jazaeri A , Lashner B , Wang Y (2019) Early introduction of selective immunosuppressive therapy associated with favorable clinical outcomes in patients with immune checkpoint inhibitor-induced colitis . J Immunother Cancer 7 (1): 93 . doi: 10.1186/s40425-019-0577-1 . 30940209 PMC6444537 45. De Martin E , Michot JM , Papouin B , Champiat S , Mateus C , Lambotte O , Roche B , Antonini TM , Coilly A , Laghouati S , Robert C , Marabelle A , Guettier C , Samuel D (2018) Characterization of liver injury induced by cancer immunotherapy using immune checkpoint inhibitors . J Hepatol 68 (6): 1181 – 1190 . doi: 10.1016/j.jhep.2018.01.033 . 29427729 46. Bernardo SG , Moskalenko M , Pan M , Shah S , Sidhu HK , Sicular S , Harcharik S , Chang R , Friedlander P , Saenger YM (2013) Elevated rates of transaminitis during ipilimumab therapy for metastatic melanoma . Melanoma Res 23 (1): 47 – 54 . doi: 10.1097/CMR.0b013e32835c7e68 . 23262440 47. Paz-Ares L , Luft A , Vicente D , Tafreshi A , Güüm M , Mazières J , Hermes B , Çay ■enler F , Cs■sz T , Fülöp A , Rodríguez-Cid J , Wilson J , Sugawara S , Kato T , Lee KH , Cheng Y , Novello S , Halmos B , Li X , Lubiniecki GM , Piperdi B , Kowalski DM ; KEYNOTE-407 Investigators (2018) Pembrolizumab plus chemotherapy for squamous non-small-cell lung cancer . N Engl J Med 379 (21): 2040 – 2051 . doi: 10.1056/NEJMoa1810865 . 30280635 48. Schmid P , Adams S , Rugo HS , Schneeweiss A , Barrios CH , Iwata H , Diéras V , Hegg R , Im SA , Shaw Wright G , Henschel V , Molinero L , Chui SY , Funke R , Husain A , Winer EP , Loi S , Emens LA ; Impassion130 Trial Investigators (2018) Atezolizumab and nab-paclitaxel in advanced triple-negative breast cancer . N Engl J Med ; 379 (22): 2108 – 2121 . doi: 10.1056/NEJMoa1809615 . 30345906 49. Rini BI , Plimack ER , Stus V , Gafanov R , Hawkins R , Nosov D , Pouliot F , Alekseev B , Soulières D , Melichar B , Vynnychenko I , Kryzhanivska A , Bondarenko I , Azevedo SJ , Borchellini D , Szczylik C , Markus M , McDermott RS , Bedke J , Tartas S , Chang YH , Tamada S , Shou Q , Perini RF , Chen M , Atkins MB , Powles T ; KEYNOTE-426 Investigators (2019) Pembrolizumab plus axitinib versus sunitinib for advanced renal-cell carcinoma . N Engl J Med 380 (12): 1116 – 1127 . doi: 10.1056/NEJMoa1816714 30779529 50. Pelster MS , Amaria RN (2019) Combined targeted therapy and immunotherapy in melanoma: a review of the impact on the tumor microenvironment and outcomes of early clinical trials . Ther Adv Med Oncol 11 : 1758835919830826 . doi: 10.1177/1758835919830826 . 30815041 PMC6384439 51. Everett J , Srivastava A , Misdraji J (2017) Fibrin ring granulomas in checkpoint inhibitor-induced hepatitis . Am J Surg Pathol 41 (1): 134 – 137 . doi: 10.1097/PAS.0000000000000759 27792061 52. Johncilla M , Misdraji J , Pratt DS , Agoston AT , Lauwers GY , Srivastava A , Doyle LA (2015) Ipilimumab-associated hepatitis: clinicopathologic characterization in a series of 11 cases . Am J Surg Pathol 39 (8): 1075 – 1084 . doi: 10.1097/PAS.0000000000000453 . 26034866 53. Zen Y , Yeh MM (2018) Hepatotoxicity of immune checkpoint inhibitors: a histology study of seven cases in comparison with autoimmune

hepatitis and idiosyncratic drug-induced liver injury . *Mod Pathol* 31 (6): 965 – 973 . doi: 10.1038/s41379-018-0013-y . 29403081 54. Kleiner DE , Berman D (2012) Pathologic changes in ipilimumab-related hepatitis in patients with metastatic melanoma . *Dig Dis Sci* 57 (8): 2233 – 2240 . doi: 10.1007/s10620-012-2140-5 . 22434096 PMC3792485 55. Kim KW , Ramaiya NH , Krajewski KM , Jagannathan JP , Tirumani SH , Srivastava A , Ibrahim N (2013) Ipilimumab associated hepatitis: imaging and clinicopathologic findings . *Invest New Drugs* 31 (4): 1071 – 1077 . doi: 10.1007/s10637-013-9939-6 . 23408334 56. Spänkuch I , Gassenmaier M , Tampouri I , Noor S , Forschner A , Garbe C , Amaral T (2017) Severe hepatitis under combined immunotherapy: Resolution under corticosteroids plus anti-thymocyte immunoglobulins . *Eur J Cancer* 81 : 203 – 205 . doi: 10.1016/j.ejca.2017.05.018 . 28641200 57. Zhang HC , Luo W , Wang Y (2019) Acute liver injury in the context of immune checkpoint inhibitor-related colitis treated with infliximab . *J Immunother Cancer* 7 (1): 47 . doi: 10.1186/s40425-019-0532-1 . 30777137 PMC6380028 58. Weber J , Thompson JA , Hamid O , Minor D , Amin A , Ron I , Ridolfi R , Assi H , Maraveyas A , Berman D , Siegel J , O'Day SJ (2009) A randomized, double-blind, placebo-controlled, phase II study comparing the tolerability and efficacy of ipilimumab administered with or without prophylactic budesonide in patients with unresectable stage III or IV melanoma . *Clin Cancer Res* 15 (17): 5591 – 5598 . doi: 10.1158/1078-0432.CCR-09-1024 . 19671877 59. Zhang ML , Neyaz A , Patil D , Chen J , Dougan M , Deshpande V (2020) Immune-related adverse events in the gastrointestinal tract: diagnostic utility of upper gastrointestinal biopsies . *Histopathology* 76 (2): 233 – 243 . doi: 10.1111/his.13963 . 31361907 PMC8386137 60. Nemakayala DR , Cash BD (2019) Excluding irritable bowel syndrome in the inflammatory bowel disease patient: how far to go? *Curr Opin Gastroenterol* 35 (1): 58 – 62 . doi: 10.1097/MOG.0000000000000493 . 30407259 61. Mooradian MJ , Wang DY , Coromilas A , Lumish M , Chen T , Giobbie-Hurder A , Johnson DB , Sullivan RJ , Dougan M (2020) Mucosal inflammation predicts response to systemic steroids in immune checkpoint inhibitor colitis . *J Immunother Cancer* 8 (1): e000451 . doi: 10.1136/jitc-2019-000451 . 32414860 PMC7239692 62. Hughes MS , Molina GE , Chen ST , Zheng H , Deshpande V , Fadden R , Sullivan RJ , Dougan M (2019) Budesonide treatment for microscopic colitis from immune checkpoint inhibitors . *J Immunother Cancer* 7 (1): 292 . doi: 10.1186/s40425-019-0756-0 . 31699151 PMC6839080 63. Hillock NT , Heard S , Kichenadasse G , Hill CL , Andrews J (2017) Infliximab for ipilimumab-induced colitis: A series of 13 patients . *Asia Pac J Clin Oncol* 13 (5): e284 – e290 . doi: 10.1111/ajco.12651 . 27981760 64. Panagiotou I , Brontzos EN , Bafaloukos D , Stoupis C , Brestas P , Kelekis DA (2002) Malignant melanoma metastatic to the gastrointestinal tract . *Melanoma Res* 12 (2): 169 – 173 . doi: 10.1097/00008390-200204000-00010 11930114 65. Bertrand F , Montfort A , Marcheteau E , Imbert C , Gilhodes J , Filleron T , Rochaix P , Andrieu-Abadie N , Levade T , Meyer N , Colacios C , Ségui B (2017) TNF α blockade overcomes resistance to anti-PD-1 in experimental melanoma . *Nat Commun* 8 (1): 2256 . doi: 10.1038/s41467-017-02358-7 . 29273790 PMC5741628 66. Perez-Ruiz E , Minute L , Otano I , Alvarez M , Ochoa MC , Belsue V , de Andrea C , Rodriguez-Ruiz ME , Perez-Gracia JL , Marquez-Rodas I , Llacer C , Alvarez M , de Luque V , Molina C , Teijeira A , Berraondo P , Melero I (2019) Prophylactic TNF blockade uncouples efficacy and toxicity in dual CTLA-4 and PD-1 immunotherapy . *Nature* 569 (7756): 428 – 432 . doi: 10.1038/s41586-019-1162-y . 31043740 67. Maneiro JR , Souto A , Gomez-Reino JJ (2017) Risks of malignancies related to tofacitinib and biological drugs in rheumatoid arthritis: Systematic review, meta-analysis, and network meta-analysis . *Semin Arthritis Rheum* 47 (2): 149 – 156 . doi: 10.1016/j.semarthrit.2017.02.007 . 28284845 68. Chen Y , Sun J , Yang Y , Huang Y , Liu G (2016) Malignancy risk of anti-tumor necrosis factor alpha blockers: an overview of systematic reviews and meta-analyses . *Clin Rheumatol* 35 (1): 1 – 18 . doi: 10.1007/s10067-015-3115-7 . 26573205 69. Wang Y , Wiesnoski DH , Helmink BA , Gopalakrishnan V , Choi K , DuPont HL , Jiang ZD , Abu-Sbeih H , Sanchez CA , Chang CC , Parra ER , Francisco-Cruz A , Raju GS , Stroehlein JR , Campbell MT , Gao J , Subudhi SK , Maru DM , Blando JM , Lazar AJ , Allison JP , Sharma P , Tetzlaff MT , Wargo JA , Jenq RR (2019) Author Correction: Fecal microbiota transplantation for refractory immune checkpoint inhibitor-associated colitis . *Nat Med* 25 (1): 188 . doi: 10.1038/s41591-018-0305-2 . 30479380 70. Costello SP , Soo W , Bryant RV , Jairath V , Hart AL , Andrews JM (2017) Systematic review with meta-analysis: faecal microbiota transplantation for the induction of remission for active ulcerative colitis . *Aliment Pharmacol Ther* 46 (3): 213 – 224 . doi: 10.1111/apt.14173 . 28612983 71. Quraishi MN , Widlak M , Bhala N , Moore D , Price M , Sharma N , Iqbal TH (2017) Systematic review with meta-analysis: the efficacy of faecal microbiota

transplantation for the treatment of recurrent and refractory *Clostridium difficile* infection . *Aliment Pharmacol Ther* 46 (5): 479 – 493 . doi: 10.1111/apt.14201 . 28707337 72. Feagan BG , Sandborn WJ , Gasink C , Jacobstein D , Lang Y , Friedman JR , Blank MA , Johanss J , Gao LL , Miao Y , Adedokun OJ , Sands BE , Hanauer SB , Vermeire S , Targan S , Ghosh S , de Villiers WJ , Colombel JF , Tulassay Z , Seidler U , Salzberg BA , Desreumaux P , Lee SD , Loftus EV Jr , Dieleman LA , Katz S , Rutgeerts P ; UNITI–IM–UNITI Study Group (2016) Ustekinumab as induction and maintenance therapy for Crohn's disease . *N Engl J Med* 375 (20): 1946 – 1960 . doi: 10.1056/NEJMoa1602773 27959607 73. Wei SC , Levine JH , Cogdill AP , Zhao Y , Anang NAS , Andrews MC , Sharma P , Wang J , Wargo JA , Pe'er D , Allison JP (2017) Distinct cellular mechanisms underlie anti-CTLA-4 and anti-PD-1 checkpoint blockade . *Cell* 170 (6): 1120 – 1133 e17 . doi: 10.1016/j.cell.2017.07.024 . 28803728 PMC5591072 74. Zaretsky JM , Garcia-Diaz A , Shin DS , Escuin-Ordinas H , Hugo W , Hu-Lieskovan S , Torrejon DY , Abril-Rodriguez G , Sandoval S , Barthly L , Saco J , Homet Moreno B , Mezzadra R , Chmielowski B , Ruchalski K , Shintaku IP , Sanchez PJ , Puig-Saus C , Cherry G , Seja E , Kong X , Pang J , Berent-Maoz B , Comin-Anduix B , Graeber TG , Tumeh PC , Schumacher TN , Lo RS , Ribas A (2016) Mutations associated with acquired resistance to PD-1 blockade in melanoma . *N Engl J Med* 375 (9): 819 – 829 . doi: 10.1056/NEJMoa1604958 . 27433843 PMC5007206 75. Hugo W , Zaretsky JM , Sun L , Song C , Moreno BH , Hu-Lieskovan S , Berent-Maoz B , Pang J , Chmielowski B , Cherry G , Seja E , Lomeli S , Kong X , Kelley MC , Sosman JA , Johnson DB , Ribas A , Lo RS (2016) Genomic and transcriptomic features of response to anti-PD-1 therapy in metastatic melanoma . *Cell* 165 (1): 35 – 44 . doi: 10.1016/j.cell.2016.02.065 . 26997480 PMC4808437 76. Rooney MS , Shukla SA , Wu CJ , Getz G , Hacohen N (2015) Molecular and genetic properties of tumors associated with local immune cytolytic activity . *Cell* 160 (1–2): 48 – 61 . doi: 10.1016/j.cell.2014.12.033 . 25594174 PMC4856474 77. Zeissig S , Petersen BS , Tomczak M , Melum E , Huc-Claustre E , Dougan SK , Laerdahl JK , Stade B , Forster M , Schreiber S , Weir D , Leichtner AM , Franke A , Blumberg RS (2015) Early-onset Crohn's disease and autoimmunity associated with a variant in CTLA-4 . *Gut* 64 (12): 1889 – 1897 . doi: 10.1136/gutjnl-2014-308541 . 25367873 PMC4512923 78. Schubert D , Bode C , Kenefeck R , Hou TZ , Wing JB , Kennedy A , Bulashevskaya A , Petersen BS , Schäffer AA , Grüning BA , Unger S , Frede N , Baumann U , Witte T , Schmidt RE , Dueckers G , Niehues T , Seneviratne S , Kanariou M , Speckmann C , Ehl S , Rensing-Ehl A , Warnatz K , Rakhmanov M , Thimme R , Hasselblatt P , Emmerich F , Cathomen T , Backofen R , Fisch P , Seidl M , May A , Schmitt-Graeff A , Ikemizu S , Salzer U , Franke A , Sakaguchi S , Walker LSK , Sansom DM , Grimbacher B (2014) Autosomal dominant immune dysregulation syndrome in humans with CTLA4 mutations . *Nat Med* 20 (12): 1410141 – 6 . doi: 10.1038/nm.3746 . PMC4668597 25329329 79. Kuehn HS , Ouyang W , Lo B , Deenick EK , Niemela JE , Avery DT , Schickel JN , Tran DQ , Stoddard J , Zhang Y , Frucht DM , Dumitriu B , Scheinberg P , Folio LR , Frein CA , Price S , Koh C , Heller T , Seroogy CM , Huttenlocher A , Rao VK , Su HC , Kleiner D , Notarangelo LD , Rampertap Y , Olivier KN , McElwee J , Hughes J , Pittaluga S , Oliveira JB , Meffre E , Fleisher TA , Holland SM , Lenardo MJ , Tangye SG , Uzel G (2014) Immune dysregulation in human subjects with heterozygous germline mutations in CTLA4 . *Science* 345 (6204): 1623 – 1627 . doi: 10.1126/science.1255904 . 25213377 PMC4371526 80. Tivol EA , Boyd SD , McKeon S , Borriello F , Nickerson P , Strom TB , Sharpe AH (1997) CTLA4Ig prevents lymphoproliferation and fatal multiorgan tissue destruction in CTLA-4-deficient mice . *J Immunol* 158 (11): 5091 – 5094 . 9164923 81. Schoenfeld SR , Aronow ME , Leaf RK , Dougan M , Reynolds KL (2020) Diagnosis and management of rare immune-related adverse events . *Oncologist* 25 (1): 6 – 14 . doi: 10.1634/theoncologist.2019-0083 . 31694890 PMC6964128 Table 1. Colitis Grade Based Table.

Abbreviations: TTG-IgA (tissue transglutaminase-immunoglobulin A), IgA (immunoglobulin A), HepB (Hepatitis B Virus), sAg (surface antigen), sAb (surface antibody), cAb (core antibody), IHC (immunohistochemistry)

Routine monitoring of comprehensive metabolic panels, complete blood counts, and TSH are recommended for all patients undergoing immunotherapy

Management recommendations for hepatic adverse events.

ALT: alanine aminotransferase; AST: aspartate aminotransferase; ALKP: alkaline phosphatase; TBILI: total bilirubin; DBILI: direct bilirubin; IgM: immunoglobulin M; IgG: immunoglobulin G; Epstein-Barr Virus (EBV); cytomegalovirus (CMV); ANA: antinuclear antibody; ASMA: anti-smooth muscle antibody; AMA: anti-mitochondrial antibody; TIBC: total iron binding capacity; A1AT: alpha-1 anti-trypsin; MRCP: magnetic resonance cholangiopancreatography; EUS: endoscopic ultrasound.