

Pan-Asian adaptation of the EHNS-ESMO-ESTRO Clinical Practice Guidelines for the diagnosis, treatment and follow-up of patients with squamous cell carcinoma of the head and neck.

Title: Pan-Asian adaptation of the EHNS-ESMO-ESTRO Clinical Practice Guidelines for the diagnosis, treatment and follow-up of patients with squamous cell carcinoma of the head and neck.

Publication year: 2021

Journal: ESMO open

Authors: Department of Internal Medicine, Seoul National University Hospital, Seoul, Korea.

Electronic address: bhumsuk@snu.ac.kr.

Society: European Society for Medical Oncology

Origin country: International (Europe)

Target countries: Europe

DOI: 10.1016/j.esmoop.2021.100309

PMID: 34844180

Source: <https://pubmed.ncbi.nlm.nih.gov/34844180/>

pmc ESMO Open ESMO Open 3084 esmo 101690685 ESMO Open 2059-7029 Elsevier
PMC8710460 PMC8710460.1 8710460 8710460 34844180 10.1016/j.esmoop.2021.100309
S2059-7029(21)00271-4 100309 1 Special Article Pan-Asian adaptation of the
EHNS–ESMO–ESTRO Clinical Practice Guidelines for the diagnosis, treatment and follow-up of
patients with squamous cell carcinoma of the head and neck Keam B. bhumsuk@snu.ac.kr 1 *
Machiels J.-P. 2 Kim H.R. 3 Licitra L. 4 Golusinski W. 5 Gregoire V. 6 Lee Y.G. 7 Belka C. 8 Guo Y.
9 Rajappa S.J. 10 Tahara M. 11 Azrif M. 12 Ang M.K. 13 Yang M.-H. 14 Wang C.-H. 15 Ng Q.S. 16
Wan Zamaniah W.I. 17 Kiyota N. 18 Babu S. 19 Yang K. 20 Curigliano G. 21 22 Peters S. 23 Kim
T.W. 24 Yoshino T. 25 Pentheroudakis G. 26 1 Department of Internal Medicine, Seoul National
University Hospital, Seoul, Korea 2 Service d'Oncologie Médicale, Institut Roi Albert II, Cliniques
Universitaires Saint-Luc, Brussels, Belgium 3 Division of Medical Oncology, Department of Internal
Medicine, Yonsei Cancer Center, Yonsei University College of Medicine, Seoul, South Korea 4
Head and Neck Cancer Medical Oncology Department, Fondazione IRCCS Istituto Nazionale dei
Tumori and University of Milan, Milan, Italy 5 Department of Head and Neck Surgery, Poznan
University of Medical Sciences, The Greater Poland Cancer Centre, Poznan, Poland 6 Department
of Radiation Oncology, Centre Léon Bérard, Lyon, France 7 Division of Hematology and Medical
Oncology, Department of Internal Medicine, Kangbuk Samsung Hospital, Sungkyunkwan University
School of Medicine, Seoul, South Korea 8 Department of Radiation Oncology, LMU Hospital,
Munich, Germany 9 Department of Medical Oncology, Shanghai East Hospital, Tongji University
School of Medicine, Shanghai, China 10 Medical Oncology, Basavataarakam Indo American Cancer
Hospital and Research Institute, Hyderabad, India 11 Department of Head and Neck Medical
Oncology, National Cancer Center Hospital East, Chiba, Japan 12 Clinical Oncology, Prince Court
Medical Centre, Kuala Lumpur, Malaysia 13 Department of Medical Oncology, National Cancer
Centre, Singapore, Singapore 14 Institute of Clinical Medicine, National Yang Ming Chiao Tung
University, Taipei, Taiwan 15 Division of Hemato-oncology, Department of Internal Medicine,
Keelung Chang Gung Memorial Hospital and Chang Gung University College of Medicine,
Taoyuan, Taiwan 16 Division of Medical Oncology, National Cancer Centre, Singapore, Singapore
17 Clinical Oncology Unit, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia 18
Oncology/Hematology, Cancer Center, Kobe University Hospital, Kobe, Japan 19 Department of
Medical Oncology, Kidwai Memorial Institute of Oncology, Bangalore, India 20 Department of
Clinical Oncology, Union Hospital, Tongji Medical College, Huazhong University of Science and
Technology, Wuhan, China 21 Department of Oncology and Hemato-Oncology, University of Milan,
Milan, Italy 22 Istituto Europeo di Oncologia, IRCCS, Milano, Italy 23 Department of Oncology,
Centre Hospitalier Universitaire Vaudois (CHUV), Lausanne, Switzerland 24 Department of

Oncology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea 25
 Department of Gastroenterology and Gastrointestinal Oncology, National Cancer Center East,
 Chiba, Japan 26 ESMO, Lugano, Switzerland * Correspondence to: Prof. Bhumsuk Keam,
 Department of Internal Medicine, Seoul National University Hospital, 101 Daehak-ro, Jongno-gu,
 Seoul 03080, Korea. Tel: +82-2-2072-7215; Fax: +82-2-2072-7379 bhumsuk@snu.ac.kr 12 2021
 26 11 2021 6 6 391737 100309 27 9 2021 26 10 2021 27 10 2021 26 11 2021 04 01 2022 05 01
 2022 © 2021 The Authors 2021 <https://creativecommons.org/licenses/by-nc-nd/4.0/> This is an open
 access article under the CC BY-NC-ND license
 (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). The most recent version of the European
 Society for Medical Oncology (ESMO) Clinical Practice Guidelines for the diagnosis, treatment and
 follow-up of squamous cell carcinoma (SCC) of the oral cavity, larynx, oropharynx and hypopharynx
 was published in 2020. It was therefore decided by both the ESMO and the Korean Society of
 Medical Oncology (KSMO) to convene a special, virtual guidelines meeting in July 2021 to adapt
 the ESMO 2020 guidelines to consider the potential ethnic differences associated with the
 treatment of SCCs of the head and neck (SCCHN) in Asian patients. These guidelines represent
 the consensus opinions reached by experts in the treatment of patients with SCCHN (excluding
 nasopharyngeal carcinomas) representing the oncological societies of Korea (KSMO), China
 (CSCO), India (ISMPO), Japan (JSMO), Malaysia (MOS), Singapore (SSO) and Taiwan (TOS). The
 voting was based on scientific evidence and was independent of the current treatment practices
 and drug access restrictions in the different Asian countries. The latter was discussed when
 appropriate. This manuscript provides a series of expert recommendations (Clinical Practice
 Guidelines) which can be used to provide guidance to health care providers and clinicians for the
 optimisation of the diagnosis, treatment and management of patients with SCC of the oral cavity,
 larynx, oropharynx and hypopharynx across Asia.

- This article provides ESMO expert recommendations adapted for the treatment of SCCHN (excluding nasopharyngeal carcinomas) in Asian patients.

The aim was to provide guidance for the optimisation of the management of such patients across Asia.

The availability and applicability of certain procedures as they relate to certain of the recommendations are discussed.

In 2020 an estimated 19.3 million new cases of cancer were diagnosed and almost 10 million cancer-related deaths recorded, worldwide. 1 Of these, squamous cell carcinomas of the head and neck (SCCHN) including carcinomas of the lip, oral cavity, larynx, oropharynx and hypopharynx, but excluding nasopharyngeal cancer, accounted for 3.9% (744 994) of new cases across both sexes, and 3.7% (357.339) of cancer deaths. 1 The highest incidence of head and neck cancer (HNC) is seen in Asia, with deaths from HNCs accounting for >5% of all cancer deaths. HNC comprises a heterogenous group of malignancies and the prevalences of the different types of HNCs vary from country to country across Asia. For example China and Taiwan have high incidences of oral cavity cancers, with the latter having increasing rates of human papilloma virus (HPV)-related oropharyngeal cancer. 2, 3 India currently contributes ~60% of the HNC cases worldwide with HNC the most common cancer in Indian men, and cancer of the oral cavity the most prevalent. In South Korea, laryngeal cancer is the commonest form of HNC with the incidence of all HNCs (tonsil, hypopharynx, oropharynx and larynx cancer) increasing, with a higher incidence in men than in women. 4 In Japan alone, there are >39 000 cases of HNC and ~10 000 deaths from the disease, annually. 5, 6 A report from the Japan Society for Head and Neck Cancer Registry Committee on 11 716 previously untreated HNC patients registered in 2016 showed ~83% of the cases to be accounted for by tumours of the oral cavity (24.9%), larynx (20.4%), hypopharynx (21.4%) and oropharynx (16.9%). 7

Although the incidence varies between countries and individual regions within countries, ~80% of cases of SCCHN worldwide are attributable to tobacco use, excessive alcohol consumption or both, and in South Asia the use of smokeless tobacco and betel quid/areca nut products. 8 Betel

quid/areca nut use has been linked to high rates of cancers of the oral cavity in India, Taiwan and some provinces of mainland China. 9 Other risk factors include environmental pollutants, especially in countries with worsening air pollution such as India 10 and China, 11 and HPV infections in the aetiology of oropharyngeal cancer and cancers of the oral cavity. 12 A recent meta-analysis looking at the prevalence of HPV-related oropharyngeal cancers in the Asia Pacific region reported an overall prevalence of 40.53% for oropharyngeal cancers. 13 In a Malaysian study of 60 patients with oropharyngeal cancer of whom 53.3% were of Chinese ethnicity, 35% of Indian ethnicity and 11.7% of Malay ethnicity, all the Indian patients had p16-negative disease. 14 This was consistent with a study of 88 patients with SCCHN conducted in South India in which only 2.6% cases were HPV/p16-positive. 15 Significantly, in the Malaysian study 14 80% of the HPV-positive cases were in Chinese patients and the prevalence of p16-positive oropharyngeal squamous cell carcinomas (SCC) across all three ethnicities was half that of a matched UK cohort (25% versus 49%). 14 A study conducted in Singapore involving 159 urban, multiethnic, South East Asian patients with SCCHN demonstrated a high prevalence of high-risk HPV variants (HPV16, 18, 31, 45, 56 and 68) and confirmed that HPV16 and p16 immunohistochemical expression were predominantly detected in the oropharyngeal carcinomas. 16 Although an ethnic predisposition cannot be excluded, the differences in HPV positivity between the different Asian populations is probably due to differences in sexual practices.

Guidelines and recommendations for the treatment and management of patients with SCCHN in Asia have been published for India, 17 , 18 China, 19 Japan, 20 Malaysia 21 and Taiwan 22 and are important for the standardisation of both diagnostic and treatment approaches, with the aim of optimising clinical outcomes for what is an increasing health care problem in Asia. The European Society for Medical Oncology (ESMO) guidelines for the diagnosis treatment and follow-up of patients with SCCs of the oral cavity, larynx, oropharynx and hypopharynx (excluding nasopharyngeal carcinomas), prepared in conjunction with the European Head and Neck Society (EHNS) and the European Society for Radiotherapy and Oncology (ESTRO), were published in October 2020, 23 and a decision was taken by ESMO and the Korean Society of Medical Oncology (KSMO) that these guidelines should be adapted for patients of Asian ethnicity. Consequently, representatives of KSMO, ESMO, ESTRO, the Chinese Society of Clinical Oncology (CSCO), the Indian Society of Medical and Paediatric Oncology (ISMPO), the Japanese Society of Medical Oncology (JSMO), the Malaysian Oncological Society (MOS), the Singapore Society of Oncology (SSO) and the Taiwan Oncology Society (TOS) convened for a virtual working meeting ('face-to-face' meeting) on 24 July 2021, hosted by KSMO, to adapt the recent EHNS-ESMO-ESTRO Clinical Practice Guidelines. 23 This manuscript summarises the Pan-Asian adapted guidelines developed at the meeting accompanied by the level of evidence (LoE), grade of recommendation (GoR) and percentage consensus reached for each recommendation.

This Pan-Asian adaptation of the current ESMO Clinical Practice Guidelines 23 was prepared in accordance with the principles of ESMO standard operating procedures (<http://www.esmo.org/Guidelines/ESMO-Guidelines-Methodology>) and was a KSMO-ESMO initiative endorsed by CSCO, ISMPO, JSMO, MOS, SSO and TOS.

Representatives from KSMO (n = 4), ESMO (n = 7), EHNS (n = 1), ESTRO (n = 2) and two experts from each of the oncological societies of China (CSCO), India (ISMPO), Japan (JSMO), Malaysia (MOS), Singapore (SSO) and Taiwan (TOS) convened for the virtual 'face-to-face' meeting. Only two of the members from KSMO (HRK and YGL) were allowed to vote on the recommendations together with the experts from each of the six other Asian oncology societies (n = 14).

A modified Delphi process was used to review, accept or adapt each of the individual recommendations in the latest EHNS-ESMO-ESTRO Clinical Practice Guidelines. 23 The 14 Asian experts were asked to vote YES or NO (one vote per society) on the 'acceptability' (agreement with the scientific content of the recommendation) and 'applicability' (availability, reimbursement and practical challenges) of each of the ESMO recommendations in a pre-meeting survey (see Supplementary Methodology , available at <https://doi.org/10.1016/j.esmoop.2021.100309>). For

recommendations, where a consensus was not reached, the Asian experts were invited to modify the wording of the recommendation(s) at the virtual 'face-to-face' meeting using rounds of voting in order to determine the definitive acceptance or rejection of an adapted recommendation and discuss the applicability challenges. The 'Infectious Diseases Society of America-United States Public Health Service Grading System' (Supplementary Table S1 , available at <https://doi.org/10.1016/j.esmoop.2021.100309>) 24 was used to define the LoE and strength (grade) of each recommendation. Any modifications to the initial recommendations were highlighted in bold text in a summary table of the final Asian recommendations and in the main text, if and as applicable. A consensus was considered to have been achieved when $\geq 80\%$ of experts voted that a recommendation was acceptable.

In the initial pre-meeting survey, the 14 Asian experts reported on the 'acceptability' and 'applicability' of the 32 recommendations for the diagnosis treatment and follow-up of patients with SCCHN of the oral cavity, larynx, oropharynx and hypopharynx from the 2020 EHNS-ESMO-ESTRO Clinical Practice Guidelines. 23 These recommendations were made in the four categories listed below: • Diagnosis and pathology/molecular biology (Recommendations 1a-f)

Staging and risk assessment (Recommendation 2)

A lack of agreement in the pre-meeting survey was initially established for 'recommendations 3i and 3v' (with no consensus for 'acceptability') (Supplementary Table S2 , available at <https://doi.org/10.1016/j.esmoop.2021.100309>) and 'recommendations 1c, 1d, 1f, 3b, 3i, 3m, 3n, 3s, 3t, 3u, 3v and 4c' (with no consensus for 'applicability') (Supplementary Table S3 , available at <https://doi.org/10.1016/j.esmoop.2021.100309>). After further consideration, however, the 'recommendations 3i and 3v', and 'recommendations 1d, 3a, 3e, 3f, 3k, 3l 3m, 3n, 3r, 3s, 3t and 4c' were identified for discussion during the 'face-to-face' meeting, based on the comments and feedback from the initial pre-meeting survey, as it was clear that some of the comments made in terms of applicability had scientific relevance. It was also decided that there was no need to discuss 'recommendations 1c, 1f, 3b and 3u' initially identified for discussion in terms of applicability.

The Pan-Asian panel of experts agreed with and 'accepted' completely (100% consensus) the EHNS-ESMO-ESTRO recommendations on screening, 'recommendations 1a-f' from a scientific point of view (see below and Table 1). 1a. Clinical examination and pathological confirmation are mandatory [IV, A].

Rigid head and neck endoscopy, head and neck contrast-enhanced computed tomography (CE-CT) and/or magnetic resonance imaging (MRI) and chest imaging (with CT and/or [18F]2-fluoro-2-deoxy- d -glucose-positron emission tomography (FDG-PET) are strongly recommended [IV, A].

For oropharyngeal cancer, p16 immunohistochemistry (IHC) is strongly recommended [I, A].

For SCCHN of unknown primary, p16 and Epstein–Barr-encoded RNA (EBER) are recommended. If p16 staining is positive, another specific HPV test may be carried out to confirm the HPV status [III, B].

On the surgical specimens, depth of invasion of oral cavity cancer, assessment of the number of invaded lymph nodes as well as the presence extracapsular extension, perineural and lymphatic infiltration and the surgical margins must be evaluated [I, A].

For recurrent and/or metastatic SCCHN, tumour programmed death-ligand 1 (PD-L1) expression should be evaluated [II, B].

CE-CT, contrast-enhanced-computed tomography; CPS, combined positive score; CRT, chemoradiotherapy; CT, computed tomography; DOI, depth of invasion; DPD, dihydropyrimidine

dehydrogenase; EMA, European Medicines Agency; EBER, Epstein–Barr-encoded RNA; ESMO-MCBS, ESMO-Magnitude of Clinical Benefit Scale; FDA, Food and Drug Administration; FDG-PET, 2-fluoro-2-deoxy- d -glucose-positron emission tomography; 5-FU, 5-fluorouracil; HPV, human papilloma virus; IHC, immunohistochemistry; IMRT, intensity modulated radiotherapy ; MDT, multidisciplinary team; MRI, magnetic resonance imaging; PCE, paclitaxel, carboplatin and cetuximab; PD-L1, programmed death-ligand 1; R/M, recurrent/metastatic; RT, radiotherapy; SCCHN, squamous cell carcinoma of the head and neck; TNM, tumour, node and metastasis; TPeX, taxane, cisplatin and cetuximab; TPF, docetaxel, cisplatin and 5 -FU; UICC, Union for International Cancer Control; VMAT, volumetric modulated arc therapy.

The experts from the oncological societies of Japan, Singapore and China, however, did not agree that an HPV test had to be carried out (Supplementary Table S3 , available at <https://doi.org/10.1016/j.esmoop.2021.100309>), and in the wording of 'recommendation 1d' above 'should' was replaced by ' may ' as per bold text above and Table 1 , and the GoR changed from A to B.

The Pan-Asian panel of experts agreed with and 'accepted' completely (100% consensus) the EHNS-ESMO-ESTRO recommendations on staging and risk assessment, 'recommendation 2a' (below and Table 1) from both a scientific (acceptability) and applicability point of view (see also Supplementary Table S2 , available at <https://doi.org/10.1016/j.esmoop.2021.100309>). 2a. The UICC TNM 8th edition staging system should be used (Supplementary Table S4 , available at <https://doi.org/10.1016/j.esmoop.2021.100309>). 25

The Pan-Asian panel of experts agreed with and 'accepted' completely (100% consensus) the EHNS-ESMO-ESTRO recommendations on treatment, 'recommendations 3b-d, g, h, j, o-q and u,' from a scientific point of view (see below and Table 1). A lack of consensus in terms of 'acceptability' for recommendations 3i and 3v was identified, however, from the time of the pre-meeting survey and the need for discussion for 'recommendations 3a, e, f, k, l, m, n, r, s and t' from a scientific perspective only after consideration of the feedback comments (Supplementary Table S5 , available at <https://doi.org/10.1016/j.esmoop.2021.100309>), as described above.

With regard to 'recommendation 3a' there was concern that not all centres in Asia have multidisciplinary teams (MDT) for the treatment of HNC, so the wording of the recommendation was revised slightly as per the bold text below with 100% consensus. After discussion at the 'face-to-face' meeting 'recommendations 3e and f' were accepted without change, with 100% consensus. 3a. Ideally the optimal treatment strategy should be discussed in an MDT including not only the treating physicians but all the supportive specialities [III, A; consensus = 100%].

Patients should be treated at high-volume facilities [II, A].

In the case of radiotherapy (RT), all patients should be treated by intensity modulated radiotherapy (IMRT) or volumetric modulated arc therapy (VMAT) [I, A].

The treatment strategy for HPV-positive SCCHN should be the same as for HPV-negative SCCHN [I, A].

The recommended treatment option should be based on patient- and treatment-related factors (e.g. side-effects, complications, etc.) since conservative surgery and RT may often provide similar locoregional control [IV, A; consensus = 100%].

Early disease should be treated as much as possible with a single-modality treatment [IV, A; consensus = 100%].

Standard options for locally advanced disease are either surgery plus adjuvant chemoradiotherapy (CRT) or primary concomitant CRT [I, A].

Primary surgical treatment followed by RT or CRT is the preferred treatment for T3/T4 oral cavity and T4 laryngeal cancers [III, A], Figures 1 and 2 . Figure 1 Management of oral cavity cancer (stage I-IVB), excluding lip carcinoma.

BSC, best supportive care; c, clinical; ChT, chemotherapy; CRT, chemoradiotherapy; DOI, depth of invasion; M, metastasis; N, node; RT, radiotherapy; T, tumour.

a If DOI <10 mm: sentinel lymph node biopsy is a valid option; if DOI <5 mm and cT1N0, active surveillance of the neck is a valid option.

Management of laryngeal cancer (stage I-IVB).

BSC, best supportive care; c, clinical; ChT, chemotherapy; CRT, chemoradiotherapy; M, metastasis; N, node; RT, radiotherapy; T, tumour.

c cT1–2N0 glottic cancer does not require neck dissection or neck RT.

d Altered fractionation (accelerated or hyperfractionated) RT is a valid option for selected T3 or T3N1.

A lack of consensus in terms of 'acceptability' for 'recommendation 3i' below, however, was identified in the pre-meeting survey. The use of a hypoxic radiosensitiser has not been approved in China and is not practised in Singapore. The Asian experts agreed that the evidence was available to support this recommendation in the form of a meta-analysis, 26 but thought as it was conducted on older studies, that the GoR should be changed from IA to IC. The wording of the original recommendation was therefore revised from: 3i. A hypoxic radiosensitiser increases locoregional control and disease-free survival compared with RT alone [I, A].

to read as per the bold text below. 3i. A hypoxic radiosensitiser, if available, might be considered to increase locoregional control and disease-free survival compared with RT alone [I, C ; consensus = 100%] .

'Recommendation 3j' was accepted without change, with 100% consensus. 3j. Concomitant CRT increases locoregional control and overall survival compared with RT alone [I, A].

A lack of consensus was also identified in the feedback comments about 'recommendations 3k, 3l, 3m and 3n'. With regard to 'recommendation 3k, there was concern over the statement that the standard of care was 100 mg/m² cisplatin given on days 1, 22 and 43 of concomitant RT (70 Gy), because a weekly dose of 40 mg/m² cisplatin is also a standard treatment option in Asian countries. Also, weekly low dose 40 mg/m² cisplatin plus RT has been shown to have similar survival to 100 mg/m² cisplatin plus RT every 3 weeks with lower toxicity. 27 A retrospective multicentre study failed to find any difference in survival between weekly versus 3-weekly cisplatin dosing and concomitant RT, 28 although these findings are not universal, 29 , 30 and more prospective clinical studies are required. Also, in a separate study, weekly cisplatin plus RT versus cetuximab plus RT, showed weekly cisplatin plus concomitant RT to have superior outcomes to cetuximab plus RT. 31 A Japanese study has also shown weekly cisplatin plus RT to be non-inferior to 3-weekly cisplatin plus RT post-operatively in Japanese patients with high-risk, locally advanced SCCN. 32 Thus, an extra line was added to 'recommendation 3k' below (see bold text below and Table 1), with 100% consensus. There was also some discussion about 'recommendation 3l' and the use of cetuximab (Supplementary Table S5 , available at <https://doi.org/10.1016/j.esmoop.2021.100309>), and the observation that Asian patients are more susceptible to RT-induced mucositis. 'Recommendation 3l' was left unchanged with 100% consensus. In relation to 'recommendation 3m' the Asian experts were concerned that it should be made clear that induction chemotherapy was only to be administered for up to three cycles,

according to response, with the patient's disease evaluated after every cycle. The text of 'recommendation 3m' was thus revised accordingly (see bold text), with 100% consensus. In the case of 'recommendation 3n' it was noted that although induction chemotherapy was not used routinely with larynx preservation, the Japanese experts considered induction chemotherapy an option in the case of patients with very advanced or rapidly progressive locally advanced HNC and strongly recommended induction chemotherapy for patients with a high risk of distant metastasis including N2c, N3 and level IV (Supplementary Table S4 , available at <https://doi.org/10.1016/j.esmoop.2021.100309>). The text of 'recommendation 3n' remained unchanged. 3k. The standard of care for chemotherapy is cisplatin at a dose of 100 mg/m² given on days 1, 22 and 43 of concomitant RT [II, A]. Weekly cisplatin 40 mg/m² is an alternative option in the post-operative setting [II, A; consensus = 100%] (Figures 3 and 4). Figure 3 Management of oropharyngeal cancer (p16-negative stage I–IVB; p16-positive stage I–III).

c, clinical; CRT, chemoradiotherapy; M, metastasis; N, node; RT, radiotherapy; T, tumour.

a Altered fractionation (accelerated or hyperfractionated) RT is a valid option for T1–N1, T2–N0 or T2–N1.

b Altered fractionation (accelerated or hyperfractionated) RT is a valid option for T1–N1 or T2–N1.

c Altered fractionation (accelerated or hyperfractionated) RT is a valid option for T1–N1 or T2–N1.

Management of hypopharyngeal cancer (stage I–IVB).

BSC, best supportive care; c, clinical; ChT, chemotherapy; CRT, chemoradiotherapy; M, metastasis; N, node; RT, radiotherapy; T, tumour.

a In the case of patients unfit for curative treatment. However, curative treatment should be considered for most patients.

In patients unfit for cisplatin, carboplatin combined with 5-fluorouracil (5-FU), or cetuximab concomitant to RT, as well as hyperfractionated or accelerated RT without chemotherapy, are treatment alternatives [II, A; consensus = 100%].

For larynx preservation, induction chemotherapy with docetaxel, cisplatin and 5-FU (up to three courses according to response) followed by RT alone is a validated treatment option [I, A; consensus = 100%].

Besides larynx preservation, induction chemotherapy is not routinely recommended (Figure 2).

Neck dissection is not recommended in cases of negative FDG-PET and normal size lymph nodes at 12 weeks post-CRT [I, A].

Post-operative RT is recommended for patients with pT3-4 tumours, resection margins with macroscopic (R2) or microscopic (R1) residual disease, perineural infiltration, lymphatic infiltration, >1 invaded lymph node and the presence of extracapsular infiltration [II, A].

Post-operative CRT is recommended for patients with an R1 resection and extranodal extension [I, A].

A lack of consensus was also identified in the feedback comments with regard to 'recommendations 3r-t, and 3v'. The Asian experts recommended that the text of 'recommendation 3r' below was revised (see bold text) to make it clear that every effort should be made to ensure that patients receive RT or CRT within 6 weeks of surgery, and definitely no later than 8 weeks after surgery.

Emphasis was placed on the importance of liaising with the surgeon in an MDT environment, where and whenever possible. In relation to 'recommendation 3s' there was much discussion about the PD-L1 combined positive score (CPS) as an indicator of response to pembrolizumab based on the data from the phase III KEYNOTE (KN)-048 trial in which previously untreated patients with recurrent or metastatic SCCHN (R/M SCCHN) were randomised to receive pembrolizumab alone or chemotherapy plus either pembrolizumab or cetuximab. 33 Pembrolizumab alone improved overall survival versus chemotherapy plus cetuximab (EXTREME regimen 34) in patients with a CPS ≥ 20 [median 14.9 months versus 10.7 months, hazard ratio (HR) 0.61 (95% confidence interval 0.45-0.83), $P = 0.0007$] and also in patients with a CPS ≥ 1 [12.3 months versus 10.3 months, HR 0.78 (0.64-0.96), $P = 0.0086$] and was non-inferior in the total population. 33 Pembrolizumab plus chemotherapy was also superior to cetuximab plus chemotherapy in terms of overall survival in the total population [13.0 months versus 10.7 months, HR 0.77 (95% confidence interval 0.63-0.93), $P = 0.0034$] with the benefit again slightly greater in patients with a CPS ≥ 20 than in those with a CPS ≥ 1 and therefore the wording of 'recommendation 3s' was revised to reflect this, and reference made to the publication by Kiyota and Imamura 2020 35 in relation to 'recommendations 3s and 3t'. Although support for the EXTREME regimen first line has been shown in a Chinese phase III study 36 and a Japanese observational study, 37 the Asian experts considered pembrolizumab plus chemotherapy to also be a valid option for the treatment of PD-L1-negative disease based on the data from the KN-048 trial, 33 for the total patient population, especially as in some countries in Asia the EXTREME regimen is not reimbursed. Docetaxel, cisplatin and cetuximab (TPeX) (although the safety data are only available for induction therapy in Asian patients), 38 and paclitaxel and carboplatin plus cetuximab (PCE) which has shown promising activity in Japanese patients, 39 are alternative treatment options, and the wording of 'recommendation 3t' below was amended to reflect this with 100% consensus. 3r. Every effort should be made to ensure that the administration of post-operative RT or CRT starts within 6 weeks of surgery [II, A; consensus = 100%].

Pembrolizumab in combination with platinum/5-FU and pembrolizumab monotherapy are two approved regimens for patients with R/M SCCHN expressing PD-L1 (CPS ≥ 1) [I, A; ESMO-MCBS v1.1 score: 4]. The choice of pembrolizumab monotherapy or chemotherapy plus pembrolizumab may be based on CPS, tumour burden and symptoms [V, B; consensus = 100%].

Platinum/5-FU/cetuximab remains the standard therapy for patients with R/M SCCHN not expressing PD-L1 [I, A; ESMO-MCBS v1.1 score: 3]. 34 Pembrolizumab plus chemotherapy [II, C], TPeX 40 [II, B], and PCE [II, B] are also treatment options in this population [consensus = 100%] (Figure 5). Figure 5 Management of recurrent and/or metastatic disease not amenable to curative RT or surgery.

BSC, best supportive care; c, clinical; ChT, chemotherapy; CRT, chemoradiotherapy; 5-FU, 5-fluorouracil; M, metastasis; MCBS, Magnitude of Clinical Benefit Scale; N, node; PCE, paclitaxel, carboplatin and cetuximab; PD-L1, programmed death-ligand 1; RT, radiotherapy; T, tumour; TPeX, cisplatin/docetaxel/cetuximab.

Nivolumab is both FDA- and EMA-approved for recurrent/metastatic patients who progress within 6 months of platinum therapy 41 , 42 , 43 [I, A; ESMO-MCBS v1.1 score: 4].

Deficiencies in the functioning of dihydropyrimidine dehydrogenase (DPD), the main enzyme involved in fluoropyrimidine metabolism, due to genetic polymorphisms, occur in 3%-5% of Western/European patients and can lead to lethal fluoropyrimidine toxicity. 44 Due to the low incidence of DPD deficiency in Asian patients, however, DPD genotyping and phenotyping is not carried out in routine daily practice in Asia, but is recommended for patients, who experience severe 5-FU toxicity during and after their first cycle of chemotherapy. The original 'recommendation 3v' below was thus revised completely, to the version in bold text and Table 1 , to reflect this. 3v. DPD testing is recommended before initiating 5-FU.

According to the specific genetic profile of the Asian patient population, DPD genotyping or phenotyping may be considered before initiating fluoropyrimidine-based therapy [III, C; consensus = 100%].

4a. Clinical follow-up, including head and neck examination by flexible endoscopy, should be carried out every 2-3 months during the first 2 years, every 6 months for years 3-5 and annually thereafter [III, A].

Imaging should be carried out if symptoms occur or in cases of abnormalities found at the clinical examination [III, A].

FDG-PET/CT is recommended 3 months after CRT for patients with node-positive disease to assess the necessity of neck dissection [I, A].

The drug and testing availability for each of the seven Asian countries is summarised in Table 2 , and the ESMO-Magnitude of Clinical Benefit Scales (ESMO-MCBSs) for the different systemic therapy options for the treatment of SCCHN are presented in Table 3 . Resource limitations are the most important barrier to offering optimal diagnosis and treatment to patients with SCCHN across the different Asian countries. Table 2 Summary of applicability (availability) of drugs, equipment and testing according to Asian country

CSCO, Chinese Society of Clinical Oncology; CT, computed tomography; DNA, deoxyribonucleic acid; DPD, dihydropyrimidine dehydrogenase; EBER, Epstein–Barr-encoded RNA, HPV, human papilloma virus; IHC, immunohistochemistry; IMRT, intensity modulated radiotherapy ; ISH, in situ hybridisation; ISMPO, Indian Society of Medical and Paediatric Oncology; JSMO, Japanese Society of Medical Oncology; KSMO, Korean Society of Medical Oncology; PET, positron emission tomography; PD-L1, programmed death-ligand 1; RNA, ribonucleic acid; SSO, Singapore Society of Oncology; TOS, Taiwan Oncology Society; VMAT, volumetric modulated arc therapy.

ESMO-MCBS table for new therapies/indications in SCCHN

CI, confidence interval; CPS, combined positive score; ESMO-MCBS, European Society for Medical Oncology-Magnitude of Clinical Benefit Scale; 5-FU, 5-fluorouracil; HR, hazard ratio; OS, overall survival; PD-L1, programmed death-ligand 1; QoL, quality of life; R/M, recurrent/metastatic; SCCHN, squamous cell carcinoma of the head and neck.

ESMO-MCBS version 1.1. The scores have been calculated by the ESMO-MCBS Working Group and validated by the ESMO Guidelines Committee.

QoL exploratory endpoint, therefore, not creditable.

Three-arm trial comparing chemotherapy plus cetuximab versus chemotherapy plus pembrolizumab versus pembrolizumab monotherapy.

The licensed indication is for CPS PD-L1 expression ≥ 1 . This score relates to a planned subgroup illustrating enhanced benefit among a subset of the approved cohort with CPS PD- L1 expression ≥ 20 .

QoL evaluated as exploratory endpoint (as distinct from primary or secondary endpoint) is not eligible for ESMO-MCBS grading.

European Medicines Agency (EMA) approval is restricted to PD-L1 $\geq 50\%$ tumour proportion score (TPS). PD-L1 ≥ 1 CPS was a secondary endpoint eligible for ESMO-MCBS scoring.

EMA indication is restricted to recurrent or metastatic head and neck cancer with PD-L1 $\geq 50\%$ TPS. This approval is based on an exploratory analysis with no adjustment for multiplicity in which the median OS control arm was 6.6 months, with a gain of 5.0 months HR 0.53 (95% CI 0.35-0.81). Although exploratory analyses can be the basis for hypothesis generation or conjecture or even licensing approvals by regulatory authorities, since they are exploratory (as distinct from primary or secondary endpoints), they are not eligible for grading using ESMO-MCBS.

The recommendations listed in Table 1 can be considered to constitute the consensus Clinical Practice Guidelines for the treatment of patients with SSCHN (excluding nasopharyngeal cancer) in Asia, and are the result of voting by the Asian experts both before and during the virtual 'face-to-face' meeting hosted by KSMO, to adapt the recently published EHNS-ESMO-ESTRO Clinical Practice Guidelines. 23

Following 'face-to-face' discussions during the virtual meeting, the revisions highlighted in bold text in Table 1 were made to the wording of the recent EHNS-ESMO-ESTRO Clinical Practice Guideline 'recommendations' initially identified in the pre-meeting survey as not having the agreement of all the Asian experts, and resulted in a 100% consensus, being achieved for all the recommendations listed.

Despite these changes, these Pan-Asian adapted recommendations show high concordance with the original EHNS-ESMO-ESTRO Clinical Practice Guideline recommendations for the treatment of patients with SCCHN, 23 with the acceptance of each recommendation by each of the Asian experts based on the available scientific evidence independently of the approval and reimbursement status of certain drugs in their individual countries.

A summary of the availability of the recommended treatment modalities and recommended drugs, as of July 2021, is presented for each participating Asian country in Table 2 and will obviously impact on some of the disease and patient management strategies that can be adopted.

Sung H. Ferlay J. Siegel R.L. Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries *CA Cancer J Clin* 71 2021 209 249 33538338 10.3322/caac.21660 2 Zhang L.-W. Li J. Cong X. Incidence and mortality trends in oral and oropharyngeal cancer in China, 2005-2013 *Cancer Epidemiol* 57 2018 120 126 30396144 10.1016/j.canep.2018.10.014 3 Hwang T.Z. Hsiao J.R. Tsai C.R. Chang J.S. Incidence trends of human papillomavirus-related head and neck cancer in Taiwan, 1995-2009 *Int J Cancer* 137 2015 395 408 25395239 10.1002/ijc.29330 4 Suh J.D. Cho J.H. Trends in head and neck cancer in South Korea Between 1999 and 2012 *Clin Exp Otorhinolaryngol* 9 2016 263 269 27136364 10.21053/ceo.2015.01123 PMC4996101 5 Vital Statistics Japan. In: Ministry of Health Law (ed) Edition 2018 Available at https://ganjoho.jp/reg_stat/statistics/index.html 6 Hori M. Matsuda T. Shibata A. Cancer incidence and incidence rates in Japan in 2009: a study of 32 population-based cancer registries for the Monitoring of Cancer Incidence in Japan (MCIJ) project *Jpn J Clin Oncol* 45 2015 884 891 26142437 10.1093/jjco/hyv088 7 Report of the Head and Neck Cancer Registry of Japan. In: Edition 2016 Available at http://www.jshnc.umin.ne.jp/pdf/2016syourei_houkoku.pdf 8 Johnson D.E. Burtness B. Leemans C.R. Head and neck squamous cell carcinoma *Nat Rev Dis Primers* 6 2020 92 33243986 10.1038/s41572-020-00224-3 PMC7944998 9 Mehrta H. Duncan K. Parascandola M. Defining a global research and policy agenda for betel quid and areca nut *Lancet Oncol* 18 2017 e767 e775 29208442 10.1016/S1470-2045(17)30460-6 10 Mishra A. Meherotra R. Head and neck cancer: global burden and regional trends in India *Asian Pac J Cancer Prev* 15 2014 537 550 24568456 10.7314/apjcp.2014.15.2.537 11 Wong I.C. Ng Y.K. Lui V.W. Cancers of the lung, head and neck on the rise: perspectives on the genotoxicity of air pollution *Chin J Cancer* 33 2014 476 480 25011457 10.5732/cjc.014.10093 PMC4198750 12 Leemans C.R. Snijders P.J.F. Brakenhoff R.H. The molecular landscape of head and neck cancer *Nat Rev Cancer* 18 2018 269 282 29497144 10.1038/nrc.2018.11 13 Shaikh M.H. McMillan N.A. Johnson N.W. HPV-associated head and neck cancers in the Asia Pacific: a critical literature review & meta-analysis *Cancer Epidemiol* 39 2015 923 938 26523982 10.1016/j.canep.2015.09.013 14 Yap L.F. Lai S.L. Rhodes A. Clinico-pathological features of oropharyngeal squamous cell carcinomas in

Malaysia with reference to HPV infection Infect Agent Cancer 13 2018 21 29942347
10.1186/s13027-018-0193-6 PMC5996537 15 Bandhary S.K. Shetty V. Saldanha M. Detection of human papilloma virus and risk factors among patients with head and neck squamous cell carcinoma attending a tertiary referral centre in South India Asian Pac J Cancer Prev 19 2018 1325 1330 29802694 10.22034/APJCP.2018.19.5.1325 PMC6031850 16 Tan L.S. Fredrik P. Ker L. High-risk HPV genotypes and P16INK4a expression in a cohort of head and neck squamous cell carcinoma patients in Singapore Oncotarget 7 2016 86730 86739 27893418 10.18632/oncotarget.13502 PMC5349949 17 Babu G. Bahl A. Bhattacharya G.S. Oncology Gold Standard® practical consensus recommendations for the use of monoclonal antibodies in the management of squamous cell carcinoma of head and neck South Asian J Cancer 6 2017 154 160 29404293 10.4103/sajc.sajc_181_17 PMC5763625 18 Prabhash K. Babu G. Chaturvedi P. Indian clinical practice consensus guidelines for the management of very advanced disease of squamous cell carcinoma of head and neck Indian J Cancer 57 2020 S22 S25 32167069 10.4103/0019-509X.278977 19 Chinese Society of Clinical Oncology (CSCO) diagnosis and treatment guidelines for head and neck cancer 2018 (English version) Chin J Cancer Res 31 2019 84 98 30996568 10.21147/j.issn.1000-9604.2019.01.05 PMC6433588 20 Nibu K.I. Hayashi R. Asakage T. Japanese clinical practice guideline for head and neck cancer Auris Nasus Larynx 44 2017 375 380 28325607 10.1016/j.anl.2017.02.004 21 Ministry of Health Malaysia Systemic Therapy Protocol Third Edition In: Edition 2016 Available at <https://nci.moh.gov.my/index.php/ms/en/main-menu-2/polisi/271-polisi> 22 [Taiwan Cooperative Oncology Group under National Health Research Institutes in Taiwan] Available at <https://tcog.nhri.org.tw/wpcontent/uploads/2020/2005/2100oralpg.pdf> 23 Machiels J.P. Rene Leemans C. Golusinski W. Squamous cell carcinoma of the oral cavity, larynx, oropharynx and hypopharynx: EHNS-ESMO-ESTRO Clinical Practice Guidelines for diagnosis, treatment and follow-up Ann Oncol 31 2020 1462 1475 33239190 10.1016/j.annonc.2020.07.011 24 Dykewicz C.A. Centers for Disease Control and Prevention (U.S.) Infectious Diseases Society of America American Society of Blood and Marrow Transplantation Summary of the Guidelines for Preventing Opportunistic Infections among Hematopoietic Stem Cell Transplant Recipients Clin Infect Dis 33 2001 139 144 11418871 10.1086/321805 25 O'Sullivan B. Head and neck tumours Brierley J. Gospodarowicz M.K. Wittekind C. UICC TNM Classification of Malignant tumours 8th ed. 2017 Wiley Chichester 17 54 26 Overgaard J. Hypoxic modification of radiotherapy in squamous cell carcinoma of the head and neck—a systematic review and meta-analysis Radiother Oncol 100 2011 22 32 21511351 10.1016/j.radonc.2011.03.004 27 Bauml J.M. Vinnakota R. Anna Park Y.H. Cisplatin every 3 weeks versus weekly with definitive concurrent radiotherapy for squamous cell carcinoma of the head and neck J Natl Cancer Inst 111 2019 490 497 30239887 10.1093/jnci/djy133 PMC6510226 28 Helfenstein S. Riesterer O. Meier U.R. 3-weekly or weekly cisplatin concurrently with radiotherapy for patients with squamous cell carcinoma of the head and neck - a multicentre, retrospective analysis Radiat Oncol 14 2019 32 30744643 10.1186/s13014-019-1235-y PMC6371614 29 Noronha V. Joshi A. Patil V.M. Once-a-week versus once-every-3-weeks cisplatin chemoradiation for locally advanced head and neck cancer: a phase III randomized noninferiority trial J Clin Oncol 36 2018 1064 1072 29220295 10.1200/JCO.2017.74.9457 30 Osman N. Elamin Y.Y. Rafee S. Weekly cisplatin concurrently with radiotherapy in head and neck squamous cell cancer: a retrospective analysis of a tertiary institute experience Eur Arch Otorhinolaryngol 271 2014 2253 2259 24121822 10.1007/s00405-013-2749-9 31 Rischin D. King M.T. Kenny L.M. Randomized trial of radiation therapy with weekly cisplatin or cetuximab in low-risk HPV-associated oropharyngeal cancer (TROG 12.01) - a Trans-Tasman Radiation Oncology Group study Int J Radiat Oncol Biol Phys 111 4 2021 876 886 34098030 10.1016/j.ijrobp.2021.04.015 32 Kiyota N. Tahara M. Fujii H. Phase II/III trial of post-operative chemoradiotherapy comparing 3-weekly cisplatin with weekly cisplatin in high-risk patients with squamous cell carcinoma of head and neck (JCOG1008) J Clin Oncol 38 suppl 15 2020 abstr 6502 10.1093/jco/hyu067 24842866 33 Burtneess B. Harrington K.J. Greil R. Pembrolizumab alone or with chemotherapy versus cetuximab with chemotherapy for recurrent or metastatic squamous cell carcinoma of the head and neck (KEYNOTE-048): a randomised, open-label, phase 3 study Lancet 394 2019 1915 1928 31679945 10.1016/S0140-6736(19)32591-7 34 Vermorken J.B. Mesia R. Rivera F. Platinum-based chemotherapy plus cetuximab in head and neck cancer N Engl J Med 359 2008 1116 1127 18784101 10.1056/NEJMoa0802656 35 Kiyota N. Imamura Y. New first-line treatment for recurrent or metastatic squamous cell carcinoma of head and neck: does one size fit

all? *Therap Radiol Oncol* 4 2020 5 36 Guo Y. Luo Y. Zhang Q. First-line treatment with chemotherapy plus cetuximab in Chinese patients with recurrent and/or metastatic squamous cell carcinoma of the head and neck: efficacy and safety results of the randomised, phase III CHANGE-2 trial *Eur J Cancer* 156 2021 35 45 34418665 10.1016/j.ejca.2021.06.039 37 Yokota T. Ota Y. Fujii H. Real-world clinical outcomes and prognostic factors in Japanese patients with recurrent or metastatic squamous cell carcinoma of head and neck treated with chemotherapy plus cetuximab: a prospective observation study (JROSG12-2) *Int J Clin Oncol* 26 2021 316 325 33125596 10.1007/s10147-020-01817-4 38 Zenda S. Ota Y. Kiyota N. A multicenter phase II trial of docetaxel, cisplatin, and cetuximab (TPEx) followed by cetuximab and concurrent radiotherapy for patients with local advanced squamous cell carcinoma of the head and neck (CSPOR HN01: ECRIPS Study) *Front Oncol* 9 2019 6 30723701 10.3389/fonc.2019.00006 PMC6349830 39 Tahara M. Kiyota N. Yokota T. Phase II trial of combination treatment with paclitaxel, carboplatin and cetuximab (PCE) as first-line treatment in patients with recurrent and/or metastatic squamous cell carcinoma of the head and neck (CSPOR-HN02) *Ann Oncol* 29 2018 1004 1009 29408977 10.1093/annonc/mdy040 40 Guigay J. Fayette J. Mesia R. TPExtreme randomized trial: TPEx versus Extreme regimen in 1st line recurrent/metastatic head and neck squamous cell carcinoma (R/M HNSCC) *J Clin Oncol* 37 suppl 15 2019 6002 41 Ferris R.L. Blumenschein G. Jr. Fayette J. Nivolumab for recurrent squamous-cell carcinoma of the head and neck *N Engl J Med* 375 2016 1856 1867 27718784 10.1056/NEJMoa1602252 PMC5564292 42 Kiyota N. Hasegawa Y. Takahashi S. A randomized, open-label, phase III clinical trial of nivolumab vs. therapy of investigator's choice in recurrent squamous cell carcinoma of the head and neck: a subanalysis of Asian patients versus the global population in checkmate 141 *Oral Oncol* 73 2017 138 146 28939066 10.1016/j.oraloncology.2017.07.023 43 Yen C.J. Kiyota N. Hanai N. Two-year follow-up of a randomized phase III clinical trial of nivolumab vs. the investigator's choice of therapy in the Asian population for recurrent or metastatic squamous cell carcinoma of the head and neck (CheckMate 141) *Head Neck* 42 2020 2852 2862 32583557 10.1002/hed.26331 PMC7540331 44 Henricks L.M. Lunenburg C.A.T.C. de Man F.M. DPYD genotype-guided dose individualisation of fluoropyrimidine therapy in patients with cancer: a prospective safety analysis *Lancet Oncol* 19 2018 1459 1467 30348537 10.1016/S1470-2045(18)30686-7 45 Mesia R. Rivera F. Kaweck A. Quality of life of patients receiving platinum-based chemotherapy plus cetuximab first line for recurrent and/or metastatic squamous cell carcinoma of the head and neck *Ann Oncol* 21 2010 1967 1973 20335368 10.1093/annonc/mdq077 PMC2946862 46 Harrington K.J. Ferris R.L. Blumenschein G. Jr. Nivolumab versus standard, single-agent therapy of investigator's choice in recurrent or metastatic squamous cell carcinoma of the head and neck (CheckMate 141): health-related quality-of-life results from a randomised, phase 3 trial *Lancet Oncol* 18 2017 1104 1115 28651929 10.1016/S1470-2045(17)30421-7 PMC6461049 47 Cohen E.E.W. Soulières D. Le Tourneau C. Pembrolizumab versus methotrexate, docetaxel, or cetuximab for recurrent or metastatic head-and-neck squamous cell carcinoma (KEYNOTE-040): a randomised, open-label, phase 3 study *Lancet* 393 2019 156 167 30509740 10.1016/S0140-6736(18)31999-8 48 Harrington K.J. Soulières D. Le Tourneau C. Quality of life with pembrolizumab for recurrent and/or metastatic head and neck squamous cell carcinoma: KEYNOTE-040 *J Natl Cancer Inst* 113 2021 171 181 32407532 10.1093/jnci/djaa063 PMC7850527 Supplementary data Supplementary Methods Supplementary Tables

The authors thank Ms K. Marinoni and Ms D. Young from the Scientific and Medical Division of ESMO, Ms Z. Othman from the ESMO Singapore Office, Dr B. Devnani from the LPG Asia Alumni and Ms M. Kim of KSMO, for their assistance in the execution of the virtual 'face-to-face' meeting of experts. Dr A. Kinsella of Cancer Communications and Consultancy Ltd, Knutsford, Cheshire, UK is acknowledged for her contribution to the preparation of the manuscript. Mrs N. Latino, ESMO Head of Scientific Affairs, is acknowledged for her contribution in the completion of the ESMO-MCBS table.

No external funding has been received for the preparation of these guidelines. Production costs have been covered by 10.13039/501100007075 ESMO from central funds.

BK declares grants or contracts from Merck Sharp & Dohme (MSD) Oncology, 10.13039/501100013170 Ono Pharmaceutical and 10.13039/100004325 AstraZeneca , consulting fees from AstraZeneca, MSD Oncology, ABL Bio, Genexine, Cellid, Handok, Celid, Trial Informatics and CBS Bio, payment or honoraria from MSD Oncology and Merck. J-PM declares consulting fees/honoraria from Pfizer, Roche, AstraZeneca, Bayer, Innate, Merck Serono, Boehringer, Bristol Myers Squibb (BMS), Novartis, Janssen, Incyte, Cue Biopharma, ALX Oncology, iTEOS, TheRNA and NEKTAR, support for attending meetings and/or travel from Amgen, Pfizer and MSD and participation at a Safety or Advisory Board for PsiOxus. LL declares institutional grants or contracts from 10.13039/100004325 AstraZeneca , BMS , 10.13039/100008349 Boehringer Ingelheim , Celgene International , 10.13039/501100003769 Eisai , 10.13039/100010544 Exelixis Inc , Debiopharm International SA , 10.13039/100007013 Hoffman-La Roche Ltd. , IRX Therapeutics Inc. , 10.13039/100007050 Medpace Inc. , Merck-Serono , MSD , 10.13039/100004336 Novartis , 10.13039/100004319 Pfizer , Roche Spa and Buran and receipt of honoraria or fees (for public speaking/teaching in medical meetings and/or for providing expert opinion in Advisory Boards) for AstraZeneca, Bayer, BMS, Eisai, MSD, Merck-Serono, Boehringer Ingelheim, Hoffman La Roche Ltd., Novartis, Roche, Debiopharm International SA, Sobi, Incyte Biosciences Italy SRL, Doxa Pharma, Amgen, Nanobiotics and GlaxoSmithKline (GSK). CB declares payment or honoraria from Merck KGaA, BMS and Roche. MT declares consulting fees from Ono Pharmaceuticals, MSD, BMS and Merck Biopharma, and honoraria from Eisai, Ono Pharmaceuticals, BMS and Merck Biopharma. MA declares consulting fees from MSD, AstraZeneca, Eli Lilly, DKSH, Eisai, Roche, Novartis and Merck, payment or honoraria from MSD, AstraZeneca, Eli Lilly, DKSH, Eisai, Roche, Novartis and Merck, support for attending meetings and/or travel from MSD, Roche, Elekta/Abex and AstraZeneca, and is the president of the Malaysian Oncological Society. MKA declares honoraria for presentations from Pfizer and Boehringer Ingelheim, sponsorship for meetings from AstraZeneca, Boehringer Ingelheim and DKSH. QSNg declares support for attending meetings and or travel from BMS, Boehringer Ingelheim, MSD and Astellas, and participation in Safety or Advisory Boards for MSD and Boehringer Ingelheim. WIWZ declares honoraria for lectures from Amgen Malaysia, DKSH Malaysia, Eisai Malaysia, Eli Lilly Malaysia, Ipsen Malaysia, MSD Corp., a subsidiary of Merck & Co., Inc., Kenilworth, NJ, USA, Merck Serono Malaysia, Pfizer Malaysia and Roche Malaysia, travel grants from Amgen Malaysia, Celgene Malaysia, Eisai Malaysia, Eli Lilly Malaysia, MSD Corp., a subsidiary of Merck & Co., Inc., Kenilworth, NJ, USA, and Roche Malaysia, participation on an Advisory Board for Celgene Malaysia, Roche Malaysia, Eli Lilly Malaysia, Eisai Malaysia and MSD Corp., a subsidiary of Merck & Co., Inc., Kenilworth, NJ, USA, and is a member of ASCO AP Regional Council and Greater Petaling Cancer City Challenge UICC. NK declares institutional grants or contracts from 10.13039/501100013170 Ono Pharmaceutical , BMS , 10.13039/100004325 AstraZeneca , 10.13039/100004319 Pfizer , 10.13039/100010795 Chugai Pharmaceutical , Rakuten Medical , 10.13039/100004326 Bayer and Adlai Nortye , payment or honoraria from Ono Pharmaceutical, BMS, Merck Biopharma, MSD, Eisai and Bayer, participation on a Data Safety Monitoring Board or Advisory Board for Bayer and Adlai Nortye. GC declares institutional grants from 10.13039/100004334 Merck , consulting fees from BMS, Pfizer, MSD, AstraZeneca, Daiichi Sankyo, Lilly, Novartis and Seattle Genetics, payment or honoraria from AstraZeneca, Roche and Daiichi Sankyo. SP declares fees for consultancy/advisory roles from AbbVie, Amgen, AstraZeneca, Bayer, Beigene, Biocartis, Boehringer Ingelheim, BMS, Clovis, Daiichi Sankyo, Debiopharm, ecancer, Eli Lilly, Elsevier, Foundation Medicine, Illumina, Imedex, Incyte, Janssen, Medscape, MSD, Merck Serono, Merrimack, Novartis, PharmaMar, Phosphatin Therapeutics, PER, Pfizer, PRIME, Regeneron, Roche/Genentech, RTP, Sanofi, Seattle Genetics, Takeda, speaker roles for AstraZeneca, Boehringer Ingelheim, BMS, ecancer, Eli Lilly, Illumina, Imedex, Medscape, MSD, Novartis, PER, Pfizer, Prime, Roche/Genentech, RTP, Sanofi, Takeda and the receipt of grants/research support: (Sub) investigator in trials (institutional financial support for trials) sponsored by 10.13039/100002429 Amgen , 10.13039/100004325 AstraZeneca , Biodesix , 10.13039/100008349 Boehringer Ingelheim , BMS , Clovis , 10.13039/100015542 GSK , 10.13039/100010905 Illumina , 10.13039/100004312 Lilly , MSD , Merck Serono , Mirati , 10.13039/100004336 Novartis , and Pfizer, Phosphatin Therapeutics , 10.13039/100004337 Roche / 10.13039/100004328 Genentech . TWK declares institutional grants or contracts from 10.13039/100004337 Roche , and 10.13039/100004339 sanofi-aventis . TY declares institutional grants or contracts from 10.13039/100009954 Taiho Pharmaceutical , Sumitomo Dainippon , 10.13039/501100013170 Ono Pharmaceutical , 10.13039/100010795 Chugai Pharmaceutical ,

10.13039/100002429 Amgen , Parexel International , MSD , 10.13039/501100002973 Daiichi-Sankyo and 10.13039/100004339 Sanofi . All other authors have declared no conflicts of interest.