

PRACTICE GUIDELINE

Primary sclerosing cholangitis in children with inflammatory bowel disease: An ESPGHAN position paper from the Hepatology Committee and the IBD Porto group



Patrick F. van Rheenen¹ | Kaija-Leena Kolho² | Richard K. Russell³ | Marina Aloï⁴ | Annamaria Deganello⁵ | Séamus Hussey⁶ | Norman Junge⁷ | Jan De Laffolie⁸ | Mark R. Deneau⁹ | Emer Fitzpatrick⁶ | Anne M. Griffiths¹⁰ | Iva Hojsak¹¹ | Emanuele Nicastro¹² | Andreia Nita¹³ | Mikko Pakarinen¹⁴ | Amanda Ricciuto¹⁰ | Lissy de Ridder¹⁵ | Aurelio Sonzogni¹⁶ | Andrea Tenca¹⁷ | Marianne Samyn¹⁸ | Giuseppe Indolfi¹⁹

Correspondence

Patrick F. van Rheenen, Department of Paediatric Gastroenterology, University of Groningen, University Medical Centre Groningen, Beatrix Children's Hospital, PO Box 30001, 9700 RB Groningen, The Netherlands.
Email: p.f.van.rheenen@umcg.nl

Funding information

European Society for Paediatric Gastroenterology Hepatology and Nutrition

Abstract

Objective: We aimed to provide an evidence-supported approach to diagnose, monitor, and treat children with inflammatory bowel disease (IBD) and primary sclerosing cholangitis (PSC).

Methods: The core group formulated seven PICO-structured clinical questions. A systematic literature search from inception to December 2022 was conducted by a medical librarian using MEDLINE and EMBASE. Core messages from the literature were phrased as position statements and then circulated to a sounding board composed of international experts in pediatric gastroenterology and hepatology, histopathology, adult gastroenterology and hepatology, radiology, and surgery. Statements reaching at least 80% agreement were considered as final. The other statements were refined and then subjected to a second online vote or rejection.

Results: Regular screening for gamma-glutamyltransferase (GGT) is essential for detecting possible biliary disease in children with IBD. MR cholangiopancreatography is the radiological modality of choice for establishing the diagnosis of PSC. Liver biopsy is relevant in the evaluation of small duct PSC or autoimmune hepatitis. Children who do not have known IBD at the time of PSC diagnosis should undergo initial screening with fecal calprotectin for asymptomatic colitis, and then at least once yearly thereafter. Children with a cholestatic liver enzyme profile can be

For affiliations refer to page 390.

CME module may be found at <https://learnonline.naspghan.org/jpgn2>

Patrick F. van Rheenen, Kaija-Leena Kolho, and Richard K. Russell contributed equally and share the first authorship.

Marianne Samyn and Giuseppe Indolfi contributed equally and share the last authorship.

Disclaimers: Although this paper is produced by the ESPGHAN Hepatology Committee and the IBD Porto Group it does not necessarily represent ESPGHAN policy and is not endorsed by ESPGHAN.

ESPGHAN is not responsible for the practices of physicians and provides guidelines and position papers as indicators of best practice only. Diagnosis and treatment is at the discretion of the healthcare provider.


This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2024 The Author(s). *Journal of Pediatric Gastroenterology and Nutrition* published by Wiley Periodicals LLC on behalf of European Society for Pediatric Gastroenterology, Hepatology, and Nutrition and North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition.

considered for treatment with ursodeoxycholic acid and can continue if there is a meaningful reduction or normalization in GGT. Oral vancomycin may have a beneficial effect on GGT and intestinal inflammation, but judicious use is recommended due to the lack of long-term studies. Children with PSC–IBD combined with convincing features of autoimmune hepatitis may benefit from corticosteroids and antimetabolites.

Conclusions: We present state-of-the-art guidance on the diagnostic criteria, follow-up strategies, and therapeutic strategies and point out research gaps in children and adolescents with PSC–IBD.

How to diagnose, monitor and treat children with Primary Sclerosing Cholangitis - Inflammatory Bowel Disease (PSC-IBD)



van Rheenen, et al. Primary sclerosing cholangitis in children with inflammatory bowel disease: An ESPGHAN position paper from the Hepatology Committee and the IBD Porto group. *J Pediatr Gastroenterol Nutr.* (2025)

JPGN
Journal of Pediatric Gastroenterology and Nutrition

KEYWORDS

calprotectin, colorectal carcinoma, MRCP, surveillance colonoscopy, ursodeoxycholic acid

1 | INTRODUCTION

Primary sclerosing cholangitis (PSC) is a progressive liver disease with a chronic course, characterized by the destruction of the intra- and extrahepatic bile ducts. Approximately 5%–10% of patients with childhood-onset inflammatory bowel disease (IBD) have or will develop PSC.^{1–3} Conversely, children who do not have known IBD at the time of PSC diagnosis are likely to develop IBD in the years thereafter. In a cohort of children with both PSC and IBD (PSC–IBD), 60% were diagnosed with both conditions simultaneously. In 26%, PSC was diagnosed after IBD, and in 14%, IBD was diagnosed after PSC.⁴ Thirty to sixty percent of children with PSC will also display features of autoimmune hepatitis (AIH).^{4,5}

With long-term complications such as cirrhosis, hepatopancreatobiliary and colorectal malignancies, which can become evident in young adulthood, children and adolescents with childhood-onset PSC–IBD carry a high disease burden with uncertain outcomes including disability and impact on their work and/or education.

Several practice guidelines have been published on the management of adult PSC,^{6–8} but currently, there is no standardized guidance for the management of patients with childhood-onset IBD and PSC. To address this, the European Society for Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN) governing board commissioned a core group of experts from the

What is Known

- Primary sclerosing cholangitis (PSC) is an uncommon condition in childhood-onset inflammatory bowel disease (IBD) with severe outcomes, and there is no consensus on diagnostic and treatment approaches.

What is New

- The European Society for Paediatric Gastroenterology, Hepatology and Nutrition Hepatology Committee and the Pediatric IBD Porto group formulated clinical questions considered to be relevant for the care of children with PSC–IBD.
- An evidence-supported approach to diagnose, monitor, and treat children with PSC–IBD is presented here, as well as suggestions for future studies based on current knowledge gaps.

Hepatology Committee and the Paediatric IBD Porto group to provide a position article. The aim of this position article is to provide pediatric gastroenterologists with an evidence-supported overview of the

BOX 1. Nomenclature

The International primary sclerosing cholangitis (PSC) Study Group recently provided the following working definitions¹²:

Large duct PSC: High-quality magnetic resonance cholangiopancreatography (MRCP) or endoscopic retrograde cholangiopancreatography (ERCP) with features compatible with sclerosing cholangitis in the absence of apparent causes of secondary sclerosing cholangitis.

Small duct PSC: histologic features typical of PSC with normal and recent (<1 year) high-quality MRCP or ERC in the absence of other cholestatic disorders.

PSC with features of autoimmune hepatitis: PSC with biochemical, serological, or histological features of autoimmune hepatitis.

In most pediatric studies the commonly used term for the overlap syndrome between sclerosing cholangitis and autoimmune hepatitis is “autoimmune sclerosing cholangitis” or “juvenile sclerosing cholangitis”, where the latter term acknowledges the significantly higher prevalence of this condition in children as compared to adults.

diagnostic and therapeutic approach of PSC associated with IBD (Box 1).

BOX 2. Clinical questions

1. In patients with suspected inflammatory bowel disease (IBD), is the risk of coexistence of primary sclerosing cholangitis (PSC) or autoimmune hepatitis (AIH) higher in the case of elevated liver enzymes? If so, what is the predictive value of the liver enzyme tests?
2. In patients with IBD and elevated gamma-glutamyltransferase (GGT), which of the two diagnostic modalities—liver biopsy or magnetic resonance cholangiography—best confirms sclerosing cholangitis? Which of the two has the best predictive value?
3. In children with PSC, with or without overlapping AIH, can fecal calprotectin be used to screen for IBD?
4. What is the time-to-cancer diagnosis in children and adolescents with PSC?
5. Should children and adolescents with PSC be treated with ursodeoxycholic acid (UDCA) to prevent or reduce liver-related complications?
6. Should children and adolescents with PSC be treated with vancomycin to prevent or reduce liver-related complications?
7. In patients with IBD–PSC with features of AIH, is long-term treatment with corticosteroids and antimetabolites justified? See Supporting Information: Table S1 for an elaborated version.

2 | METHODOLOGY

The core group formulated seven clinical questions (see Box 2), structured by the Population-Intervention-Comparator-Outcome or Population-Exposure-Outcome format, to assimilate evidence from the literature and to prepare core messages as *position statements*. The position statements and supporting text were circulated to a sounding board composed of international experts in pediatric gastroenterology and hepatology, histopathology, adult gastroenterology and hepatology, radiology, and surgery. Following evaluation through an online platform, position statements reaching at least 80% agreement were considered as final. The other statements were refined and then subjected to a second online vote or rejection.

2.1 | Data sources and searches

We searched MEDLINE (through PubMed) and EMBASE from inception to December 2, 2022. The search

strategies were developed in collaboration with a medical information specialist. The search strategies for each of the electronic databases are shown in Supporting Information: Tables S2–S8. No language restrictions were applied.

2.2 | Classification of the quality of evidence

The quality of evidence was classified into four categories following the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach (Table 1). We categorized recommendations as weak or strong, in favor of or against an intervention. Weak recommendations imply that there was a significant variation in the decision of the voters and that physicians need to engage in a shared decision-making process with patients. Strong recommendations suggest that almost all voters would choose that intervention. In such a case, physicians usually do not need to present an alternative intervention.

TABLE 1 GRADE certainty ratings.

Certainty	What it means
High	We have a lot of confidence that the true effect is similar to the estimated effect
Moderate	We believe that the true effect is probably close to the estimated effect
Low	The true effect might be markedly different from the estimated effect
Very low	The true effect is probably markedly different from the estimated effect

2.3 | Position statements

For readability, this paper is divided into four main sections: diagnostic criteria, follow-up strategies, therapeutic strategies, and research agenda. Each position statement (with the quality of evidence and, if applicable, the strength of recommendation) is framed and followed by a discussion of the evidence. Practical guidance sections complement the evidence by providing additional information not covered by the position statements.

2.4 | Practice points

Practice points are included based on the consensus of the core group and reflect common practice where evidence is lacking. The practice points are framed, but they are not accompanied by statements about the quality of the evidence and the strength of the recommendation.

3 | RESULTS

3.1 | Screening process

Supporting Information: Figures S1–S7 show the literature screening process. A total of 2121 records were screened in duplicate, after which 309 reports were sought for retrieval. In total, 39 studies with sufficient methodological quality to answer our clinical questions were used to formulate position statements (Supporting Information: Tables S9–S15).

4 | DIAGNOSTIC CRITERIA

4.1 | Basic liver test panel

Position statement 1

In children with suspected or confirmed IBD the risk of coexistent liver disease is significantly higher in patients with elevated liver enzymes.

We recommend testing liver enzymes (ALT and GGT) regularly to screen for associated liver disease. Significant or persistent elevation of liver enzymes prompts further diagnostic work-up into an underlying liver disease.

Moderate certainty | Agreement: 95% | Strong recommendation

Practice point

Screening is usually performed at 3 to 6 months intervals and a work-up for underlying liver disease is most commonly initiated when liver enzymes exceed 2x the upper limit of normal.

4.1.1 | Evidence

Liver enzyme elevations are common in children with IBD, but little is known about the action threshold that justifies a diagnostic work-up for underlying liver disease. We searched for studies that assessed the diagnostic accuracy of elevated liver enzymes in patients who were not known to have liver disease. We found four studies that explored liver enzyme testing in IBD (Supporting Information: Table S9). There was considerable heterogeneity among the study cohorts regarding differences in the prevalence of PSC (ranging from 3% to 47%), group size, and the timing of blood sampling relative to the IBD diagnosis.^{3,9–11} Nevertheless, the negative predictive value of GGT was high in all cohorts (99% to 100%), indicating that a value below 50 U/L can reliably exclude cholestatic liver disease.

4.1.2 | Practical guidance

In adults, it is recommended to evaluate a diagnosis of PSC when serum markers of cholestasis, including alkaline phosphatase, GGT and/or bilirubin, are elevated.⁸ Due to its variation with age because of increased bone turnover and growth, alkaline phosphatase is not a reliable marker of biliary injury in children, hence should not be included in diagnostic testing.¹²

In children with IBD, abnormalities in alanine aminotransferase (ALT) and GGT frequently occur.¹ This usually involves mild elevations (less than twice the upper limit of normal [ULN]) associated with a transient and undefined etiology. If mild liver enzyme elevations persist for more than 1 month, further evaluation is required. Moderate to marked liver enzyme elevations ($\geq 2 \times$ ULN) warrant prompt investigation of etiology.^{13–15} Table 2 illustrates the common causes of liver enzyme abnormalities in IBD. As corticosteroids and antimetabolites could affect the presentation and course of the liver disease, liver enzymes should be checked at the time of IBD diagnosis and at regular intervals thereafter. All patients with IBD should have their liver enzymes tested half yearly as a minimum,¹⁴ although in practice the majority of the children

with IBD will have them screened much more frequently than this.

The most common cause of significantly elevated liver enzymes in children with IBD is drug-induced liver injury (DILI). Most drugs used for the treatment of IBD,¹⁶ as well as antibiotics, are known to be hepatotoxic and can be found on a searchable database maintained by the National Institute of Diabetes and Digestive, and Kidney Diseases, called LiverTox.¹⁷ Attributing abnormalities in ALT and GGT to any of these drugs requires systematic evaluation, including time to liver test abnormality after the implicated drug has been started (latency), resolution after the drug is stopped (dechallenge), recurrence on re-exposure (rechallenge) and information on the drug's potential for hepatotoxicity (likelihood).^{13,18}

TABLE 2 Common causes of liver enzyme elevations in children with inflammatory bowel disease.

Causes	Biochemical profile		
	Hepatocellular	Cholestatic	Mixed type
	ALT $\uparrow\uparrow$ to $\uparrow\uparrow\uparrow$ GGT normal to $\uparrow$	ALT $\uparrow$ GGT $\uparrow\uparrow$ – $\uparrow\uparrow\uparrow$	
Immune-related liver disease			
Autoimmune hepatitis (AIH)	☒	☐	☐
Primary sclerosing cholangitis (PSC)	☐	☒	☐
PSC with features of AIH	☐	☐	☒
Viral hepatitis			
Primary infection (Hepatitis A/B/C/E, EBV, CMV)	☒	☐	☐
Reactivation (Hepatitis B, EBV, CMV) associated with anti-TNF or steroid use	☒	☐	☐
Drug-induced liver injury^a			
Mesalamine, thiopurines, allopurinol, methotrexate and anti-TNF	☒	☒	☐
metabolic-dysfunction associated steatotic liver disease			
Hepatic steatosis caused by obesity or starvation	☒	☐	☐
Biliary obstruction			
Bile stones (unrelated to PSC)	☐	☒	☐
Other, e.g.^b			
Coeliac disease	☒	☐	☐
Wilson disease	☒	☐	☐
Alpha-1-antitrypsin deficiency	☒	☐	☐
Undefined			
Rapid normalisation of liver enzymes	☐	☐	☒

Note: $\uparrow$: mild elevation ($<2 \times$ ULN); $\uparrow\uparrow$: moderate elevation (2 to $5 \times$ ULN); $\uparrow\uparrow\uparrow$: marked elevation ($>5 \times$ ULN).

Abbreviations: ALT, alanine aminotransferase; CMV, cytomegalovirus; EBV, Epstein-Barr virus; GGT, gamma-glutamyltransferase; TNF, tumor necrosis factor; ULN, upper limit of normal.

^aHere, the focus is on IBD medication. However, many more drugs are potentially hepatotoxic. For a complete overview see livertox.nih.gov.

^bThese diseases may present with mild ALT elevation.

The ratio between ALT and GGT determines the biochemical profile and may be useful to narrow down the extensive etiology checklist (Table 2). A hepatocellular biochemical profile is characterized by a disproportionate elevation of ALT compared with GGT. Common causes include viral hepatitis and AIH. The latter has a weak bidirectional association with IBD. In adult patients with IBD, AIH coexists in less than 0.5% of cases, but the prevalence is 3–5 times higher in children and adolescents.^{14,19} In patients with IBD and AIH, the presence of additional sclerosing cholangitis should be actively evaluated with magnetic resonance cholangiopancreatography (MRCP) (see Section 4.3), as PSC–IBD with AIH features is more common than IBD with isolated AIH.²⁰

A cholestatic biochemical profile is a disproportionate elevation of GGT compared with ALT, which is typically seen in PSC. PSC is characterized by inflammation and fibrosis of the entire biliary tree, which ultimately leads to multifocal bile duct strictures and dilations. PSC with features of AIH exhibits a mixed-type biochemical profile.

4.2 | Immunology

AIH is characterized by elevated immunoglobulin G levels (>1x ULN) and presence of circulating autoantibodies. Autoantibodies can be present in other liver disorders, including PSC, and are not diagnostic in isolation.²¹ Testing for autoantibodies by indirect immunofluorescence on a freshly prepared rodent substrate should include antinuclear antibody (ANA), antismooth muscle antibody (anti-SMA), antiliver kidney microsomal type 1 antibody (anti-LKM-1), antiliver cytosol type 1 antibody (anti-LC1), and antisoluble liver antigen (anti-SLA), if available.²¹ Antinuclear cytoplasmic antibody (ANCA) positivity is common in colonic IBD and therefore not useful when screening for AIH.⁸ Remission of AIH is considered complete when transaminases and IgG levels are normal, and when autoantibodies are negative or low-titer (i.e., <1:20 for ANA and SMA and <1:10 for anti-LKM1 and anti-LC1).²²

4.3 | Imaging

Position statement 2

Use MRCP as the radiological modality of choice for diagnosing PSC.

Moderate certainty | Agreement: 100% | Strong recommendation

4.3.1 | Evidence

Transabdominal ultrasound is recommended as a baseline imaging study when evaluating elevated liver enzymes. In PSC, bile duct wall thickening and focal bile duct dilatations may be demonstrated on ultrasound, however formal imaging of the biliary tree is essential for the diagnosis and classification of PSC (Box 1). Historically, direct cholangiography was performed via the endoscopic route (ERCP), which has now been largely replaced by MRCP. The latter is less invasive and easier to perform. The role of ERCP is currently mainly therapeutic, which will be further discussed in Section 6.5. Supporting Information: Table S10 shows the results of four pediatric studies in which the diagnostic yield of MRCP (index test) was compared with liver biopsy (reference test).^{23–26} It is unsurprising that children presenting solely with small duct PSC, either alone or with overlapping features of AIH, had a negative MRCP.

4.3.2 | Practical guidance

The presence of a cholestatic liver enzyme profile and typical radiological findings of intra- or extrahepatic bile duct abnormalities on MRCP (Figure 1), in the absence of biochemical and serological evidence of AIH, supports a diagnosis of PSC. MRCP should be reviewed by experienced radiologists to exclude other possible diagnoses such as choledochal malformations. The Imaging working group of the International PSC Study Group recently provided a consensus document with definitions and suggested reporting standards for MRCP features of PSC, which allows for a standardized approach to diagnosis, assessment of disease severity, follow-up, and detection of complications.²⁷

4.4 | Liver biopsy

Position statement 3

Consider performing a liver biopsy in children with IBD and suspected PSC in the following circumstances:

- Normal biliary tree at MRCP, or
- Raised immunoglobulin G and the presence of liver-specific autoantibodies, or
- Clinical uncertainty before steroid induction therapy for IBD.

Low certainty | Agreement: 84% | Weak recommendation




FIGURE 1 Images from a magnetic resonance cholangiopancreatography (A) and an endoscopic retrograde cholangiopancreatography (B) taken from the same patient with primary sclerosing cholangitis demonstrating irregular intrahepatic and extrahepatic bile ducts. (Courtesy of Dr. Patrick van Rheenen, University Medical Center Groningen, The Netherlands).

4.4.1 | Evidence

Typical histological features of PSC are periductal fibrosis, ductular reaction, periductal inflammation and less commonly fibro-obliterative cholangitis (Figure 2A). A liver biopsy specimen showing interface hepatitis, characterized by a dense infiltrate of plasma cells and lymphocytes which invade the surrounding parenchyma beyond the limiting plate, confirms AIH (Figure 2B) or PSC with features of AIH (Figure 2C). It is worth noting that portal inflammation and interface hepatitis may also be observed in patients with PSC.¹² Histological criteria to decide whether inflammatory abnormalities in the presence of periductal fibrosis justify treatment with corticosteroids and antimetabolites do not exist. In addition to standard hematoxylin and eosin (H&E) staining as is shown in Figure 2, cytokeratin 7 immunohistochemical staining is employed to facilitate a more comprehensive examination of biliary structures.²⁸

4.4.2 | Practical guidance

Liver histology is required to evaluate the presence of small duct PSC in children with a predominant cholestatic biochemical profile and a normal biliary tree at MRCP, and it is the most accurate tool to demonstrate AIH (Figure 3). Children with a pure cholestatic liver enzyme profile and large duct PSC on MRCP may not require a liver biopsy, but the decision whether a liver biopsy is appropriate in this scenario is at the discretion of the professionals managing the patient.

4.5 | Fecal calprotectin

Practice point

Children who do not have known IBD at the time of PSC diagnosis should undergo initial screening with fecal calprotectin for asymptomatic colitis.

Position statement 4

Perform fecal calprotectin screening at least once yearly in children with isolated PSC and/or AIH to select patients for diagnostic endoscopy for suspected inflammatory bowel disease.

Low certainty | Agreement: 90% | Strong recommendation

Practice point

There is no clear cutoff value of fecal calprotectin to predict colonic inflammation in patients with PSC. The guideline panel suggests that values ≥ 150 $\mu\text{g/g}$ usually justify an ileocolonoscopy.

4.5.1 | Evidence

Most children with PSC and IBD have colitis with a predilection for more severe inflammation in the right colon, as opposed to ulcerative colitis (UC) which is characterized by uniform inflammation of the colon and rectum, or increasing severity of inflammation towards the rectum. Children with PSC whose IBD is labeled Crohn's disease usually have colonic involvement. Isolated ileitis or extensive small bowel involvement is very rare in PSC-IBD.²⁹ Figure 4 shows the typical distribution of inflammation with rectal sparing and/or backwash ileitis in the setting of pancolitis.^{29,30}

Persistent rectal bleeding or perianal disease should prompt endoscopic assessment of the gut.³¹ When the indication for endoscopy is less obvious, as in the case of children with chronic abdominal pain or nonbloody diarrhea, measuring fecal calprotectin may help to distinguish who is in need of a colonoscopy. A raised value signifies the presence of active inflammation warranting colonoscopy, whereas a normal value reduces the probability of finding inflammation to almost zero.³² It is unknown if this test strategy performs equally well in children with PSC and asymptomatic colitis, where there is typically mild intestinal disease, a reversed right to left gradient of inflammation, and rectal sparing.^{29,33} Despite milder colonic inflammation in PSC-IBD, fecal calprotectin values may be very similar to that in patients with "non-PSC IBD."³⁴ This apparent paradox has been provisionally explained in a study among adult patients with active PSC, in




FIGURE 2 Hematoxylin and eosin stained pathology slides. (A) is taken from a patient with primary sclerosing cholangitis (PSC) showing periductal onion skin fibrosis. (B) is taken from a patient with autoimmune hepatitis (AIH) showing an infiltrate of plasma cells and lymphocytes extending beyond the limiting plate. (C) is taken from a patient with PSC and features of AIH showing bile duct injury and interface hepatitis. Bar = 50 μ m. (Courtesy of Dr. Marius van den Heuvel, histopathologist, University Medical Center Groningen, The Netherlands).




FIGURE 3 Conceptual diagnostic model for autoimmune liver disease in children with inflammatory bowel disease. AIH, autoimmune hepatitis; MRCP, magnetic resonance cholangiopancreatography; PSC, primary sclerosing cholangitis.

whom endoscopically quiescent colitis (assessed with the UC endoscopic index of severity [UCEIS]) was correlated with fecal calprotectin, biliary calprotectin (samples collected during ERCP) and markers of cholestasis.³⁵ It was hypothesized that fecal calprotectin may not necessarily reflect only the severity of colonic inflammation, but also the severity of the concurrent biliary inflammation.

4.5.2 | Practical guidance

There is no consensus on the ideal fecal calprotectin cutoff point to justify a colonoscopy. In a prospective

case-control study performed at the Hospital for Sick Children in Toronto fecal calprotectin values in children with PSC and colitis (cases, $n = 37$) and children with colitis without liver disease (controls, $n = 50$) were correlated with the UCEIS. The colonoscopy was executed as part of routine clinical care. The decision to perform a colonoscopy was irrespective of the fecal calprotectin result. The authors concluded that a fecal calprotectin level $<93 \mu\text{g/g}$ best predicted endoscopic healing (UCEIS = 0) in children with PSC-IBD.³⁶ This application of fecal calprotectin (to predict endoscopic remission) differs from the intended use as stated in Box 2, where calprotectin screening is used to predict




FIGURE 4 The IBD phenotype of PSC-IBD. IBD, inflammatory bowel disease; PSC, primary sclerosing cholangitis.

active inflammation in the colon. The guideline panel suggests that values $\geq 150 \mu\text{g/g}$ usually justify endoscopy, which is in line with the European Crohn's and Colitis Organization (ECCO)–European Society for Gastrointestinal and Abdominal Radiology (ESGAR) guideline for diagnostic assessment in IBD.³⁷

5 | FOLLOW-UP STRATEGIES

5.1 | Natural progression of PSC

A considerable proportion of children and adolescents with PSC-IBD have a slowly progressive liver disease characterized by hepatobiliary fibrosis, biliary strictures, intermittent bacterial cholangitis, and, eventually, the development of cirrhosis and end-stage liver disease. In a large multicenter cohort study in which 781 patients with childhood-onset PSC were followed longitudinally, approximately 50% developed these complications within the first 10 years after the diagnosis of PSC. Once these complications arose, the median time to liver transplantation was 3–4 years.⁵

Patients with PSC who have features of AIH and those with small-duct PSC seem to have a more extended period until the development of complications. There is a debate among experts whether these patients have an earlier stage of the same disease or rather a distinct PSC phenotype.³⁸ The first concept assumes that the stage with immune-mediated hepatitis can be suppressed with steroids and thiopurines, while the biliary disease continues to deteriorate. In the end, the extent of biliary damage determines the long-term outcome.

5.2 | Counseling

Various prognostic markers have been explored to predict patient outcomes. Achieving spontaneous or treatment-associated normalization of GGT (i.e., $<50 \text{ IU/L}$) or a percentage decrease of $>25\%$ within the first year after a PSC diagnosis has been linked to a reduced occurrence of complications in the first 5-year period.^{39,40}

In the adult PSC literature, the presence of fibrosis or cirrhosis on liver biopsy, as well as increased liver stiffness (measured through vibration-controlled transient elastography), are cited as unfavorable predictors.⁷ The use of elastography for monitoring liver fibrosis in children is rapidly advancing but currently lacks clear guidance on interpreting test results.

The Pediatric PSC Consortium developed the Sclerosing Cholangitis Outcomes in Pediatrics (SCOPE) index based on the retrospective data of over 1000 children. The SCOPE index combines the results of four blood markers (total bilirubin, albumin, platelet count, and GGT) with cholangiography and may help to assess the child's risk for liver transplantation at diagnosis and over time.⁴¹ Children in the low-risk SCOPE category are unlikely to require a liver transplant in the next decade, while those in the medium- or high-risk SCOPE category probably require a referral to a pediatric hepatologist with expertise in PSC. The SCOPE index has not yet been prospectively validated. Until that time, the decision to refer a child to a liver transplant center should not be based on the SCOPE index. More information about referral criteria for liver transplantation can be found in Section 5.6.

5.3 | Monitoring of disease progression

Monitoring of the progression of PSC is based on the regular reappraisal of clinical signs and symptoms, inflammatory activity in the gut and the liver, biliary involvement, and parenchymal fibrosis/cirrhosis (Table 3). The latter may be evaluated with elastography or radiological imaging. Liver biopsy to evaluate progression of fibrosis in isolation is not recommended.

Performing an annual MRCP in children with PSC is not indicated due to the extremely low risk of cholangiocarcinoma (CCA) before the age of 18 (see

TABLE 3 Monitoring disease progression of PSC.

Category	Test	Test frequency
Clinical evaluation	Fatigue	Every 3–6 months
	Pain right upper quadrant	
	Bleeding tendency	
	Jaundice	
	Pruritis	
	Fever of unknown origin	
	Nutritional state	
Blood tests	Quality of life	Every 3–6 months
	ALT	
	AST	
	GGT	
	TSB	
	Albumin	
	INR	
	Platelets	
	CRP	
	IgG	Every 12 months
	AFP	
Imaging	Lipid soluble vitamins	In patients with negative tests at diagnosis and increased IgG during follow-up
	Autoantibodies (ANA, anti-SMA, anti-LKM-1, anti-LC1, and anti-SLA)	
	Serum bile acids	Optional
	Ultrasonography (liver parenchyma and bile ducts); in case of cirrhosis combined with doppler of portal circulation	Every 12 months
	MRCP	In case of progression of biliary disease
	ERCP	For therapeutic interventions or tissue sampling for evaluation of dysplasia or cholangiocarcinoma
Elastography	Liver stiffness	Optional

Abbreviations: AFP, alpha-fetoprotein; ALT, alanine aminotransferase; ANA, antinuclear antibody; AST, aspartate aminotransferase; anti-LC1, antiliver cytosol type 1 antibody; anti-LKM-1, antiliver kidney microsomal type 1 antibody; anti-SLA, antisoluble liver antigen; anti-SMA, antismooth muscle antibody; CRP, C-reactive protein; ERCP, endoscopic retrograde cholangiopancreatography; GGT, gamma-glutamyl transferase; IgG, immunoglobulin G; INR, international normalized ratio; MRCP, magnetic resonance cholangiopancreatography; PSC, primary sclerosing cholangitis; TSB, total serum bilirubin.

Section 5.4.2). In the case of biochemical deterioration of cholestasis, an MRCP is recommended to assess the presence of a dominant stricture.

5.4 | PSC and cancer risk

In adults with PSC-IBD, the inherent risk of colorectal carcinoma (CRC) and CCA plays a significant role in both morbidity and mortality. The likelihood of these malignancies manifesting before the 18th year of life is extremely low (see Sections 5.4.1 and 5.4.2). There is a significant difference between pediatric and adult practice regarding cancer screening. Due to the lower risk and practicalities for organizing endoscopy, yearly surveillance in children with PSC-IBD is less self-evident. Once the transfer

to adult-oriented care has taken place, annual surveillance colonoscopy with biopsies is recommended in all adults with PSC-IBD.⁸ It is important to sensitively educate children and adolescents with PSC-IBD and their families about their elevated cancer risk at the appropriate time and prepare them for this more intense screening strategy.

5.4.1 | CRC surveillance

Position statement 5

In children with PSC-IBD the risk of high-grade dysplasia or colorectal cancer is greater than in isolated IBD, however, the absolute risk of

being diagnosed with colorectal cancer before the age of 18 is very low.

Moderate certainty | Agreement: 100%

Practice point

Surveillance colonoscopy should be considered in children with PSC-IBD and the following risk factors of colorectal cancer:

- persistent active colonic inflammation, or
- longstanding colitis (≥ 8 years), or
- a family history of colorectal cancer in a first-degree relative < 50 years.

5.4.1.1 | Evidence

The risk of high-grade dysplasia and CRC in adult patients with PSC is 4–5 times greater compared to patients with IBD without PSC.⁷ Leading adult guidelines recommend to perform yearly surveillance colonoscopy from the time of diagnosis of PSC. Dye-based or virtual chromoendoscopy with targeted biopsies is increasingly recommended, although it has not been proven that these techniques are superior to high-definition endoscopy without chromoendoscopy and random biopsies. Children with PSC-IBD seem to develop CRC at similar rates as adults, with approximately 5% affected 10 years after PSC diagnosis (see Figure 5). However, the absolute risk of being diagnosed with CRC before the age of 18 is low (0.2%).⁴²

It is important to note that there is a lack of substantial evidence supporting the effectiveness of surveillance colonoscopy. Current practice guidelines offer little guidance regarding the age to start cancer surveillance, the colonoscopy technique, and the screening interval (as outlined in Supporting Information:

Table S16). As a result, there is a large variation in clinical practice, as was shown in a recently published survey among Dutch pediatric gastroenterologists.⁴³

5.4.1.2 | Practical guidance

Recommendations for malignancy screening in adults with PSC-IBD are outlined in the ECCO Malignancy in IBD Guidelines.⁴⁴ Strictly following these recommendations for children and adolescents with PSC-IBD would lead to unacceptably frequent procedures with no treatment consequences. We recommend including only those patients with a high-risk profile in a surveillance colonoscopy program. This high-risk profile includes individuals with ongoing colonic inflammation (fecal calprotectin > 150 $\mu\text{g/g}$ or histologic active disease), a history of long-standing colitis (8 years or more), or a family history of CRC in a first-degree relative under the age of 50. When one of these risk factors is present, surveillance colonoscopy is usually conducted every other year. Pediatricians involved in CRC screening are advised to familiarize themselves with the “best practice advice” to enhance the quality of screening.^{45,46} Some centers involve endoscopists from the adult GI department in CRC screening for adolescents.

5.4.2 | CCA surveillance

In adults CCA can occur any time during the disease course, but the most consistent risk factor for CCA is older age. CCA is rarely diagnosed in the pediatric population or in those with small-duct PSC.⁷ In a multicenter, international cohort of 781 children with PSC, eight were diagnosed with CCA at an age range of 15–18 years, a median of 6 years after initial PSC diagnosis.⁵ Three patients with metastatic cancer died




FIGURE 5 Summary of CRC-free survival in patients with PSC and IBD at 5 and 10 years after diagnosis. CRC, colorectal cancer; IBD, inflammatory bowel disease; PSC, primary sclerosing cholangitis.

under palliative care within 5 months of CCA diagnosis. Five patients with localized disease underwent surgery or liver transplantation, with all five alive at a median follow-up of 2.5 years after diagnosis. All patients who developed CCA had PSC without AIH overlap. Routine CCA surveillance for patients with PSC under 18 years of age is not recommended. New onset jaundice with a cholestatic biochemical profile should prompt MRCP and carbohydrate antigen 19.9 (CA 19.9) testing, and findings of distal common bile duct strictures will require ERCP with biliary brushings for cytology. CA 19.9 testing is not suitable for surveillance purposes due to its insufficient accuracy.^{7,8}

5.5 | Referral criteria for liver transplantation

As mentioned in Section 5.1, approximately 50% of the patients with PSC eventually require a liver transplantation. A population-based study from the Netherlands among adults showed a transplant-free interval of 15–20 years after the diagnosis of PSC,⁴⁷ whereas in specialized hepatology centers transplant-free survival is usually shorter (10–15 years), due to a different case mix.⁴⁸ Although liver transplantation is more likely to occur after transfer to adult services, assessment for liver transplantation should be considered in patients with PSC and intractable severe pruritus, recurrent bacterial cholangitis, portal hypertension or relevant bile-duct strictures. This advice is consistent with a recently published consensus guideline of the European Society of Organ Transplantation.⁴⁹ Prioritization for liver transplantation was traditionally based on the model for end-stage liver disease (MELD) score that included three blood parameters: international normalized ratio, creatinine, and bilirubin. The current revised MELD also includes serum sodium. The MELD-score, which is a proxy for the severity of hepatic dysfunction, underestimates the severity of PSC complicated by recurrent cholangitis.^{50,51} Consequently, waiting times on transplant lists for patients with PSC have been typically longer and outcomes worse than those for other forms of liver disease.^{50,51} We suggest referral of children with newly diagnosed PSC–IBD to a pediatric hepatologist with expertise in liver transplantation at an early stage. Thereafter, joint care with the IBD team and the local pediatric hepatologist should continue. This advice is in accordance with the latest ECCO guideline on extraintestinal manifestations in IBD.¹⁴

Recurrence of PSC (rPSC) after liver transplantation is not uncommon. The International Pediatric PSC Consortium described a cohort of 140 children with PSC who were transplanted before 18 years of age. rPSC occurred in 27% of the patients in the first 5 years after liver transplantation. IBD was more prevalent in

the rPSC group as compared to those without recurrence.⁵² In an international retrospective adult cohort study of 531 liver transplant recipients, those with increased IBD activity after transplantation had a greater risk of rPSC compared to those with stable IBD (hazard ratio 1.7, 95% confidence interval 1.1–2.8; $p = 0.021$).⁵³ A systematic review published in 2018 suggested a protective role of colectomy when carried out before or at the time of liver transplantation.⁵⁴ The quality of the included papers was too low to routinely recommend colectomy for rPSC prevention, but colectomy may be considered on a case-by-case basis, especially if IBD has been difficult to manage.

6 | THERAPEUTIC STRATEGIES

Currently, there is no medication available to slow down PSC disease progression, although some improvement in liver biochemistry may occur. We present here the most frequently used therapies.

6.1 | UDCA

Position statement 6

UDCA may be prescribed at doses of 15–20 mg/kg/day. Despite evidence of improvement of liver enzymes, its long-term effect on disease progression has not been demonstrated.

Low certainty | Agreement: 90% | Weak recommendation

Practice point

Consider a 6-months therapeutic trial of UDCA, either immediately after PSC diagnosis or when spontaneous normalization of GGT does not occur in the first 6 months postdiagnosis. Continue UDCA treatment if there is a meaningful reduction or normalization of GGT or improvement of symptoms.

6.1.1 | Evidence

UDCA is a hydrophilic bile acid with anticholestatic, anti-inflammatory, and antifibrotic actions. It is widely used in the treatment of PSC in children and adolescents. Similar to adult studies, pediatric research shows an improvement in liver biochemistry after introducing the drug (Supporting Information: Table S13). However, there is currently no evidence that UDCA slows down the progression of PSC.

There is no agreement on what constitutes a minimum effective dosage. In a retrospective multicenter study including 46 international centers the median UDCA dose administered to children with PSC was 15 mg/kg/day (interquartile range 15–19).⁵⁵ Notably, an adult trial with high-dose UDCA (28–30 mg/kg/day) was prematurely terminated for its negative impact on esophageal varices, time-to-transplantation and survival, despite improvement in liver biochemistry.^{56,57} Provided that UDCA is adequately dosed, its long-term safety profile is excellent. Up to 10% of users report loose stools and it has few interactions with other drugs.⁵⁸

A prospective study among 22 children with PSC evaluated UDCA withdrawal. The participants had been on UDCA maintenance therapy with normal biochemistry. The UDCA dose was reduced by 50% for 4 weeks, after which it was discontinued for 8 weeks. A third of the participants showed a GGT increase above 100 IU/L during withdrawal with a normalization of GGT upon UDCA reinstitution. In another third GGT and ALT remained in the normal range during withdrawal. The remaining group had a mild increase in GGT up to a maximum of 100 IU/L. The significance of a biochemical flare is unclear and underscores the need for a more reliable biomarker of disease progression.⁵⁹

6.1.2 | Practical guidance

Despite lack of evidence of long-term effects, UDCA is prescribed long-term in over 80% of children with PSC across many countries.⁵ Many experts feel there is at least some role for a 6-month therapeutic trial of UDCA, with continuation of the drug in patients with a substantial biochemical response,⁶⁰ while others recommend monitoring GGT for 6 months before initiating UDCA because levels can spontaneously normalize.⁷ Patients with persistently elevated GGT levels can then be considered for UDCA treatment at 15–20 mg/kg/day, and treatment can be continued if there is a meaningful reduction or normalization of GGT or improvement of symptoms with 6 months of treatment.

6.2 | Oral vancomycin

Position statement 7

Oral vancomycin may be prescribed for a potential improvement in liver biochemistry as well as bowel inflammation. Its long-term effect on disease progression has not been demonstrated.

Low certainty | Agreement: 85% | Weak recommendation

6.2.1 | Evidence

Dysregulation of the gut microbiome has been postulated as one facet of the intricate pathogenesis of PSC, and modulation of the gut microbiome may have a favorable effect on the course of PSC.⁶¹ Oral vancomycin is the most extensively investigated antibiotic in the context of childhood-onset PSC. It is poorly absorbed from the gut and may diminish the concentration of bacteria in the gut, or their potentially harmful metabolites in the portal circulation. Prospective and retrospective studies conducted in children (Supporting Information: Table S14) have demonstrated improvements in liver biochemistry following oral vancomycin treatment. In a small-scale retrospective study, liver function tests improved or normalized in 12 out of 17 children with PSC who received oral vancomycin for a minimum of 3 months. Moreover, a significant portion of these children also achieved biochemical endoscopic and histologic remission of colitis.⁶² A article from the Pediatric PSC Consortium that was recently published is worth mentioning in this context as well.⁶³ A group of 70 children with PSC-IBD, who in addition to conventional IBD medication also received oral vancomycin for at least 3 months, was matched with a group of 210 children with PSC-IBD who were not exposed to vancomycin. The probability of reaching IBD clinical remission was higher in the vancomycin-treated subgroup.

Two small prospective studies (respectively 45 and 14 participants) showed significant changes in liver biochemistries in the majority of children enrolled.^{64,65} In the largest retrospective case-control study so far, 264 children diagnosed with PSC were matched 1:1:1 to those treated with vancomycin, UDCA, or observed without therapy.⁶⁶ After 1 year, no discernible improvement in outcomes was observed across the treatment groups. Specifically, GGT levels normalized in 53%, 49%, and 52%, while the liver fibrosis stage improved in 20%, 13%, and 18% among the vancomycin, UDCA, and observation-only groups, respectively. However, liver fibrosis worsened in 11%, 29%, and 18%, respectively. Furthermore, the 5-year likelihood of liver transplant listing stood at 21%, 10%, and 12%, respectively. Notably, spontaneous normalization of liver biochemistry was frequently observed in children who did not receive therapy, particularly those with a mild phenotype and minimal fibrosis.

6.2.2 | Practical guidance

There are many uncertainties regarding the use of oral vancomycin in patients with PSC-IBD, including optimal dosage, efficacy across different stages of fibrosis and the length of safe treatment. In an open-label prospective clinical trial children with PSC were treated at

a dose of 50 mg/kg/day divided into three times per day if weight was <30 kg, and at a dose of 500 mg three times a day if weight was ≥30 kg.⁶⁴ Until more data is available, we recommend judicious use of the drug. Patients who experience a rapid reduction and subsequent normalization of GGT within 12 weeks may be considered for longer-term therapy with periodic evaluation for vancomycin-resistant enterococcus.⁶⁰

6.3 | Corticosteroids and antimetabolites

Position statement 8

In children with PSC–IBD and biochemical, serological, and histological features of AIH, the use of corticosteroids and antimetabolites may suppress immune-mediated hepatitis.

In the absence of convincing AIH features, the use of corticosteroids and antimetabolites is not indicated to manage PSC.

Low certainty | Agreement: 95% | Weak recommendation

6.3.1 | Evidence

Some studies address long-term use of corticosteroids and antimetabolites to control autoimmunity in patients with PSC and features of AIH based on biochemical improvement or reduction of parenchymal inflammation in the liver. However, the bile duct disease progresses in about 50% of cases, leading to end-stage liver disease requiring transplantation more frequently than in isolated AIH.²¹

Maintenance therapy with corticosteroids, usually between 5 and 7.5 mg per day,²¹ daily, may cause adrenal suppression.⁶⁷ Also, the effect of corticosteroids on growth are well documented. However, a recent retrospective study including 74 children with autoimmune liver disease (PSC not mentioned separately) treated with a daily maintenance dose of steroids for a median duration of 11 years (interquartile range 8.4–13.8) showed mean z-scores for weight, height, and body mass index within the normal range.⁶⁸

6.3.2 | Practical guidance

There are no convincing data that use of corticosteroids and antimetabolites improves long-term outcomes in children with PSC and overlapping features of AIH. However, in children with biochemical, serological, as

well as histological features of AIH it is reasonable to trial corticosteroids and antimetabolites using a similar approach as in AIH.²¹ The prednisolone starting dose is weight-dependent and should be tapered once aminotransferases start to decrease, but not later than 4 weeks after initiation. Frequently, a low-dose prednisolone maintenance dose (2.5–5 mg daily) is necessary to keep aminotransferase levels in range.

A steroid-sparing antimetabolite such as azathioprine or mercaptopurine is usually introduced a few weeks after the initiation of prednisolone, or when aminotransferases stop decreasing. A recent adult study on the use of mycophenolate mofetil with prednisolone as induction therapy for AIH showed better outcomes than the combination prednisolone–azathioprine.⁶⁹ However, patients with PSC and IBD were excluded from participating in this study. Currently, there is insufficient evidence to recommend the use of mycophenolate mofetil in children with PSC–IBD and convincing AIH features. There may be a role for repeated liver biopsy in the patient with PSC–AIH who continues to display elevated liver enzymes to help clarify whether there is ongoing inflammation in the liver parenchyma, or only biliary damage. In the latter situation withdrawal of corticosteroids and antimetabolites rather than long-term use could be considered. Immunosuppressive therapy is not recommended for PSC without AIH features.

6.4 | Biologicals

In a retrospective analysis of 141 adults with PSC–IBD, antitumor necrosis factor- α (anti-TNF- α) agents were primarily prescribed for the IBD indication. Anti-TNF- α medication was effective in reducing intestinal inflammation, although not as effective as in patients with non-PSC IBD, with no specific safety signals.⁷⁰ One double-blind, randomized controlled study in adult patients with UC and PSC was specifically designed to evaluate the efficacy of infliximab on PSC symptom relief, liver biochemistry and histology. When an interim analysis revealed that infliximab did not provide a significant treatment benefit on PSC symptoms and liver histology, the study was halted prematurely, due to the disproportionately high risk to participants (resulting from liver biopsies at baseline and at 26 weeks).⁷¹

In a spin-off study of the Pediatric PSC Consortium, a retrospective multicenter research registry, 37 children with PSC–IBD on vedolizumab therapy were followed for a year. IBD activity improved on vedolizumab at rates similar to non-PSC IBD, but hepatobiliary outcomes did not improve.⁷²

Currently, there is insufficient evidence to recommend one biological treatment over another to stop progression of PSC. For decisions around IBD treatment, we refer to the latest ECCO–ESPGHAN treatment guidelines for Crohn's disease and UC.^{73,74} The effect of IBD activity on PSC progression is unclear.




FIGURE 6 Summary flowchart of diagnosis, monitoring and treatment. The numbers between brackets refer to the relevant sections. AIH, autoimmune hepatitis; ALT, alanine aminotransferase; CRC, colorectal cancer; FC, fecal calprotectin; GGT, gamma-glutamyltransferase; IBD, inflammatory bowel disease; MRCP, magnetic resonance cholangiopancreatography; PSC, primary sclerosing cholangitis; UDCA, ursodeoxycholic acid; ULN, upper limit of normal.

However, as in non-PSC IBD, mucosal healing should be the treatment goal in PSC-IBD, as it improves long-term outcomes of IBD and possibly mitigates the increased risk of CRC.

6.5 | ERCP and strictures

Practice point

Children with PSC, relevant bile-duct strictures and cholestatic symptoms should be assessed for liver transplantation. When their symptoms are likely to improve following biliary intervention, ERCP can be considered. Referral to a centre with expertise in pediatric interventional ERCP is highly recommended.

As discussed in Section 4.3 in connection with diagnostic imaging, MRCP has supplanted the role of ERCP to identify focal narrowing and ballooning of bile ducts. ERCP is now mainly a means to manage dominant biliary strictures, or to sample tissue for suspected CCA.^{8,12}

The possible presence of a dominant stricture in the bile ducts should be investigated when a patient with known PSC-IBD suddenly develops cholestatic symptoms (including jaundice, bacterial cholangitis, itching, or pain in the right upper abdomen) and a cholestatic biochemical profile. When narrowing in the extrahepatic or first-order intrahepatic ducts is observed on MRCP, with or without upstream dilatation, there is an indication for ERCP. When ERCP confirms a dominant stricture, primary treatment should consist of balloon dilation. Dilation followed by stent placement does not lead to better outcomes but is associated with significant adverse events, such as bacterial cholangitis and pancreatitis.⁷⁵

Based on the invasiveness of the procedure, the considerable exposure to radiation and its potential for severe complications, ERCP in children is only justified in the situation of a favorable benefit-risk ratio, such as a high suspicion of a dominant stricture or malignancy on MRCP.⁷⁶ Referral to a centre with expertise in pediatric interventional ERCP is highly recommended.⁸

6.6 | Colectomy

Colectomy in adults with IBD, specifically UC, alongside PSC, presents a complex clinical scenario. There are data supporting that IBD (and its inflammatory activity) is important for PSC progression. Transplant- and cancer-free survival is poorer in patients with UC when compared to patients with no IBD or

Crohn's disease.⁴⁷ Furthermore, the longer time-to-transplantation in adults who underwent colectomy before diagnosis of PSC compared to those with their colon in situ supports the notion that gut inflammation may affect PSC progression.⁷⁷ In adults with UC and PSC, the contemplation of colectomy often arises due to the heightened risk of CRC associated with PSC. The decision to proceed with colectomy depends on various factors, including disease severity, response to medical treatments, and the presence of dysplasia or cancer. While evidence indicates that colectomy can reduce the risk of CRC, it does not stop PSC progression, and individuals may still require liver transplantation in the future. Colectomy may also offer relief from UC-related symptoms, but this surgical intervention carries inherent risks and significant lifestyle implications.

The use of colectomy in adults with IBD and PSC has been debated as a pre-emptive measure to reduce the risk of recurrent PSC after liver transplantation. Currently, the European Association for the Study of the Liver advises against performing a pre-emptive colectomy before liver transplantation in adults with PSC-IBD. However, in refractory cases, colectomy should be considered at a slightly lower threshold than in individuals with PSC who have not yet undergone a liver transplant.⁸

In children with IBD and PSC, colectomy is less frequently considered compared to adults.³³ The appropriateness of colectomy in children with PSC-IBD requires individualized assessment taking into account their growth, development and overall quality of life. Decisions regarding colectomy should be made in consultation with a multidisciplinary team.

7 | CONCLUSION

The aim of this ESPGHAN position paper is to guide clinicians' decisions with the best evidence available to establish the right diagnosis, follow-up and treatment strategies for children with IBD-associated PSC. Summary flowcharts are shown in Figure 6. It is up to every clinician to make adaptations to local regulations and to the patient's individual characteristics and needs.

8 | KNOWLEDGE GAPS

The number of patients with PSC is increasing along with the substantial increase in the incidence of IBD and autoimmune diseases in general during the past decade.⁷⁸ However, the absolute number of patients with PSC in each pediatric center is still low pointing out the need for multicenter prospective studies to increase knowledge of the pediatric phenotype of PSC. To date,

we lack the means to identify the young patients at risk for the development of end-stage liver disease and CCA, and we do not have medications to stop disease progression. UDCA is widely prescribed, and it may have an effect in normalizing liver biochemistry, but there is no evidence that it indeed improves the outcomes of the patients. This is true for other drugs with some reported benefit on the levels of liver enzymes as well, such as vancomycin. Also, it is an unanswered question whether PSC with features of AIH comprises a disease subtype of its own with different outcomes or, as recently suggested, a stage on the disease continuum between AIH and PSC. As most children with PSC have concomitant IBD, they are at risk for developing CRC. Reassuringly, the absolute risk for colon cancer in PSC is low at a young age. Unfortunately, we do not know how to stratify those at substantial risk. Adult guidelines on PSC-IBD recommend annual surveillance endoscopies that seem overcautious for young patients and there is a need for guidelines adjusting the recommendations according to the age and duration of disease history in childhood-onset PSC. Disease progression in childhood-onset PSC is individual and the optimal follow-up with MRCP is to be defined. Follow-up is needed for timely detection of disease progression which may be an indication for ERCP and brush cytology to diagnose dysplasia that precedes the development of cancer. Patient selection for ERCP remains a challenge. Future research will hopefully identify novel prognostic biomarkers to better characterize the subtype of childhood-onset PSC. Such markers would aid in focusing the most intensive follow-up on those patients with the highest risk for adverse outcomes and spare the other patients from invasive investigations (Box 3).

BOX 3. Suggestions for future studies based on current knowledge gaps

- Prospective international registry using agreed clinical definitions, combined with the collection of biological samples (including -80°C frozen stool, urine, blood, and DNA) to assess the natural history of PSC-IBD and explore new biomarkers.
- Nested diagnostic study to predict the presence of colonic inflammation in children diagnosed with isolated PSC.
- Controlled study of vancomycin use to evaluate short and medium-term outcomes.

AFFILIATIONS

¹Department of Paediatric Gastroenterology, Hepatology, and Nutrition, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

²Children's Hospital, University of Helsinki and HUS, Helsinki, Finland

³Department of Paediatric Gastroenterology, and Nutrition, Royal Hospital for Children and Young People, Edinburgh, UK

⁴Sapienza University of Rome - Umberto I Hospital, Rome, Italy

⁵Department of Radiology, King's College Hospital, School of Biomedical Engineering and Imaging Sciences, King's College London, London, UK

⁶Children's Health Ireland and University College Dublin, Dublin, Ireland

⁷Division for Pediatric Gastroenterology and Hepatology, Department of Pediatric Kidney, Liver, and Metabolic Diseases, Hannover Medical School, Hannover, Germany

⁸General Paediatrics and Neonatology, Gastroenterology, Justus Liebig University Giessen, Giessen, Germany

⁹University of Utah and Intermountain Healthcare Primary Children's Hospital, Salt Lake City, Utah, USA

¹⁰Faculty of Medicine, IBD Centre, SickKids Hospital, University of Toronto, Toronto, Ontario, Canada

¹¹Children's Hospital Zagreb, University of Zagreb Medical School, Zagreb, Croatia

¹²Pediatric Hepatology, Gastroenterology and Transplantation, Hospital Papa Giovanni XXIII, Bergamo, Italy

¹³Department of Paediatric Gastroenterology, Great Ormond Street Hospital, London, UK

¹⁴Department of Pediatric Surgery, The New Children's Hospital, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

¹⁵Department of Paediatric Gastroenterology, Erasmus University Medical Center Sophia Children's Hospital, Rotterdam, The Netherlands

¹⁶Department of Pathology, ASST Bergamo Est, Bergamo, Italy

¹⁷Helsinki University and Helsinki University Hospital HUS, Abdominal Center, Helsinki, Finland

¹⁸Paediatric Liver, GI and Nutrition Service, King's College Hospital, London, UK

¹⁹Meyer Children's Hospital IRCCS, Florence, Italy

ACKNOWLEDGMENTS

We thank Hanna van Rheenen for creating the illustration in Figure 4. This project was funded by the European Society for Pediatric Gastroenterology Hepatology and Nutrition.

CONFLICT OF INTEREST STATEMENT

Patrick F. van Rheenen: Research Grant—European Crohn & Colitis Organisation. Leadership Role—Chair of Dutch PIBD Guideline (2022-2024). Receipt of Donated Equipment—BÜHLMANN Laboratories AG. Kaija-Leena Kohlo: Grants or Contracts: Helsinki University Hospital Research Fund—ongoing research projects on fecal microbiome; Pediatric Research Foundation (Finland)—ongoing research project on pediatric IBD Consultancy fees: Abbvie; Blocohex. Richard K. Russell: Grant/Contract: Nestle Health Science—ongoing; Ferring—past. Consultancy fees: Celtrion and Lily. Participated in Data Monitoring Committee: OCEAN Trial. Medical Writing: Pfizer.

Honoraria Lecture: Janssen. Payment for Meeting attendance: Abbvie and Celltrion. Marina Aloï: Honoraria for Lectures—Takeda, Nestle. Safety Monitoring Board—Pfizer. Annamaria Deganello: none. Séamus Hussey: Research Grant—Janssen. Norman Jung: Speaker at annual meeting/HEPCOM member—ESPGHAN. Data Monitoring Committee—ERN Rare Liver member for Rare Liver Registry. Leadership Roles—ERN Rare Liver, ERN Transplant Child. Jan De Laffolie: Consulting Fees—Mirum, Takeda, Sanofi. Lecture Honoraria—Mirum, Baxter, Sanofi. Safety monitoring board member—Sanofi. Advisor to patient organizations—DCCV, DZG. Mark R. Deneau: none. Emer Fitzpatrick: Grant/Contract—Health Research Board, Ireland (author's institution). Board member—ESPGHAN. Anne M. Griffiths: Grants/Research Support—Abbvie. Member of advisory board—Abbvie, Janssen, Lily. Honoraria for Lectures—Abbvie, Janssen, Nestle. Honoraria for Consultation—Abbvie, Amgen, Bristol Meyers Squib, Pfizer, Merck, Janssen, Lily. Iva Hojsak: Consultancy Fees—Abbot and Biocodex. Honoraria for Lectures—Sandoz, Nestle, BioGaia, Hipp, GM Pharma. Emanuele Nicastro: Served as member on advisory board—Mirum Pharmaceuticals. Andreia Nita: none. Mikko Pakarinen: none. Amanda Ricciuto: none. Lissy de Ridder: Collaboration (such as involved in industry-sponsored studies, investigator-initiated study, consultancy) with Abbvie, Eli Lilly, Takeda, Janssen, Medtronic and Pfizer. Aurelio Sonzogni: none. Andrea Tenca: none. Marianne Samyn: none. Giuseppe Indolfi: Payment/honorarium Kedrion Pharma 2020.

ORCID

Patrick F. van Rheenen  <http://orcid.org/0000-0003-3867-2665>

REFERENCES

1. Pusateri AJ, Kim SC, Dotson JL, et al. Incidence, pattern, and etiology of elevated liver enzymes in pediatric inflammatory bowel disease. *J Pediatr Gastroenterol Nutr.* 2015;60(5):592-597. doi:10.1097/MPG.0000000000000672
2. Daniluk U, Kwiatek-Sredzinska K, Jakimiec P, Daniluk J, Czajkowska A, Lebensztejn D. Liver pathology in children with diagnosed inflammatory bowel Disease: a single center experience. *J Clin Med.* 2021;10(22):5359. doi:10.3390/jcm10225359
3. Chandrakumar A, Loeppky R, Deneau M, El-Matary W. Inflammatory bowel disease in children with elevated serum gamma glutamyltransferase levels. *J Pediatr.* 2019;215:144-151.e3. doi:10.1016/j.jpeds.2019.07.065
4. Joosse ME, Haisma SM, Sterk MFM, et al. Disease progression in paediatric- and adult-onset sclerosing cholangitis: results from two independent Dutch registries. *Liver Int.* 2019;39(9):1768-1775. doi:10.1111/liv.14159
5. Deneau MR, El-Matary W, Valentino PL, et al. The natural history of primary sclerosing cholangitis in 781 children: a multicenter, international collaboration. *Hepatology.* 2017;66(2):518-527. doi:10.1002/hep.29204
6. Chapman MH, Thorburn D, Hirschfield GM, et al. British Society of Gastroenterology and UK-PSC guidelines for the diagnosis and management of primary sclerosing cholangitis. *Gut.* 2019;68(8):1356-1378. doi:10.1136/gutjnl-2018-317993
7. Bowlus CL, Arrivé L, Bergquist A, et al. AASLD practice guidance on primary sclerosing cholangitis and cholangiocarcinoma. *Hepatology.* 2022;77(2):1-44. doi:10.1002/hep.32771
8. Chazouilleres O, Beuers U, Bergquist A, et al. EASL clinical practice guidelines on sclerosing cholangitis. *J Hepatol.* 2022;77(3):761-806. doi:10.1016/j.jhep.2022.05.011
9. Goyal A, Hyams JS, Lerer T, et al. Liver enzyme elevations within 3 months of diagnosis of inflammatory bowel disease and likelihood of liver disease. *J Pediatr Gastroenterol Nutr.* 2014;59(3):321-323. doi:10.1097/MPG.0000000000000409
10. Ordonez F, Lacaille F, Canioni D, et al. Pediatric ulcerative colitis associated with autoimmune diseases: a distinct form of inflammatory bowel disease. *Inflamm Bowel Dis.* 2012;18(10):1809-1817. doi:10.1002/ibd.22864
11. Valentino PL, Feldman BM, Walters TD, et al. Abnormal liver biochemistry is common in pediatric inflammatory bowel disease: prevalence and associations. *Inflamm Bowel Dis.* 2015;21(12):2848-2856. doi:10.1097/MIB.0000000000000558
12. Ponsioen CY, Assis DN, Boberg KM, et al. Defining primary sclerosing cholangitis: results from an International Primary Sclerosing Cholangitis Study Group Consensus Process. *Gastroenterology.* 2021;161(6):1764-1775.e5. doi:10.1053/j.gastro.2021.07.046
13. van Rheenen PF. Managing abnormal liver tests in children with inflammatory bowel disease. *Curr Opin Pediatr.* 2021;33(5):521-529. doi:10.1097/MOP.0000000000001045
14. Gordon H, Burisch J, Ellul P, et al. ECCO guidelines on extra-intestinal manifestations in inflammatory bowel disease. *J Crohns Colitis.* 2023;18(1):1-37. doi:10.1093/ecco-jcc/jjad108
15. Hensel KO, Kyrana E, Hadzic N, et al. Sclerosing cholangitis in pediatric inflammatory bowel disease: early diagnosis and management affect clinical outcome. *J Pediatr.* 2021;238:50-56.e3. doi:10.1016/j.jpeds.2021.07.047
16. Shamberg L, Vaziri H. Hepatotoxicity of inflammatory bowel disease medications. *J Clin Gastroenterol.* 2018;52(8):674-684. doi:10.1097/MCG.0000000000001084
17. LiverTox. *Clinical and Research Information on Drug-Induced Liver Injury.* National Institute of Diabetes and Digestive and Kidney Diseases; 2012.
18. Hoofnagle JH, Björnsson ES. Drug-induced liver injury—types and phenotypes. *N Engl J Med.* 2019;381(14):1395-1396. doi:10.1056/NEJMc1911063
19. Ghersin I, Khateeb N, Katz LH, Daher S, Shamir R, Assa A. Comorbidities in adolescents with inflammatory bowel disease: findings from a population-based cohort study. *Pediatr Res.* 2020;87(7):1256-1262. doi:10.1038/s41390-019-0702-3
20. Di Giorgio A, Hadzic N, Dhawan A, et al. Seamless management of juvenile autoimmune liver disease: long-term medical and social outcome. *J Pediatr.* 2020;218:121-129.e3. doi:10.1016/j.jpeds.2019.11.028
21. Mieli-Vergani G, Vergani D, Baumann U, et al. Diagnosis and management of pediatric autoimmune liver disease: ESPGHAN hepatology committee position statement. *J Pediatr Gastroenterol Nutr.* 2018;66(2):345-360. doi:10.1097/MPG.0000000000001801
22. Vergani D, Alvarez F, Bianchi FB, et al. Liver autoimmune serology: a consensus statement from the committee for autoimmune serology of the International Autoimmune Hepatitis Group. *J Hepatol.* 2004;41(4):677-683. doi:10.1016/j.jhep.2004.08.002
23. Patil K, Ricciuto A, Alsharief A, et al. Magnetic resonance cholangiopancreatography severity predicts disease outcomes in pediatric primary sclerosing cholangitis: a reliability and

- validity study. *Hepatology Commun.* 2020;4(2):208-218. doi:10.1002/hep4.1454
24. Miloh T, Arnon R, Shneider B, Suchy F, Kerkar N. A retrospective single-center review of primary sclerosing cholangitis in children. *Clin Gastroenterol Hepatol.* 2009;7(2):239-245. doi:10.1016/j.cgh.2008.10.019
 25. Noble-Jamieson G, Heuschkel RB, Torrente F, Hadzic N, Zilbauer M. Colitis-associated sclerosing cholangitis in children: a single centre experience. *J Crohn's Colitis.* 2013;7(10):e414-e418. doi:10.1016/j.crohns.2013.01.016
 26. Rojas CP, Bodicharla R, Campuzano-Zuluaga G, Hernandez L, Rodriguez MM. Autoimmune hepatitis and primary sclerosing cholangitis in children and adolescents. *Fetal Pediatr Pathol.* 2014;33(4):202-209. doi:10.3109/15513815.2014.898721
 27. Venkatesh SK, L.Welle C, H.Miller F, et al. Correction to: reporting standards for primary sclerosing cholangitis using MRI and MR cholangiopancreatography: guidelines from MR Working Group of the International Primary Sclerosing Cholangitis Study Group. *Eur Radiol.* 2022;32(4):2860. doi:10.1007/s00330-021-08333-7
 28. Bateman AC, Hübscher SG. Cytokeratin expression as an aid to diagnosis in medical liver biopsies. *Histopathology.* 2010;56(4):415-425. doi:10.1111/j.1365-2559.2009.03391.x
 29. Ricciuto A, Kamath BM, Griffiths AM. The IBD and PSC phenotypes of PSC-IBD. *Curr Gastroenterol Rep.* 2018;20(4):16. doi:10.1007/s11894-018-0620-2
 30. de Vries AB, Janse M, Blokzijl H, et al. Distinctive inflammatory bowel disease phenotype in primary sclerosing cholangitis. *World J Gastroenterol.* 2015;21(6):1956-1971. doi:10.3748/wjg.v21.i6.1956
 31. Franke M, Geiß A, Greiner P, et al. The role of endoscopy in pediatric gastrointestinal bleeding. *Endosc Int Open.* 2016; 04(9):E1011-E1016. doi:10.1055/s-0042-109264
 32. Van de Vijver E, Heida A, Ioannou S, et al. Test strategies to predict inflammatory bowel disease among children with non-bloody diarrhea. *Pediatrics.* 2020;146(2):e20192235. doi:10.1542/peds.2019-2235
 33. Ricciuto A, Hansen BE, Ngo B, et al. Primary sclerosing cholangitis in children with inflammatory bowel diseases is associated with milder clinical activity but more frequent subclinical inflammation and growth impairment. *Clin Gastroenterol Hepatol.* 2020;18(7):1509-1517.e7. doi:10.1016/j.cgh.2019.08.048
 34. Bjarnason I, Hayee B, Pavlidis P, et al. Contrasting pattern of chronic inflammatory bowel disease in primary and autoimmune sclerosing cholangitis. *EBioMedicine.* 2015;2(10):1523-1527. doi:10.1016/j.ebiom.2015.08.041
 35. Pavlidis P, Joshi D, El Sherif Y, et al. Faecal calprotectin is a surrogate marker of biliary inflammation in primary sclerosing cholangitis associated inflammatory bowel disease. *Frontline Gastroenterol.* 2022;13(6):497-502. doi:10.1136/flgastro-2021-102053
 36. Ricciuto A, Fish J, Carman N, et al. Symptoms do not correlate with findings from colonoscopy in children with inflammatory bowel disease and primary sclerosing cholangitis. *Clin Gastroenterol Hepatol.* 2018;16(7):1098-1105.e1. doi:10.1016/j.cgh.2018.01.020
 37. Maaser C, Sturm A, Vavricka SR, et al. ECCO-ESGAR guideline for diagnostic assessment in IBD part 1: initial diagnosis, monitoring of known IBD, detection of complications. *J Crohn's Colitis.* 2019;13(2):144-164K. doi:10.1093/ecco-jcc/ijy113
 38. Ricciuto A, Kamath BM, Hirschfield GM, Trivedi PJ. Primary sclerosing cholangitis and overlap features of autoimmune hepatitis: a coming of age or an age-ist problem. *J Hepatol.* 2023;79(2):567-575. doi:10.1016/j.jhep.2023.02.030
 39. Deneau MR, Mack C, Abdou R, et al. Gamma glutamyl-transferase reduction is associated with favorable outcomes in pediatric primary sclerosing cholangitis. *Hepatol Commun.* 2018;2(11):1369-1378. doi:10.1002/hep4.1251
 40. Berhane B, van Rheenen PF, Verkade HJ. Gamma-glutamyl transferase and disease course in pediatric-onset primary sclerosing cholangitis: a single-center cohort study. *Health Sci Rep.* 2023;6(2):e1086. doi:10.1002/hsr2.1086
 41. Deneau MR, Mack C, Perito ER, et al. The sclerosing cholangitis outcomes in pediatrics (SCOPE) index: a prognostic tool for children. *Hepatology.* 2021;73(3):1074-1087. doi:10.1002/hep.31393
 42. Olén O, Askling J, Sachs M, et al. Childhood onset inflammatory bowel disease and risk of cancer: a Swedish nationwide cohort study 1964-2014. *BMJ.* 2017;358:j3951. doi:10.1136/bmj.j3951
 43. Jagt JZ, van Schie DA, Benninga MA, van Rheenen PF, de Boer NKH, de Meij TGJ. Endoscopic surveillance for colorectal cancer in pediatric ulcerative colitis: a survey among Dutch pediatric gastroenterologists. *JPGN Reports.* 2023; 4(3):e341. doi:10.1097/PG9.0000000000000341
 44. Gordon H, Biancone L, Fiorino G, et al. ECCO guidelines on inflammatory bowel disease and malignancies. *J Crohn's Colitis.* 2023;17(6):827-854. doi:10.1093/ecco-jcc/ijac187
 45. Keswani RN, Crockett SD, Calderwood AH. AGA clinical practice update on strategies to improve quality of screening and surveillance colonoscopy: expert review. *Gastroenterology.* 2021;161(2):701-711. doi:10.1053/j.gastro.2021.05.041
 46. Pouw RE, Bisschops R, Gecse KB, et al. Endoscopic tissue sampling - part 2: lower gastrointestinal tract. European Society of Gastrointestinal Endoscopy (ESGE) guideline. *Endoscopy.* 2021;53(12):1261-1273. doi:10.1055/a-1671-6336
 47. Boonstra K, Weersma RK, van Erpecum KJ, et al. Population-based epidemiology, malignancy risk, and outcome of primary sclerosing cholangitis. *Hepatology.* 2013;58(6):2045-2055. doi:10.1002/hep.26565
 48. Weismüller TJ, Trivedi PJ, Bergquist A, et al. Patient age, sex, and inflammatory bowel disease phenotype associate with course of primary sclerosing cholangitis. *Gastroenterology.* 2017;152(8):1975-1984.e8. doi:10.1053/j.gastro.2017.02.038
 49. Carbone M, Della Penna A, Mazzarelli C, et al. Liver transplantation for primary sclerosing cholangitis (PSC) with or without inflammatory bowel disease (IBD): a European Society of Organ Transplantation (ESOT) consensus statement. *Transpl Int.* 2023;36:11729. doi:10.3389/ti.2023.11729
 50. Nagai S, Safwan M, Kitajima T, Yeddula S, Abouljoud M, Moonka D. Disease-specific waitlist outcomes in liver transplantation - a retrospective study. *Transpl Int.* 2021;34(3):499-513. doi:10.1111/tri.13814
 51. Goet JC, Hansen BE, Tieleman M, et al. Current policy for allocation of donor livers in the Netherlands advantages primary sclerosing cholangitis patients on the liver transplantation waiting list-a retrospective study. *Transpl Int.* 2018;31(6):590-599. doi:10.1111/tri.13097
 52. Martinez M, Perito ER, Valentino P, et al. Recurrence of primary sclerosing cholangitis after liver transplant in children: an international observational study. *Hepatology.* 2021;74(4):2047-2057. doi:10.1002/hep.31911
 53. Visseren T, Erler NS, Heimbach JK, et al. Inflammatory conditions play a role in recurrence of PSC after liver transplantation: an international multicentre study. *JHEP Reports.* 2022; 4(12):100599. doi:10.1016/j.jhepr.2022.100599
 54. Buchholz BM, Lykoudis PM, Ravikumar R, Pollok JM, Fusai GK. Role of colectomy in preventing recurrent primary sclerosing cholangitis in liver transplant recipients. *World J Gastroenterol.* 2018;24(28):3171-3180. doi:10.3748/wjg.v24.i28.3171
 55. Deneau M, Perito E, Ricciuto A, et al. Ursodeoxycholic acid therapy in pediatric primary sclerosing cholangitis: predictors of gamma glutamyl-transferase normalization and favorable clinical

- course. *J Pediatr*. 2019;209:92-96.e1. doi:10.1016/j.jpeds.2019.01.039
56. Abbas N, Quraishi MN, Trivedi P. Emerging drugs for the treatment of primary sclerosing cholangitis. *Curr Opin Pharmacol*. 2022;62:23-35. doi:10.1016/j.coph.2021.11.003
 57. Lindor KD, Kowdley KV, Luketic VAC, et al. High-dose ursodeoxycholic acid for the treatment of primary sclerosing cholangitis. *Hepatology*. 2009;50(3):808-814. doi:10.1002/hep.23082
 58. Leoni MC, Amelung L, Lieveld FI, et al. Adherence to ursodeoxycholic acid therapy in patients with cholestatic and autoimmune liver disease. *Clin Res Hepatol Gastroenterol*. 2019;43(1):37-44. doi:10.1016/j.clinre.2018.08.006
 59. Black DD, Mack C, Kerker N, et al. A prospective trial of withdrawal and reinstitution of ursodeoxycholic acid in pediatric primary sclerosing cholangitis. *Hepatol Commun*. 2019;3(11):1482-1495. doi:10.1002/hep4.1421
 60. Laborda TJ, Jensen MK, Kavan M, Deneau M. Treatment of primary sclerosing cholangitis in children. *World J Hepatol*. 2019;11(1):19-36. doi:10.4254/wjh.v11.i1.19
 61. Trivedi PJ, Hirschfield GM, Adams DH, Vierling JM. Immunopathogenesis of primary biliary cholangitis, primary sclerosing cholangitis and autoimmune hepatitis: themes and concepts. *Gastroenterology*. 2024;166(6):995-1019. doi:10.1053/j.gastro.2024.01.049
 62. Tan LZ, Reilly CR, Steward-Harrison LC, Balouch F, Muir R, Lewindon PJ. Oral vancomycin induces clinical and mucosal remission of colitis in children with primary sclerosing cholangitis-ulcerative colitis. *Gut*. 2019;68(8):1533-1535. doi:10.1136/gutjnl-2018-316599
 63. Ricciuto A, Liu K, El-Matary W, et al. Oral vancomycin is associated with improved inflammatory bowel disease clinical outcomes in primary sclerosing cholangitis-associated inflammatory bowel disease (PSC-IBD): a matched analysis from the paediatric PSC Consortium. *Aliment Pharmacol Ther*. 2024;59:1236-1247. doi:10.1111/apt.17936
 64. Ali AH, Damman J, Shah SB, et al. Open-label prospective therapeutic clinical trials: oral vancomycin in children and adults with primary sclerosing cholangitis. *Scand J Gastroenterol*. 2020;55(8):941-950. doi:10.1080/00365521.2020.1787501
 65. Davies YK, Cox KM, Abdullah BA, Safta A, Terry AB, Cox KL. Long-term treatment of primary sclerosing cholangitis in children with oral vancomycin: an immunomodulating antibiotic. *J Pediatr Gastroenterol Nutr*. 2008;47(1):61-67. doi:10.1097/MPG.0b013e31816fee95
 66. Deneau MR, Mack C, Mogul D, et al. Oral vancomycin, ursodeoxycholic acid, or no therapy for pediatric primary sclerosing cholangitis: a matched analysis. *Hepatology*. 2021;73(3):1061-1073. doi:10.1002/hep.31560
 67. Sidoroff M, Kolho KL. Screening for adrenal suppression in children with inflammatory bowel disease discontinuing glucocorticoid therapy. *BMC Gastroenterol*. 2014;14:51. doi:10.1186/1471-230X-14-51
 68. Manwani K, Mieli-Vergani G, Mancell S, Dhawan A, Hadzic N, Samyn M. Long-term growth in children and young people with autoimmune liver disease treated with daily steroids. *J Pediatr Gastroenterol Nutr*. 2022;75(3):252-256. doi:10.1097/MPG.0000000000003513
 69. Snijders RJALM, Stoelinga AEC, Gevers TJG, et al. An open-label randomised-controlled trial of azathioprine vs. mycophenolate mofetil for the induction of remission in treatment-naïve autoimmune hepatitis. *J Hepatol*. 2024;80(4):576-585. doi:10.1016/j.jhep.2023.11.032
 70. Hedin CRH, Sado G, Ndegwa N, et al. Effects of tumor necrosis factor antagonists in patients with primary sclerosing cholangitis. *Clin Gastroenterol Hepatol*. 2020;18(10):2295-2304.e2. doi:10.1016/j.cgh.2020.02.014
 71. Hommes DW, Erkelens W, Ponsioen C, et al. A double-blind, placebo-controlled, randomized study of infliximab in primary sclerosing cholangitis. *J Clin Gastroenterol*. 2008;42(5):522-526. doi:10.1097/MCG.0b013e3181662426
 72. Laborda TJ, Ricciuto A, Aumar M, et al. Vedolizumab therapy in children with primary sclerosing cholangitis: data from the pediatric primary sclerosing cholangitis consortium. *J Pediatr Gastroenterol Nutr*. 2020;71(4):459-464. doi:10.1097/MPG.0000000000002855
 73. van Rheenen PF, Aloï M, Assa A, et al. The medical management of paediatric Crohn's disease: an ECCO-ESPGHAN guideline update. *J Crohn's Colitis*. 2020;15:171-194. doi:10.1093/ecco-jcc/jjaa161
 74. Turner D, Ruemmele FM, Orlanski-Meyer E, et al. Management of paediatric ulcerative colitis, part 1: ambulatory care: an evidence-based guideline from European Crohn's and Colitis Organization and European Society of Paediatric Gastroenterology, Hepatology and Nutrition. *J Pediatr Gastroenterol Nutr*. 2018;67(2):257-291. doi:10.1097/MPG.0000000000002035
 75. Ponsioen CY, Arnelo U, Bergquist A, et al. No superiority of stents vs balloon dilatation for dominant strictures in patients with primary sclerosing cholangitis. *Gastroenterology*. 2018;155(3):752-759.e5. doi:10.1053/j.gastro.2018.05.034
 76. Vanhanen E, Ovissi A, Kolho KL, Tenca A. Magnetic resonance cholangiography in the diagnosis of dominant strictures in pediatric-onset primary sclerosing cholangitis. *Dig Liver Dis*. 2023;55:1496-1501. doi:10.1016/j.dld.2023.05.024
 77. Nordenvall C, Olén O, Nilsson PJ, et al. Colectomy prior to diagnosis of primary sclerosing cholangitis is associated with improved prognosis in a nationwide cohort study of 2594 PSC-IBD patients. *Aliment Pharmacol Ther*. 2018;47(2):238-245. doi:10.1111/apt.14393
 78. Trivedi PJ, Bowlus CL, Yimam KK, Razavi H, Estes C. Epidemiology, natural history, and outcomes of primary sclerosing cholangitis: a systematic review of population-based studies. *Clin Gastroenterol Hepatol*. 2022;20(8):1687-1700.e4. doi:10.1016/j.cgh.2021.08.039

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Van Rheenen PF, Kolho K-L, Russell RK, et al. Primary sclerosing cholangitis in children with inflammatory bowel disease: an ESPGHAN position paper from the Hepatology Committee and the IBD Porto group. *J Pediatr Gastroenterol Nutr*. 2025;80:374-393. doi:10.1002/jpn3.12378