

Amsterdam UMC

The E-AHPBA – ESSO – Innsbruck consensus recommendations on peri- and postoperative management following liver resection

Maier, Eva; Stättner, Stefan; Carrion-Alvarez, Lucia; di Martino, Marcello; Olthof, Pim; Primavesi, Florian; Sochorova, Dana; van Laarhoven, Stijn; Balakrishnan, Anita; Breitskopf, Robert; Buis, Carlijn I.; Cipriani, Federica; Erdmann, Joris; Frampton, Adam; Fuks, David; Gilg, Stefan; Gulla, Aiste; Lancellotti, Francesco; Margreiter, Christian; Melloul, Emmanuel; Oberkofler, Christian; Petritsch, Stefan; Raab, Helmut; Rahbari, Nuh N.; Rappold, Daniela; Reiberger, Thomas; Ruzzenente, Andrea; Sallinen, Ville; Schäfer, Benedikt; Schnitzbauer, Andreas A.; Serrablo, Alejandro; Soreide, Kjetil; Sparrelid, Ernesto; Starlinger, Patrick; Stavrou, Gregor A.; Tinguely, Pascale; Aldrighetti, Luca; Dasari, Bobby V. M.; Donadon, Matteo; Dopazo, Cristina; Gruenberger, Thomas; Jonas, Eduard; Malik, Hassan; Viganó, Luca; Siriwardena, Ajith K.; Maglione, Manuel

Published in:
HPB

DOI:
[10.1016/j.hpb.2025.10.008](https://doi.org/10.1016/j.hpb.2025.10.008)

E-pub ahead of print: 01/01/2025

Document Version
Publisher's PDF, also known as Version of record

Document license
CC BY

[Link to publication](#)

Citation for published version (APA):

Maier, E., Stättner, S., Carrion-Alvarez, L., di Martino, M., Olthof, P., Primavesi, F., Sochorova, D., van Laarhoven, S., Balakrishnan, A., Breitskopf, R., Buis, C. I., Cipriani, F., Erdmann, J., Frampton, A., Fuks, D., Gilg, S., Gulla, A., Lancellotti, F., Margreiter, C., ... Maglione, M. (2025). The E-AHPBA – ESSO – Innsbruck consensus recommendations on peri- and postoperative management following liver resection. *HPB*. Advance online publication. <https://doi.org/10.1016/j.hpb.2025.10.008>

General rights

It is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), other than for strictly personal, individual use, unless the work is under an open content license (like Creative Commons).

Disclaimer/Complaints regulations

If you believe that digital publication of certain material infringes any of your rights or (privacy) interests, please let the Amsterdam UMC Medical Library know, stating your reasons. In case of a legitimate complaint, the Medical Library will make the material inaccessible and/or remove it from the website. Contact address: outputregistratie@amsterdamumc.nl

Download date: 24. Jul. 2026

CLINICAL PRACTICE GUIDELINE

The E-AHPBA – ESSO – Innsbruck consensus recommendations on peri- and postoperative management following liver resection

Eva Maier^{1,*}, Stefan Stättner^{2,3,*}, Lucia Carrion-Alvarez⁴, Marcello Di Martino^{5,6}, Pim Olthof⁷, Florian Primavesi¹, Dana Sochorova⁸, Stijn van Laarhoven⁹, Anita Balakrishnan¹⁰, Robert Breitkopf¹¹, Carlijn I. Buis¹², Federica Cipriani¹³, Joris Erdmann¹⁴, Adam Frampton^{15,16}, David Fuks^{17,18}, Stefan Gilg¹⁹, Aiste Gulla²⁰, Francesco Lancellotti^{21,22}, Christian Margreiter²³, Emmanuel Melloul^{17,18}, Christian Oberkofler²⁴, Stefan Petritsch²⁵, Helmut Raab¹¹, Nuh N. Rahbari²⁶, Daniela Rappold²⁵, Thomas Reiberger²⁷, Andrea Ruzzenente²⁸, Ville Sallinen^{29,30}, Benedikt Schäfer³¹, Andreas A. Schnitzbauer³², Alejandro Serrablo³³, Kjetil Soreide^{34,35}, Ernesto Sparrelid¹⁹, Patrick Starlinger^{36,37}, Gregor A. Stavrou³⁸, Pascale Tinguely³⁹, Luca Aldrighetti¹³, Bobby V.M. Dasari⁴⁰, Matteo Donadon^{5,6}, Cristina Dopazo⁴¹, Thomas Gruenberger⁴², Eduard Jonas⁴³, Hassan Malik⁴⁴, Luca Viganò^{45,46}, Ajith K. Siriwardena^{21,**} & Manuel Maglione^{1,*}

¹Department of Visceral, Transplant and Thoracic Surgery, Medical University of Innsbruck, Innsbruck, ²Department of Visceral and General Surgery, Hepatobiliary Center, Kepler University Hospital GmbH, ³Johannes Kepler University, Medical Faculty, Linz, Austria, ⁴HPB Unit, General Surgery Department, Fuenlabrada University Hospital, Madrid, Spain, ⁵Department of Health Sciences, Università del Piemonte Orientale, ⁶Division of General Surgery, Azienda Ospedaliera Universitaria Maggiore della Carità, Novara, Italy, ⁷Department of Surgery, Erasmus MC Cancer Institute, Rotterdam, the Netherlands, ⁸Department of Surgery, Tomas Bata Hospital Zlin, Czech Republic, ⁹Department of Surgery, Leiden University Medical Center, the Netherlands, ¹⁰Cambridge Hepatopancreatobiliary Unit, Cambridge University Hospitals NHS Foundation Trust, Cambridge, United Kingdom, ¹¹Department of Anaesthesiology and Intensive Care Medicine, Medical University Innsbruck, Innsbruck, 6020, Austria, ¹²Department of Hepato-Pancreato-Biliary Surgery and Liver Transplantation, University Medical Centre Groningen, Groningen, the Netherlands, ¹³Hepatobiliary Surgery Division, San Raffaele Scientific Institute, Milan, Italy, ¹⁴Department of Surgery, Amsterdam UMC, Amsterdam, the Netherlands, ¹⁵HPB Surgical Unit, Royal Surrey NHS Foundation Trust, ¹⁶Section of Oncology, Dept. of Clinical & Experimental Medicine, University of Surrey, Guildford, Surrey, United Kingdom, ¹⁷Department of Visceral Surgery, Lausanne University Hospital (CHUV), ¹⁸Faculty of Biology and Medicine, University of Lausanne (UNIL), Lausanne, Switzerland, ¹⁹Department of Clinical Science, Intervention and Technology, Division of Surgery and Oncology, Karolinska Institutet, Karolinska University Hospital, Stockholm, Sweden, ²⁰Faculty of Medicine, Institute of Clinical Medicine, Vilnius University, Vilnius, Lithuania, ²¹Regional Hepato-Pancreato-Biliary Unit, Manchester Royal Infirmary, Manchester, United Kingdom, ²²Robotic and Minimally Invasive digestive Surgery, Arcispedale Sant'Anna, Ferrara, Italy, ²³Department of General, Visceral and Transplant Surgery, Medical University of Graz, Graz, Austria, ²⁴Vivévis AG - Visceral, Tumor and Robotic Surgery, Clinic Hirslanden Zurich, Zurich, Switzerland, ²⁵Department for General-, Visceral- and Vascular Surgery, Salzkammergut Klinikum Vöcklabruck, Vöcklabruck, Austria, ²⁶Department of Surgery, University Medical Center Ulm, Ulm, Germany, ²⁷Division of Gastroenterology and Hepatology, Department of Medicine III and CD-Lab for Portal Hypertension and Liver Fibrosis, Medical University of Vienna, Vienna, Austria, ²⁸General and Hepatobiliary Surgery, Department of Surgery, Dentistry, Gynecology and Pediatrics University of Verona, GB Rossi Hospital, Verona, Italy, ²⁹Department of Gastroenterological Surgery, ³⁰Department of Transplantation and Liver Surgery, Helsinki University Hospital and University of Helsinki, Helsinki, Finland, ³¹Department of Gastroenterology, Medical University Innsbruck, Innsbruck, Austria, ³²Knappschaft Clinics University Hospital Bochum, Ruhr-University Bochum, Bochum, Germany, ³³HPB Unit of Miguel Servet University Hospital, Zaragoza, Spain, ³⁴Department of Gastrointestinal Surgery, HPB Unit, Stavanger University Hospital, Stavanger, ³⁵Department of Clinical Medicine, University of Bergen, Bergen, Norway, ³⁶Department of Surgery, Division of Hepatobiliary and Pancreatic Surgery, Mayo Clinic, Rochester, MN, USA, ³⁷Center of Physiology and Pharmacology, Medical University of Vienna, Vienna, Austria, ³⁸Department of General, Visceral and Thoracic Surgery, Surgical Oncology, Klinikum Saarbruecken, Saarbruecken, Germany, ³⁹Department of Visceral and Transplantation Surgery, University Hospital Zurich, University of Zurich, Zurich, Switzerland, ⁴⁰Department of HBP and Liver Transplantation Surgery, University of Birmingham, Birmingham, United Kingdom, ⁴¹Department of General Surgery, Division of HPB Surgery and Transplants, Vall d'Hebron Hospital Universitari, Vall d'Hebron Institut de Recerca (VHIR), Universitat Autònoma de Barcelona Vall d'Hebron Barcelona Hospital Campus, Barcelona, Spain, ⁴²Department of Surgery, Clinic Favoriten, HPB Center, Health Network Vienna and Sigmund Freud Private University, Vienna, Austria, ⁴³Surgical Gastroenterology Unit, Division of General Surgery, Department of Surgery, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa, ⁴⁴Liver Surgery Unit, Liverpool University Hospitals NHS Foundation Trust, Liverpool, United Kingdom, ⁴⁵Department of Biomedical Sciences, Humanitas University, Pieve Emanuele, Milan, and ⁴⁶Hepatobiliary Unit, Department of Minimally Invasive General & Oncologic Surgery, Humanitas Gavazzeni University Hospital, Bergamo, Italy

* These authors contributed equally to this work.

** Shared corresponding authorship.

Abstract

Background: Liver surgery carries a high risk of complications due to the interplay of patient-related factors, disease characteristics and liver function. This European consensus provides evidence-based guidance on selected aspects of perioperative care.

Methods: A modified Delphi process was used to achieve consensus, with a 70 % agreement threshold. The expert panel comprised hepatobiliary surgeons, anaesthetists, hepatologists, a nurse and physiotherapist. Systematic literature search was conducted in PubMed/Medline, EMBASE, Web of Science, and Cochrane databases. Evidence appraisal and statement development followed Scottish Intercollegiate Guidelines Network methodology.

Results: Six topics were addressed: (i) thromboprophylaxis, (ii) antibiotics, (iii) prehabilitation-nutrition-mobilisation, (iv) bile leaks, including bilio-enteric anastomosis, (v) post-hepatectomy haemorrhage, (vi) post-hepatectomy liver failure (PHLF). Screening yielded 204 included publications (initial 6514) and thirty-two statements were formulated (median evidence-level:2). Evidence strength varied by topic with lower evidence-levels in complex surgery settings and subcohorts. Study heterogeneity and specific inclusion criteria resulted in some topics in conditional recommendations despite high-level evidence. Weakest evidence was found for thromboprophylaxis and PHLF management. Strong recommendations were formulated for prehabilitation, mobilisation, and avoidance of routine drainage. Several evidence gaps warranting multicentre studies were identified.

Conclusion: Optimising perioperative care after hepatectomy remains challenging. Standardising key practices and addressing evidence gaps through collaborative research are vital to improve outcomes.

Received 22 September 2025; accepted 17 October 2025

Correspondence

Manuel Maglione, Department of Visceral, Transplant and Thoracic Surgery, Medical University of Innsbruck, Anichstrasse 35, 6020, Innsbruck, Austria. E-mail: manuel.maglione@i-med.ac.at (M. Maglione)

Correspondence

Ajith K. Siriwardena, Regional Hepato-Pancreato-Biliary Unit, Manchester Royal Infirmary, Oxford Road, Manchester, M13 9WL, United Kingdom. E-mail: Ajith.Siriwardena@mft.nhs.uk (A.K. Siriwardena)

Introduction

Liver resection is a fundamental intervention for the treatment of primary and secondary hepatic malignancies, as well as selected benign liver disorders. In spite of advances in surgical technique and perioperative care that have improved outcomes, liver resections are characterised by a complex interplay of patient-, malignancy- and liver function-related factors, resulting in a high risk of complications. These include bile leaks, haemorrhage, venous thromboembolism (VTE), surgical site infections (SSI), and post-hepatectomy liver failure (PHLF) eventually leading to worsened short- and long-term outcomes.^{1–5}

Despite the growing clinical experience, the evidence guiding peri- and postoperative management is frequently of moderate quality, largely derived from retrospective studies, institutional protocols, or expert opinion.

Professional societies such as the ERAS® (Enhanced Recovery after Surgery) Society have published guidelines with the aim of optimising perioperative patient management.⁶ However, with respect to prevention and management of

post-hepatectomy complications, to date, no consensus recommendations guiding patient care are available.

Considering the persistent inconsistency in clinical practices, the overarching objective of this consensus project is to furnish clinical guidance on specific perioperative matters and the management of postoperative complications, categorized in six topics.

The first addresses perioperative thromboprophylaxis, its timing and duration, and the management of occurring portal vein thrombosis. In the context of liver resections, there is often a delicate balance to be considered between the risk of bleeding and the occurrence of thrombotic complications.⁷ This is further complicated by the fact that patients frequently exhibit hypercoagulability during the perioperative period.⁸

In light of the safe, abbreviated perioperative regimens and reduced SSIs following tailored antibiotic administration,^{9,10} the second topic covers perioperative antibiotics, including recommendations on timing and length of antibiotic prophylaxis for standard liver resections and those for patients considered at high risk.

Given the expansion of boundaries in liver resection, the importance of preoperative multimodal prehabilitation protocols is increasing.^{6,11} Themes pertaining to pre-operative screening and perioperative nutritional optimisation were selected as third issue for the purpose of this consensus.

Postoperative bile leakage is a prevalent complication, with incidence rates ranging from 5 % to 15 %, and even higher following perihilar resections.^{12,13} The fourth topic discusses issues pertaining to leakage prevention, early detection, and treatment strategies.

The fifth subject pertains to post-hepatectomy haemorrhage (PHH). Rather than emphasising the management of this life-threatening complication, which is undoubtedly influenced by patients' conditions^{13,14}, this consensus focuses on various pre- and intraoperative strategies designed to avert this deleterious complication.

Finally, the sixth topic is PHLF. Following the recently published Innsbruck Consensus on preoperative liver function assessments¹⁵ this consensus focuses on diagnosis, management and monitoring of patients experiencing PHLF.

The objective of this European consensus project is to harmonise practices across these six domains. It should guide clinicians through evidence-informed recommendations for optimal peri- and postoperative care following liver resection.

Methods

Selection of committee members

The steering committee (EM, MM, AS, SS) selected 46 members for the expert panels, validation committee, and writing group based on their experience and expertise in the field of pre-, intra- and postoperative management of liver surgery patients and on their scientific contribution on challenges surrounding liver resections and complication management. Care was taken to ensure a balanced mix of participants' characteristics in terms of demographics (age, sex, country), clinical speciality and experience. The consensus development involved a broad variety of specialities (hepatobiliary surgeons, anaesthetists, hepatologists, a specialist nurse, and a physiotherapist) and was well balanced in terms of age and gender balance, and also nationalities with input from 21 different countries. All individuals made a declaration of interest regarding potential financial and non-financial conflicts related to the consensus recommendation content.

Selection of key topics to develop in the consensus

After an exploratory PUBMED literature review by the steering committee and following several online meetings (10/2023), six key areas were identified covering different topics on peri- and postoperative management following liver resection: (i) Perioperative thromboprophylaxis; (ii) Perioperative antibiotics; (iii) Prehabilitation, nutrition, mobilisation; (iv) Post-hepatectomy bile leak including bilio-enteric anastomosis leaks; (v) Post-hepatectomy haemorrhage (PHH); (vi) Post-

hepatectomy liver failure (PHLF). Each area was further stratified into sub-topics.

The experts were assigned to individual groups according to their scientific and clinical profile. Under the direction of elected group leaders, each group could either perform a systematic literature search by itself or was provided by the results of a centralized literature search querying MEDLINE®, EMBASE, Web of Science and Cochrane databases between 01/2010 and 02/2024 according to pre-specified Medical subject headings (MeSH) search terms, expanded by individual keywords related to sub-topics. Papers deemed to be pivotal to the subject matter, but not supplied by the centralized research due to their publication date were also included following a careful review by the expert panel. The publications were prefiltered according to the inclusion and exclusion criteria ([Supplementary Material: Supplementary Methods - Systematic Literature Review and Supplementary Table S1](#)).

Methodology for guideline development

The methodology followed a previously described process.¹⁶ Three validated methods were integrated: the SIGN (Scottish Intercollegiate Guidelines Network) methodology for the assessment of evidence and development of guideline statements¹⁷; the modified Delphi method for achieving expert consensus^{18,19}; and the AGREE II-GRS (Global Rating Scale) instrument for assessment of methodological quality and externally validating final statements.²⁰

Inclusion and grading of evidence

Included studies were assessed and graded according to the SIGN methodology by evaluating the "study quality" and "evidence level" according to the SIGN grading system ([Supplementary Tables S2 and S3](#)). The resulting "evidence tables" were then reviewed by the steering committee to ensure correct application of the inclusion and exclusion criteria and gradings. The working groups then created "considered judgement forms" to summarize the evidence, quality ratings, limitations, and strength of individual studies, resulting statements, strength of recommendations, and future areas of scientific interest.

Modified Delphi process

The key questions and proposed guideline statements with attached judgement forms were sent out for a stepwise Delphi process for anonymous voting to the whole expert panel. It allowed every member to agree or disagree with the statement, and to make comments and recommendations for changes. An agreement level of 70 % per Delphi stage was considered appropriate to ensure a balance between consensus and voting progress. In a total of three Delphi rounds, statements reaching 70 % agreement were excluded from further discussion, while statements below this level were reviewed by the steering committee and working group leaders and revised accordingly to enter the next Delphi round. At the final virtual preconference meeting the remaining statements were again discussed, reviewed, and then adopted for the onsite

conference. This conference was held as part of the Surger-Innsbruck International Meeting on Liver Surgery on December 12–13, 2024 in Innsbruck, Austria. Here, all key questions, statements and preliminary paragraphs drafted by the six groups were summarized with their underlying evidence and presented to the committee and the audience. All congress participants could discuss and ultimately vote electronically on their individual support for the statements. The presentations, discussions and voting results were reviewed by the validation committee, which provided final thoughts and comments on each statement before endorsing it. This final voting of the conference participants and the validation committee was considered by the writing and steering committee for refining the wording of individual statements and expert comment paragraphs for the published manuscript.

Validation

The entire guideline process was quality-controlled by the validation committee (Supplementary Figures). All members of the panel and a patient representative (Gerhard Lobenscheg, patient support group “L(i)eberleben”, Tyrol, Austria) reviewed the draft before submitting it for publication. The editorial staff of the publisher was involved during the process in terms of formal structure and layout. The consensus development was endorsed by the Scientific and Research Committee of the European-African Hepato-Pancreato-Biliary Association (E-AHPBA) and the Board of Directors of the European Society of Surgical Oncology (ESSO).

Overall results of Delphi process

A total of 6514 publications were reviewed by the six working groups, resulting in a final number of 204 included manuscripts (Supplementary Results – Literature Review).

After two online Delphi rounds all but one statement had achieved 70 % agreement level. This one was reviewed, discussed, and then finally included after reaching sufficient agreement during the onsite conference. All statements ($n = 32$) were supported by the validation committee and included in the paper after final revision. The overall methodological quality of the guideline development was considered 6.6 to 7.0 by all members of the validation committee (Supplementary Figures).

Each clinical question is answered with a short overall statement, including the grade of recommendation and the level of evidence, followed by an expert panel comment including a selection of relevant publications related to the question. A complete list of references for each topic can be found in Supplementary Results – Literature Review. An outlook for future research questions has been generated.

Topic 1: perioperative thromboprophylaxis

Q1 When should perioperative thromboprophylaxis be commenced in liver surgery?

Pharmacological thromboprophylaxis is recommended for all patients undergoing liver surgery and can be initiated either pre- or postoperatively. Mechanical thromboprophylaxis (socks, pneumatic) should be considered routinely.

Recommendation grade: conditional | median level of evidence: 2+ (range: 2++ to 3)

Expert panel comment

The incidence of post-hepatectomy venous thromboembolism (VTE) has been reported to range between 2 % and 6 %, which is higher than the incidence observed for many other abdominal surgeries and is directly proportional to the magnitude of the hepatectomy, with extended hepatectomies exhibiting the highest risk.²¹ Published hepatectomy bleeding event rates range from 1 % to 8 %, ¹³ with most series focusing their data on intra-operative bleeding and transfusion rates. An ACS-NSQIP analysis reported that the gradient between VTE event rate and bleeding complications ranged from 3.5 to 8.3 times and increased significantly with the magnitude of hepatectomy.²¹

Pharmacological thromboprophylaxis with low-molecular weight heparin (LMWH) or unfractionated heparin should be initiated as a standard postoperative care measure, unless deemed contraindicated by exceptional circumstances.⁷ There is no high-quality evidence to guide the optimal timing of thromboprophylaxis in liver surgery specifically; in particular, evidence is lacking on whether it should be initiated preoperatively or postoperatively.²²

Retrospective uncontrolled studies have provided low-quality evidence suggesting that preoperatively initiated pharmacological thromboprophylaxis using LMWH could decrease VTE events.^{21,23} However, regarding bleeding, the reports are conflicting. Therefore, the current clinical practice is highly variable between centers.²⁴ While postoperatively initiated thromboprophylaxis is associated with lower VTE rates without an increase in bleeding complications, no strong recommendation can be made in the preoperative setting. A randomized controlled trial (RCT) (PREPOSTEROUS trial, NCT04731558) comparing preoperative to postoperative thromboprophylaxis in liver surgery is currently recruiting.^{25,26} Until high-quality evidence emerges, either approach is acceptable.

Q2 What is the optimal duration of perioperative thromboprophylaxis in liver surgery?

A 4-week period of administration of LMWH should be considered after major liver surgery. However, prolonged (>4 weeks) prophylaxis may be indicated in high-risk patients based on individual risk stratification.

Recommendation grade: conditional | median level of evidence: 2- (range: 2- to 5)

Expert panel comment

High-level evidence for perioperative thromboprophylaxis specifically in liver surgery, adjusted to patient's factors, the extent of surgery and open versus minimally invasive approach is lacking. The term "prolonged" prophylaxis is used variably in the literature, referring to 4 weeks or more than 4 weeks, depending on the study. Additionally, most of the data are derived from the in-hospital stay with limited information on post-discharge VTE and efficacy of its prevention. Since there are studies indicating an increased risk of venous thromboembolism after liver surgery compared to other oncological surgery, many advocate prolonged prophylaxis of four weeks to all patients undergoing liver surgery for malignancy and this seems to be feasible without increased risk of bleeding complications. Others only recommend prolonged duration to high-risk patients using individual risk stratification tools²⁷ (e.g. Padua prediction score²⁸) to define high-risk patients. Antithrombin III (AT III) deficiency and PHLF must be considered as confounding factors. Liver resections of a limited extent done by minimally invasive approach carry less risks and these patients could be considered for less than 4 weeks of LMWH, unless individual other risk factors exist related to chronic conditions.

Q3 What anticoagulation should be used when performing hepatectomy with portal vein resection and reconstruction?

There is no strong evidence for specific perioperative prevention strategy of portal vein thrombosis in patients undergoing liver surgery with portal vein resection and reconstruction. The approach should be tailored according to the patient-specific factors and clinical judgement.

Recommendation: conditional | median level of evidence: 2-range: 1+- to 2-)

Expert panel comment

Miyazaki in 2016 analysed 270 patients who underwent a portal vein resection (including end-to-end anastomosis, patch or segmental graft and venoraphy) after 74 hepatectomies and 196 pancreatectomies.²⁹ They introduced a systemic heparinization (200–240 u/kg/day) from the first postoperative day to 7 postoperative days. They experienced a postoperative portal vein thrombosis in 24 cases (8.9 %), 16 out of 74 patients (21.6 %) after hepatectomy and 8 out of 188 patients (4.3 %) after pancreatectomy. Molina *et al.* reported a policy of a prophylactic LMWH to all patients who underwent a hepatectomy for peri-hilar cholangiocarcinoma, moving to administration every 12 h for patients with vascular resection.³⁰ Considering lack of evidence on this topic, a tailored approach may be applied according to patient-specific factors. Ejaz *et al.* found that patients with previous history of VTE, prolonged operative time (OR 1.17 per additional hour, 95 % CI 1.04–1.32, $P = 0.009$) and increased length of stay were independent risk factors for VTE.³¹ Unfortunately, no direct mention has been

made regarding portal vein thrombosis in patients who received a vascular resection and reconstruction during hepatectomy.

Q4 Is there a role for defining early and late portal vein thrombosis (PVT) after liver resection?

Defining early and late PVT after liver resection should be considered with the aim to standardize outcomes. A widely accepted definition specific to liver resection is lacking. A 30-day cut-off, as used in liver transplantation, is a pragmatic approach that may be adopted, with limitations related to partially different pathophysiological causes of PVT after liver resection compared to liver transplantation.

Recommendation: conditional.

Expert panel comment

Early PVT is typically related to immediate postoperative factors, such as direct surgical manipulation or technical failure, intraoperative hemodynamic changes, or pro-thrombotic conditions. The detection of very early postoperative PVT would necessitate a routine imaging protocol, such as Doppler ultrasonography within 48 h of major liver surgery. Late PVT, occurring after 30 days, often reflects long-term processes, including liver regeneration and altered portal venous flow. Proposed post-resection PVT classification system³² include categories according to the site of the thrombus but do not reflect the time aspect. The decision for a 30-day cut-off is a practical solution to distinguish these phases but remains somewhat arbitrary and may require further validation. Defining PVT according to the time of onset may be useful for academic purposes. From a clinical point of view, the cut-off should also consider the management strategy for PVT, such as anticoagulation or invasive interventions, as these approaches may differ based on the timing, location and nature of the thrombosis. Early PVT may more often require reoperation, late PVT is often asymptomatic, detected on follow-up imaging and can be mainly managed by anticoagulation.

Q5 Are there differences in the treatment of early and late portal vein thrombosis after liver resection?

There are no high-quality comparative studies to establish standardized treatment algorithms for early and late PVT after liver resection. Treatment options include anticoagulation, surgical intervention, and interventional radiology procedures. The decision on treatment strategy should be adjusted according to the time onset, thrombus extent and site, patient-specific factors, and institutional resources.

Recommendation: conditional.

Expert panel comment

Early PVT poses a significant risk of complications such as liver failure, portal hypertension, or bowel ischemia and requires

timely intervention. Treatment strategies depend not only on the extent (partial versus total), timing (early versus late) and duration of existence of the thrombosis, but also on the site and on the patient's conditions at the time of diagnosis. Early thrombosis at the site of reconstruction can be considered for surgical approach, as one of the possible causes is a technical factor following the index procedure. The administration of anticoagulation aims to prevent thrombosis extension and related decrease in portal venous flow, which may result in portal hypertension and liver failure. Late PVT management focuses on addressing long-term complications, such as chronic portal hypertension, with tailored anticoagulation strategies based on follow-up imaging. While institutional resources play a crucial role in determining the feasibility of specific interventions, such as advanced interventional radiology or surgical expertise, there remains a significant gap in high-level evidence to support specific recommendations.

Topic 2: perioperative antibiotics

Q1 Which antibiotics should be used as standard prophylaxis for liver surgery?

For liver surgery prophylaxis, antibiotic choice should align with local pathogen resistance patterns. First-generation cephalosporins such as cefazolin are preferred for livers with naive bile ducts, administered within 60 min before incision. Broader-spectrum agents, like ceftriaxone or Piperacillin-Tazobactam, are conditionally recommended for high-risk cases. Consultation with a microbiologist should be considered in MRSA-risk settings. In beta-lactamic allergic patients Fosfomycin or aminoglycosides should be selected.

Recommendation: conditional | median level of evidence: 2++ (range: 1++ to 2+)

Expert panel comment

Guidelines for prophylactic antibiotics in liver surgery emphasize the need to balance infection prevention with antibiotic stewardship. Currently, there is no conclusive evidence to either support or refute the routine use of antibiotic prophylaxis for all liver resections.^{33,34} However, cefazolin, a first-generation cephalosporin, is recommended for standard liver resections due to its effectiveness in reducing surgical site infections (SSIs) and its minimal impact on 30-day readmission rates. Administered 1 h before surgery, cefazolin provides reliable coverage for clean-contaminated procedures, addressing the infection risk in most patients undergoing liver resection.^{35,36}

For higher-risk patients, broader-spectrum antibiotics like third generation cephalosporins or beta-lactam/beta-lactamase inhibitors may be appropriate, providing enhanced coverage against gram-negative and anaerobic pathogens. The decision to use these alternatives should consider patient risk factors and

local resistance trends. Although effective, these agents are recommended conditionally, given the moderate-quality evidence supporting their use in specific cases.^{37–43}

For patients at risk of MRSA infection, adding vancomycin to the prophylactic regimen is suggested, but should be guided by local resistance data and individual patient factors as well as consultation with an infectious disease specialist to prevent resistance development. Aminoglycosides are reserved due to nephrotoxicity concerns and are recommended only for cases with specific coverage needs. This framework reflects a careful, evidence-based approach prioritizing effective prophylaxis while mitigating antibiotic resistance risks.^{44,45}

Q2 Which patients should be defined as high-risk for developing clinically relevant SSI following hepatectomy?

Patients at higher risk of SSI can be identified by bacterial colonization resistant to commonly used prophylactic antibiotics, as well as by factors which make them generally more at risk for postoperative complications like underlying liver disease, preoperative biliary drainage, and synchronous resections of the gastrointestinal tract. In patients with preoperative biliary drainage cultures should be obtained at the time of the intervention. In case of bile duct resection and reconstruction intraoperative biliary cultures should be obtained during the surgical procedure.

Recommendation: conditional | median level of evidence: 2+ (range: 1++ to 2+)

Expert panel comment

Several risk factors are associated with a higher risk of developing SSI following hepatectomy. Most of the evidence comes from multivariable analyses of retrospective observational studies, with very few RCTs addressing this issue.⁴³ Patients at high risk for developing clinically relevant SSIs after hepatectomy can be categorized based on the preoperative, surgical, and postoperative risk factors.

Preoperative risk factors (moderate to high level of evidence) include patient related factors (obesity, diabetes mellitus, chronic pulmonary diseases, and chronic cardiac diseases,^{46,47} preoperative anaemia, malnutrition, or hypoalbuminemia,⁴⁸ colonization with bacteria resistant to commonly used perioperative antibiotics^{37,46,49}), liver-related factors (hepatolithiasis, chronic liver disease, biliary obstruction, or preoperative biliary drainage^{50–53}) and technical or logistic factors (previous liver resection, planned bilio-enteric anastomosis,^{37,54,55} prior hospitalization^{49,52}).

Surgical risk factors (moderate level of evidence) comprehend high volumes of intravenous fluids, blood transfusion, prolonged operation time, and concomitant bowel surgery.^{37,48,51,52,55} Finally, persistent postoperative elevation of inflammatory markers such as C-reactive protein (CRP) or bile

leakage^{56,57} are examples of postoperative risk factors (moderate level of evidence).

Despite these risk factors, clear criteria for identifying such high-risk patients are not yet well defined.

Q3 Which patients are likely to benefit from extended antibiotic prophylaxis?

Although antibiotic prophylaxis is recommended, extended postoperative antibiotic prophylaxis for more than 24 h has not demonstrated any additional benefit in preventing postoperative infections after standard hepatectomies. Patients at high risk for developing clinically relevant SSI may benefit from extended antimicrobial duration (three to five days postoperatively).

Recommendation: conditional | median level of evidence: 2+ (range: 1+ to 2+)

Expert panel comment

RCT/high evidence data is mainly available for standard hepatectomies. There is lack of evidence supporting routine extended (three to five days postoperatively) antibiotic prophylaxis after standard liver resection, aligning with principles of antibiotic stewardship.^{34,58–61} Patients at high risk for developing clinically relevant SSI may benefit from extended antimicrobial duration. The emphasis on tailoring prolonged prophylaxis to high-risk patients, particularly those with preoperative biliary drainage, or complex procedures, is consistent with findings from recent studies.^{40,43,62,63} However, the recommendation would benefit from a clearer stratification of risk factors and more specific guidance on how intraoperative cultures should influence postoperative antimicrobial strategies. Additionally, defining thresholds (e.g., bile contamination, duration of surgery, or intraoperative findings) for initiating extended prophylaxis would enhance clinical applicability. Future prospective studies are needed to refine high-risk criteria and validate targeted protocols.⁶⁴

Q4 What is the role of antibiotic treatment in cases of postoperative fever of unknown origin and no suspicion of infection or sepsis following liver resection?

Antibiotic therapy is not routinely recommended for postoperative fever of unknown origin if diagnostic workup excluded any signs of infection or sepsis. High-risk patients—such as those with preoperative biliary drainages, bilioenteric anastomosis, immunosuppression, or comorbidities like diabetes mellitus—may benefit from targeted antibiotic therapy if early signs of infection are present.

Recommendation: strong | median level of evidence: 1+ (range: 2+ to 1++)

Expert panel comment

Postoperative fever after hepatectomy can arise from both infectious and non-infectious causes. Routine use of antibiotics for fever of unknown origin is generally discouraged due to the risk of promoting multidrug-resistant bacteria and complicating future antibiotic selection for septic events.^{43,45,58} If indicated, empirical antibiotic treatment should be guided by the severity of infection, patient-specific risk factors, and local antibiotic sensitivity patterns, covering common pathogens like *Enterococcus* species and gram-negative bacteria until specific pathogens are identified.^{36,43,60} Assessing preoperative bacterial colonization in patients with biliary abnormalities is crucial and is supported by high-level evidence, aiding in implementing targeted antibiotic strategies when necessary.^{33,36,41} This tailored approach aims to improve perioperative outcomes in selected high-risk patients while minimizing the development of antibiotic resistance.

Topic 3: prehabilitation, nutrition, mobilisation

Q1 What is the value of a prehabilitation program in patients undergoing hepatectomy?

Multimodal prehabilitation programs (exercise training, nutritional optimization, psychological interventions) should be considered for patients prior to hepatectomy for both benign and malignant liver disease following assessment to reduce postoperative morbidity and length of hospital stay.

Recommendation: strong | median level of evidence: 1+ (1- to 2++)

Expert panel comment

Prehabilitation programs (including physical exercise training, diet optimization, psychological interventions, smoking cessation, alcohol cessation) may have positive impact on outcomes after hepatectomy, especially for malignant liver disease (30-day morbidity reduced: trend,^{65,66} no effect^{67–69}; length of hospital stay shortened: significant,⁷⁰ trend,^{65,66} no effect^{68,69,71}; length of intensive care stay shortened⁷¹; quality of life improved⁷²). No effect of prehabilitation on postoperative mortality was shown in 2 systematic reviews (SR)/metaanalysis (MA).^{68,69} The content of prehabilitation programs evaluated for liver or HPB surgery is heterogenous, with 1 RCT evaluating exercise training after cardiopulmonary exercise testing (CPET) as unimodal intervention⁷³ and 7 SR/MA/RCT evaluating multimodal programs including in-hospital or home-based physical exercise training, dietary interventions or psychological support.^{66,67,70,74,75}

There is inconsistent literature on the duration of a prehabilitation program, with recently published guidelines

suggesting 3–6 weeks before major oncologic surgery.⁷⁶ For exercise training prior to liver surgery, a minimum of 4 weeks preoperatively is evaluated in 5 SR/MA/RCT.^{65,68,72,74,75} Alcohol and smoking cessation as part of prehabilitation is recommended for surgery in general,⁷⁶ with at least 4 weeks of cessation prior to hepatectomy.⁶ Due to the heterogeneity of the outcomes assessed in the included SR/MA/RCT, no clear recommendation can be made, indicating the need for further standardised research to assess prehabilitation in liver surgery.

Q2 What is the influence of preoperative body composition for outcomes in patients undergoing hepatectomy?

Preoperative body composition influences outcomes in patients undergoing hepatectomy for various conditions of liver malignancies. Therefore, preoperative screening for low skeletal muscle mass (LSMM) should be done in every patient considered for hepatectomy for liver malignancies.

Recommendation strong/median level of evidence: 1+ (1–2+)

Expert panel comment

At least 7 scientific papers (Meta analysis and Systematic Review) have pointed out “sarcopenia” or LSMM as major risk factors for OS (overall survival) and DFS (disease free survival) in patients undergoing hepatectomy for HCC^{77–83} and for postoperative complication rates in liver resection for CRLM (colorectal-liver metastasis).^{82,84} Only one high quality paper MA investigated the role of obesity on OS and DFS in patients undergoing hepatic resection for CRLM – no significant impact was found.⁸⁴

Preoperative screening for LSMM should be done in every patient undergoing hepatectomy for liver malignancies, as it shows significant impact on postoperative outcomes. Therefore CT-scans seem to be a good option, as they are routinely used in oncologic patients for staging examination and broadly accessible. As sarcopenia is defined as loss of muscle mass, reduced function and impaired muscle strength, the term “sarcopenia” should only be used if diagnosed according to the guidelines.⁸⁵ If diagnosed preoperatively, addressing LSMM and sarcopenia with nutritional support (e.g. protein supplementation) and physical activity should be considered.⁸⁶ There is only limited high quality evidence for explaining the influence of obesity on outcome in patients undergoing hepatic resection, hence further research is needed.

Q3 What is the role of fitness screening and assessment in patients undergoing hepatectomy?

As aerobic fitness is an important modifiable risk factor and can be improved by a prehabilitation program, screening for this risk factor can be incorporated as a routine part of the preoperative assessment for patients undergoing liver surgery,

particularly for those with borderline or reduced fitness, as it provides critical information for optimizing perioperative care and improving postoperative outcomes.

Recommendation: conditional | median level of evidence: 1 (1- to 2+)

Expert panel comment

Fitness screening and assessment allow for risk stratification and the identification of potential modifiable risk factors.⁸⁷ Aerobic fitness has a strong relationship with postoperative outcomes.^{88–91} Ideally it is measured by Cardiopulmonary Exercise Testing (CPET), alternatively by Steep Ramp Test (STRT), Six-Minute Walk Test (6MWT), or Thirty-Second Sit-to-Stand Test (30ST).^{92–97}

Lower CPET-derived thresholds, specifically AT values below 9.0–11.5 ml/kg/min and VO₂ peak below 15 ml/kg/min, are associated with increased risks of postoperative complications, longer hospital stays, and poorer overall survival.^{73,91,97,98} These markers serve as reliable predictors for identifying high-risk patients and are used to guide clinical decisions regarding the need for prehabilitation interventions and perioperative management.

CPET offers superior predictive accuracy compared to simpler tests such as the Incremental Shuttle Walk Test (ISWT) or the STRT, which have shown limited precision in predicting postoperative outcome.^{97,98} Despite their practicality, these alternative tests are not as sensitive or specific as CPET in evaluating fitness for liver surgery.

Q4 Does perioperative immunonutrition offer a benefit in patients undergoing hepatectomy?

Immunonutrition can be administered as perioperative nutritional supplementation in patients undergoing hepatectomy as it has a positive effect on postoperative liver function and might foster the regenerative potential especially in vulnerable populations.

Recommendation: conditional | median level of evidence: 1- (1- to 2+)

Expert panel comment

A review of 9 RCTs with 966 patients, including administration of omega-3-fatty acids (FA), Arginine/Glutamin, and RNA applications, found improvement in complication rates, overall postoperative infectious complications, and wound infections and a beneficial effect on the length of hospital stay.⁹⁹

No effects of immunonutrition could be found on the recurrence in HCC patients,¹⁰⁰ however, in an RCT of 320 patients undergoing liver resection for HCC, the intervention group profited significantly from postoperative parenteral administration of omega-3-FA with regard to complications and infections.¹⁰¹

The meta-analysis by Wong *et al.* confirmed the findings in 11 RCTs (>1.000 patients) with an improved clinical course reflected by lower complications. The timing of intervention was heterogeneous (pre/post), also the duration which lasted from 30 days preoperatively until 3 weeks postoperatively. Regimens were also heterogeneous and consisted of omega-3-FAs alone $\pm$ ribonucleic acids $\pm$ glutamine/arginine supplementation either enterally or fully parenterally.¹⁰²

Q5 What is the role of prokinetic policies to improve time to first bowel movement/full standard diet?

Routine use of postoperative prokinetics or laxatives is recommended to stimulate bowel movement after hepatectomy, while prokinetics may shorten the time to first flatus or stool. In patients who have had a simultaneous liver and colonic resection, use of prokinetic treatment should be used with caution and based upon established colorectal protocols.

Recommendation: conditional |median level of evidence: 1+ (1–2+)

Expert panel comment

At least 4 RCTs and 1 case control prospective study have shown that the use of prokinetics such as postoperative laxatives, gum chewing, herbal medicine, or decoction after hepatectomy reduce the time to first flatus or stool.^{103–107} However, prokinetics did not reduce secondary outcomes, such as delayed gastric emptying, length of stay, or time to functional recovery.

While some members of the expert panel suggested that time to first flatus or stool is not a significant end point in liver surgery, others argued that this might have a positive impact on patients' well-being and quality of life (QoL).

Q6 When and how should artificial nutrition be added to oral nutrition in the postoperative period?

Early oral nutrition should be preferred after hepatectomy. In patients requiring artificial nutrition support (malnourished patients, patients with complications causing several days of fasting), enteral nutrition through a nasofeeding tube should be preferred over total parenteral nutrition (TPN). Nutritional specialist assessment is recommended for patients with delayed oral intake.

Recommendation: conditional |median level of evidence: 1- (1- to 2++)

Expert panel comment

Early oral diet has been shown to be safe and to decrease the time to first bowel movement in abdominal and liver surgery.¹⁰⁸ The ESPEN guidelines on clinical nutrition in surgery recommend oral nutrition postoperatively.¹⁰⁹

A recent RCT of patients undergoing major abdominal surgery reported beneficial effects on nosocomial infections from

early parenteral nutritional support in addition to oral intake, compared with delayed parenteral nutrition.¹¹⁰ Two meta-analyses suggest that for patients at risk of impaired oral intake over a prolonged period, enteral nutrition (EN) through a naso-feeding tube should be preferred over TPN whenever possible.^{111,112}

Q7 How should (early) postoperative mobilization after hepatectomy be defined?

The ideal prescription on postoperative mobilization after hepatic surgery is not clear. Early mobilization after liver surgery should start immediately on postoperative day one with assisted active mobilization and sitting out of bed. Walking (with no precise target) should start at least on second postoperative day adapted to the patient's functional capacity and carefully supervised especially in elderly or frail patients.

Recommendation: strong |median level of evidence: 1+ (1–2)

Expert panel comment

Either resistance or endurance training should be part of the mobilization for this patient population. No literature was found on the realistic endpoints in early mobilization of elderly or frail patients. In 2 RCTs and 2 prospective cohort studies postoperative mobilization has shown an improvement on functional capacity.^{113–116} Only one study showed a reduction of the hospital length and consequently a cost-saving effect.¹¹³ No meta-analysis on this topic was found. 2 prospective cohort studies recommend activity tracking which might detect post-surgical complication in vulnerable patients and enhance the recovery.^{117,118}

Even though no study was found that provides a precise answer on this question, we agree that patients with a preoperative reduced level of physical activity require additional attention in the postoperative period. Especially for elderly or frail patients undergoing major surgery monitoring postoperative activity might be useful in identifying postsurgical complications. One RCT with 108 patients showed that early mobilization reduces the risk for inability to walk without human assistance at postoperative day 5 or hospital discharge.¹¹⁴ However, no differences were found regarding clinical outcomes or complications.

Topic 4: post-hepatectomy bile leak including bilio-enteric anastomosis leaks

Q1 Is there a role for intraoperative bile leak test (i.e. white test, methylene blue, indocyanine green (ICG) ...) after hepatectomy without bilio-enteric anastomosis?

Intraoperative bile leak tests can be considered after liver resection, especially in major resections or complex procedures,

recording its use in the operation report. No specific test has shown clear superiority regarding identification of leaks, although the gauze (surface) test and the “white-test” might be preferred. Therefore, the choice of test should depend on centre expertise and availability.

Recommendation: conditional (major/complex resections: strong) | median level of evidence: 1- (range: 1++ to 4)

Expert panel comment

Intraoperative bile leak testing plays an important role in identifying bile leaks at the resection plane during hepatectomy, especially after complex resections.^{119,120} Bile leaks, if undetected, can lead to potentially serious complications such as bile peritonitis and intra-abdominal abscesses. There are several tests available, including the injection of different substances into the bile ducts, such as air, saline, methylene blue, lipid-based solutions (“white test”), indocyanine green (ICG), standard cholangiography or a simple stain test using a gauze.

The “white test” is a cost-effective technique that has been shown in several studies,^{120,121} including two RCTs, to reduce postoperative bile leak rates by enabling intraoperative detection and immediate intervention.^{122,123} It is important to be aware this test may not detect disconnected segments. There is an ongoing RCT comparing the white test to the gauze test, with results expected in 2025.¹²⁴

ICG fluorescence imaging offers near-infrared visualization of bile ducts. While some retrospective observational studies and smaller trials suggest potential advantages of ICG in complex cases due to improved anatomical delineation,^{120,125} available RCT data indicate that ICG has not shown clear superiority over simpler and more accessible techniques techniques regarding bile leak detection rates and clinical outcomes. Despite one ongoing prospective trial,¹²⁶ further large-scale RCTs are necessary to conclusively determine the role of ICG for bile leak prevention.

Q2 Is there a role for drainage after hepatectomy without bilio-enteric reconstruction?

Routine abdominal drainage after an uncomplicated hepatectomy should be avoided.

Recommendation: strong | median level of evidence: 1+ (range: 1++ to 2+)

Expert panel comment

The role of routine abdominal drainage after hepatectomy without bilio-enteric reconstruction has been extensively debated, with current evidence strongly suggesting that it should be avoided in uncomplicated cases. A meta-analysis by Hajibandeh *et al.*,¹²⁷ which included seven RCTs involving 1064 patients, demonstrated that postoperative drainage not only fails to reduce the incidence of intra-abdominal collections requiring intervention but also increases the risk of overall and wound-related

complications. This finding is supported by earlier publications, such as the Cochrane review by Gurusamy *et al.*,¹²⁸ which concluded that routine drainage does not provide significant benefits in uncomplicated liver resections. Similarly, Sun *et al.*¹²⁹ conducted a RCT and found no advantage to routine drainage after elective hepatectomy using the crush clamp method. Furthermore, Liu *et al.*¹³⁰ highlighted that abdominal drainage is particularly contraindicated in patients with chronic liver diseases, as it may exacerbate complications without improving outcomes. Despite these findings, there are specific high-risk scenarios where drain placement might still be warranted. For instance, Dezfouli *et al.*¹³¹ and Brooke-Smith *et al.*¹³² suggest that drainage could be considered in cases involving difficult anatomical locations, such as the right superior segments, or when intraoperative bile leaks are detected and there are concerns about adequate control. Additionally, in settings where interventional radiology is not readily available, drainage may serve as a precautionary measure. However, the consensus, reinforced by systematic reviews such as those by Gavriilidis *et al.*¹³³ and Anweier *et al.*,¹³⁴ is that routine drainage after uncomplicated hepatectomy is unnecessary and may even be harmful.

Q3 What is the value of biliary trans-anastomotic stents in complex bilio-enteric anastomoses in hepatectomy?

Trans-anastomotic stents can be considered in complex bilio-enteric anastomoses (e.g., extended resections, small bile ducts, friable tissue) but are not recommended for routine use. If used, they should be removed within 3 months to minimize complications. The choice of trans-anastomotic stent depends on centre availability and expertise.

Recommendation: conditional | median level of evidence: 2- (range: 2++ to 4)

Expert panel comment

Bilio-enteric anastomotic bile leakage remains a challenging complication and it is associated with prolonged hospitalization and mortality. The decision to use trans-anastomotic stents in small ducts or high-risk anastomoses is still common practice aimed to reduce the risk of complications. However, it remains a topic of debate and should be based on the specific risks associated with the patient and the surgical procedure. Recent studies have identified several factors associated with bilio-enteric anastomotic bile leakage. The risk factors can be related to the surgical procedure (complex surgeries/compromised tissue) and to patient’s factors (comorbidities/previous surgery or radiation treatment).^{135,136}

Removal of these stents within 3 months is recommended to minimize side effects such as cholangitis and strictures.¹³⁷ Whether their routine use is indicated remains unclear, even if most (low evidence) studies have shown no benefit. Currently, there is no evidence supporting the use self-resorbable trans-anastomotic stents.

A large multicenter study found that preoperative biliary drainage was independently associated with a lower risk of anastomotic bile leakage, while postoperative external biliary drainage showed an association with higher leakage rates on univariate analysis but was not an independent risk factor in multivariable analysis.¹³⁵ Separately, the use of postoperative external biliary drains has also been identified as an independent risk factor for clinically relevant PHLF in patients undergoing major liver resection for perihilar cholangiocarcinoma.¹³⁸

The choice of anastomotic technique has not shown to be a relevant factor for its incidence in this study.¹³⁵ Other studies underscore the value of specific surgical techniques and modifications in enhancing biliary reconstruction success, such as basin-shaped hepaticojejunostomy portoenterostomy and a modified hepato-jejunostomy technique using internal and external stents.^{135,139}

Q4 Is there a role for routine imaging (computed tomography (CT), magnetic resonance (MR) cholangiography) in high-risk patients for bile leaks?

Routine use of advanced imaging modalities such as CT, MR, or single-photon emission computed tomography (SPECT) is not recommended for asymptomatic high-risk patients due to their cost and limited availability in most healthcare settings. For patients with suspected or detected bile leaks, the threshold for performing radiological imaging should be low. The choice of modality should be guided by the resources available at the treating centre.

Recommendation: conditional | median level of evidence: 3 (range: 3)

Expert panel comment

Patients with suspected bile leakage originating from the resection margin or a central bile duct, whether presenting with high or low output, benefit from prompt radiological detection using, depending on local availability, ultrasound, CT scan or more specialized imaging like bile-specific Gadolinium-enhanced MR cholangiography (Gd-MR)^{140–142} or Mebrofenin SPECT-CT.¹⁴³ MR cholangiography, particularly contrast-enhanced MR, is especially reliable for identifying bile leaks following hepatobiliary surgery and may reduce the need for other, potentially risky invasive diagnostic procedures.¹⁴²

The definition and grading of post-hepatectomy bile-leak has been defined by the International Study Group of Liver Surgery (ISGLS)¹³² and allows objective comparison between the degree and severity of bile leaks. The following risk factors for postoperative bile leakage are widely accepted and have been identified by most authors: a) preoperative: anaemia, liver cirrhosis, mechanical jaundice, chemotherapy with anti-VEGF-antibodies

(Bevacizumab), positive bacterial culture (bile swab) and re-operations (for other reasons than bile leakage). b) Intra-operative and surgical risk factors: major or complex hepatic resections including meso-hepatectomies, vascular resections or reconstructions, re-resections and central bile duct resections. Additional risk factors are operation time, blood loss and Pringle-manoeuvre.^{140–143}

Q5 What is the role of endoscopy and interventional radiology in patients with clinically relevant bile leakage after hepatectomy?

In case of a clinically relevant bile leakage (>100 ml bile/24 h) when a drain is in place, an early (endoscopic retrograde cholangiopancreatography) ERCP and stent placement seems sensible as a strategy to resolve bile leakage quickly. When the patient has a bilioma with no drain in place, the first step should be percutaneous drainage; ERCP/stent should follow on the same conditions as above.

Recommendation: conditional | median level of evidence: 2- (1+ to 4)

Expert panel comment

Management of bile leakage relies strongly on locally available resources and experience. Recommendations are made in the context of healthcare systems where all available interventions are in place. This might not always be the case in clinical practice; therefore, recruitment of external expertise might be needed occasionally. RCTs comparing early vs late ERCP/Stent could so far not conclusively answer which approach is superior to the other.^{144–150}

Topic 5: post-hepatectomy haemorrhage (PHH)

Q1 What is the role of pre-operative strategies to reduce blood loss associated with liver resection (fat-reducing diet, nutritional changes, hypovolaemic phlebotomy)?

In patients who are overweight or obese, the implementation of a low-calorie, low-fat diet should be considered, provided that the surgical timeline permits adequate intervention. Pre-operative management of metabolic dysfunction-associated steatotic liver disease (MASLD) should align with the recommendations of the EASL–EASD–EASO Clinical Practice Guidelines. In selected patients at high risk of significant intra-operative bleeding, pre-operative hypovolaemic phlebotomy may be considered as a strategy to reduce transfusion requirements.

Recommendation: strong | median level of evidence: 2- (range: 1++ to 3)

Expert panel comment

There is growing evidence that targeted pre-operative strategies can reduce blood loss in patients undergoing liver resection, particularly those with MASLD. Patients with MASLD have been shown to have a 1.8-fold higher risk of postoperative bleeding, as highlighted in a recent meta-analysis^{151,152} underscoring the need for tailored perioperative approaches.

Nutritional interventions, especially short-term dietary restrictions, have demonstrated promising results. A small RCT found that a one-week, 800-kcal/day diet with 20 g fat and 70 g protein significantly reduced intra-operative blood loss¹⁵³. These findings are supported by retrospective data showing similar benefits with calorie- and fat-restricted diets.¹⁵⁴ Such regimens likely reduce hepatic steatosis and liver volume, improving resectability and surgical safety.

Additionally, hypovolaemic phlebotomy has re-emerged as a strategy to control intra-operative bleeding. Meta-analytic data and the recent PRICE-2 trial both demonstrated that pre-operative venesection reduces transfusion requirements without increasing complication rates.^{155,156} These findings support its selective use in high-risk patients.

Q2 What is the role of anaesthetic management to reduce blood loss in patients undergoing hepatectomy?

Intraoperative low central venous pressure (CVP) should be maintained until completion of parenchymal transection to reduce blood loss and transfusion requirements. No specific anaesthetic agent is recommended. Tranexamic acid is not advised in liver resection for cancer as it does not decrease blood loss during liver resection. The role of acute normovolaemic haemodilution remains unclear.

Recommendation: moderate to strong| median level of evidence: 2++ (range: 1+ to 2-)

Expert panel comment

Anaesthetic expertise in liver surgery is critical in reducing intraoperative blood loss during liver resection. Data from meta-analyses of RCTs supports the use of low CVP until completion of parenchymal transection. This strategy significantly reduces blood loss, shortens operating time, lowers transfusion requirements, and decreases postoperative complications, and is considered both safe and effective in adult patients undergoing hepatectomy.¹⁵⁷

While certain anaesthetic agents may influence hepatic blood flow and mitigate ischaemia-reperfusion injury, particularly volatile inhalational agents metabolised by the liver, current data do not support a definitive recommendation for one agent over another in terms of bleeding outcomes. More targeted studies are needed to establish clear guidance.¹⁵⁸

The use of tranexamic acid (TXA) during liver resection for cancer is not supported by recent evidence. The HeLix trial ($n = 2145$) found no reduction in intraoperative or early post-operative bleeding, transfusion needs, or venous thromboembolism. Notably, TXA was associated with a higher rate of complications.¹⁵⁹

The role of acute normovolaemic haemodilution (ANH) remains uncertain. While ANH may reduce the proportion of patients requiring transfusion, it does not appear to lower total transfused volume or improve postoperative outcomes, including morbidity and mortality.^{157,159}

Q3 What is the role for intra-operative liver inflow occlusion in reducing intra-operative blood loss?

Inflow occlusion techniques such as the Pringle maneuver (PM) and hemihepatic occlusion (HHO) can reduce intra-operative blood loss and should be considered when major bleeding is anticipated. While intermittent Pringle occlusion is safe for up to 120 min, prolonged clamping may increase the risk of ischemia-reperfusion injury, particularly in patients with underlying liver dysfunction.

Recommendation: moderate| median level of evidence: 2+ (range: 1++ to 2-)

Expert panel comment

Intra-operative liver inflow occlusion, most commonly via the PM, is a widely adopted technique to minimize blood loss during hepatic transection. Comparative studies have evaluated PM against HHO, but findings remain mixed.

A 2011 meta-analysis of four RCTs involving 338 patients found no significant differences in intraoperative outcomes between PM and HHO, aside from higher aspartate aminotransferase (AST) levels on postoperative day one in the PM group, indicating transient hepatic injury.¹⁶⁰ A 2019 meta-analysis including eight RCTs and 688 patients reported significantly lower perioperative morbidity in the HHO group. However, liver-specific complications such as bile leak, PHLF, postoperative hemorrhage, and transfusion rates showed no significant differences, though transfused volume per patient was higher in the PM group.¹⁶¹

There is also some evidence from a propensity-matched analysis suggesting that patients with normal liver function may experience worse outcomes with PM compared to no occlusion, potentially due to ischemia-reperfusion injury.¹⁶²

While inflow occlusion is effective in limiting blood loss, it is not without risk. Prolonged continuous clamping can lead to ischemia-reperfusion injury, particularly in patients with liver dysfunction. For this reason, intermittent PM—typically in 20-min cycles with 5-min reperfusion intervals—has been shown to

be safe for total occlusion times up to 120 min, particularly in patients with preserved liver function.^{161,163}

Q4 Is there an influence of parenchymal transection technique on blood loss?

Meta-analyses of RCTs suggest that bipolar energy devices may reduce intraoperative blood loss during parenchymal transection. However, due to the wide range of available techniques, the limited number of direct comparisons, and methodological variability across studies, no specific recommendation can currently be made regarding the optimal transection technique.

Recommendation: moderate | median level of evidence: 1+ (range: 1++ to 2-)

Expert panel comment

Parenchymal transection represents a critical phase of partial hepatectomy, with the potential to significantly influence intraoperative blood loss and postoperative complications. Multiple techniques and devices have been developed to facilitate safe and efficient transection. These range from the traditional clamp-crushing technique, which involves blunt dissection of the parenchyma with a Kelly clamp followed by clipping or ligation of vascular and biliary structures, to energy-based devices such as bipolar cautery, ultrasonic dissectors, and vascular staplers.

Over the past two decades, numerous RCTs have compared these techniques with respect to key perioperative outcomes, particularly intraoperative blood loss and postoperative complication rates—most notably bile leakage. However, individual trials often yield conflicting results due to differences in methodology and patient populations.

A 2020 systematic review and network meta-analysis of 22 RCTs sought to synthesize this evidence and found that bipolar energy devices were associated with the greatest reduction in intraoperative blood loss.¹⁶⁴ Nevertheless, the authors noted that cross-trial comparisons were limited by heterogeneity in surgical practices such as CVP management and use of vascular inflow occlusion.

In conclusion, while certain transection techniques may offer modest advantages, particularly in reducing blood loss, the overall evidence remains inconclusive due to variability in study design and surgical context.

Q5 What is the value of haemostatic agents to reduce blood loss in patients undergoing hepatectomy?

No specific recommendations can be made as to the use of topical haemostats. The time to haemostasis is shorter when using topical haemostatic agents (fibrin sealant) but there is no

significant difference in peri operative blood transfusion. There is no evidence of one individual topical sealant being superior to another.

Recommendation: moderate | median level of evidence: 1+ (range: 1++ to 2-)

Expert panel comment

Topical haemostatic agents, including matrix-based products, fibrin sealants, and their combination—known as carrier-bound fibrin sealants—are widely used in liver surgery for secondary haemostasis. Despite their popularity, the level of evidence supporting their effectiveness in reducing blood loss remains low.

A meta-analysis evaluating carrier-bound fibrin sealants in open and laparoscopic liver surgery, as well as transplantation, included RCT, large retrospective cohort studies, and case-control studies.¹⁶⁵ While time to haemostasis was shorter in the carrier-bound fibrin sealant group, the use of these agents did not significantly affect the risk of intraoperative blood transfusion, postoperative fluid collections, or bile leakage.

Additionally, a network meta-analysis of 20 RCTs comparing different topical haemostatic agents reported that fibrin glue and fibrin patches were the most effective in achieving haemostasis at both 4 and 10 min.¹⁶⁶ However, no significant differences were observed among agents in terms of overall blood loss, transfusion requirements, bile leak, postoperative complications, reoperation rates, or mortality.

Overall, while certain agents may improve time to haemostasis, their routine use does not appear to influence key perioperative outcomes in liver surgery.¹⁶⁵

Q6 Does minimally-invasive surgery influence blood loss in patients undergoing hepatectomy?

Minimally-invasive surgery (MIS) does not reduce blood loss compared to open liver surgery for minor nor for major hepatectomy. MIS liver surgery may be associated with less blood loss in some conditions (i.e. HCC) based on non-randomized observations.

Recommendation: strong (minor/major hepatectomy), conditional (HCC) | median level of evidence: 1- (range: 1++ to 2-)

Expert panel comment

In selected patients, MIS may decrease perioperative blood loss compared to the open approach. However, this is only found when looking at specific resections (i.e. HCC) and, most prominently in left lateral resections and in uncontrolled data comparisons.^{167–169} Neither the Orange III RCT,¹⁷⁰ nor the COMET RCT,¹⁷¹ demonstrated statistically significant difference in blood loss attributable to the surgical approach.

Topic 6: post-hepatectomy liver failure (PHLF)

Q1 Is there a clinical relevance to sub classify PHLF by the main underlying pathomechanism?

PHLF involves complex and heterogenous pathophysiological mechanisms and classification as primary and secondary PHLF can be considered, because it impacts on preventive and therapeutic interventions and prognosis.

Recommendation: conditional | median level of evidence: 2+ (range: 1+ to 3)

Expert panel comment

Primary PHLF defines liver failure (increased International normalized ratio (INR) and Bilirubin according to local laboratory normal value ranges) and clinical signs of hepatic encephalopathy that occurs within days after liver resection (usual time frame within 3 weeks)¹⁷² Primary PHLF includes „small-for-size“ PHLF, in which the liver remnant is insufficient to maintain essential hepatic function, and iatrogenic PHLF, in which surgical injury of vital structures or excessive blood loss prevent sufficient function of the liver remnant.^{172,173} Secondary PHLF occurs after an initial partial stabilization/recovery of hepatic function (usual time frame within 3 months after surgery) and is often complicated/aggravated by infections or hepatic vasculature thrombosis.^{172,174}

While postoperative outcome research has predominantly focused on the first 90 postoperative days, it should be noted that delayed hepatic decompensation after this period can occur, particularly if infectious complications arise. Recent evidence supports this concept, as impaired preoperative liver function and/or occurrence of non-PHLF postoperative hepatic dysfunction is associated with long term survival after hepatic resection.^{175–177} This is particularly important as treatment strategies might differ between PHLF types. For example, Molecular Adsorbent Recirculating System (MARS) achieved better results as therapy of primary PHLF¹⁷⁸ (especially if used early), while aggressive infection management is key in avoiding secondary PHLF.¹⁷² Further research is required to strengthen the evidence of subclassifying PHLF, particularly for defining delayed PHLF occurring after 90 days.

Q2 Should patients suffering from PHLF be monitored beyond 90 days after hepatectomy?

Primary and secondary PHLF usually occur within 90 days after surgery. Based on indirect evidence, PHLF monitoring beyond 90 days can be considered for patients who have recovered from PHLF independent of oncological follow-up.

Recommendation: conditional | median level of evidence: 4 (range: 2+ to 5)

Expert panel comment

Primary and secondary PHLF occur within 90 days after liver resection. Delayed PHLF after 90 days — after initial stabilization/recovery of liver function — may occur, but further research is required to strengthen evidence for its definition, prediction and management recommendations.

Recent evidence suggests that preoperative liver function and/or the development of postoperative non-PHLF hepatic dysfunction represents a critical predictor for long term survival after hepatic resection.^{175–177,179} Delayed PHLF may account for a significant proportion of post-resection mortality within 1 year, but strong supporting evidence is yet missing. Monitoring beyond 90 days can be considered for those who recovered from PHLF, but also for high-risk patients, who should be defined as per recent consensus guidelines.¹⁵

Q3 Which postoperative markers should be considered to monitor postoperative liver function recovery in patients at high risk of PHLF?

The following parameters should be used to monitor liver function in patients at high risk of PHLF: bilirubin, INR, albumin, lactate, ammonia levels (in patients with signs of hepatic encephalopathy), AST, alanine aminotransferase (ALT), glucose and temperature. Furthermore, infection monitoring by complete blood count (CBC) and CRP should be performed, as well as kidney function monitoring using creatinine.

Recommendation: strong | median level of evidence: 2+ (range: 2+ to 4)

Expert panel comment

After hepatic resection the following parameters should be used to monitor liver function in patients at high risk of PHLF: bilirubin, INR, albumin, ammonia levels (in patients with signs of hepatic encephalopathy), glucose, temperature, transaminases (AST/ALT), infection markers (leukocytes, procalcitonin and CRP), lactate and kidney function (creatinine).

If clinically indicated, radiological assessment of the liver perfusion (Ultrasound or CT scan) can be helpful to identify vascular complications. In this context, high risk patients should be defined as per recent consensus guidelines.¹⁵

Q4 Is there a value for early postoperative markers to predict PHLF?

Several biomarkers (e.g. lactate, phosphate, platelets) have been reported to allow for early postoperative identification of PHLF and their measurement can be considered in high-risk patients after hepatic resection to initiate supportive treatment early. There are no comparative studies, and future research is needed to identify the optimal biomarker.

Recommendation: conditional | median level of evidence: 2+ (range: 2+ to 4)

Expert panel comment

The postoperative parameters reported to be useful for early identification of PHLF patients include lactate levels,^{180–182} fibrosis-4 (FIB-4) score,¹⁸³ platelets,¹⁸⁴ ICG clearance (intra- and postoperative),^{185,186} phosphate,^{187,188} Antithrombin III and CRP,¹⁸⁹ scintigraphy,^{190,191} thrombopoietin,¹⁹² intrahepatic neutrophil accumulation.¹⁹³ At this point in time direct comparisons of these markers are missing. Accordingly, no recommendation can be made regarding which parameter is superior. Given that supportive treatments (particularly for primary PHLF) should be initiated early,¹⁷⁸ the biomarker identification and comparison should become a key area of future research. Moreover, these markers will be instrumental for establishing and eventually initiating therapeutic interventions in the future.^{194,195} In this context, high risk patients should be defined as per recent consensus guidelines.¹⁵

Q5 Which treatments are available to support hepatic regeneration in PHLF?

Currently, no specific therapeutic interventions are available or approved to promote liver regeneration and avoid or treat progression of PHLF.

Recommendation: strong | median level of evidence: 2+ (range: 1+ to 5)

Expert panel comment

Supportive care and aggressive management of potential causes of PHLF are recommended. While several treatments have been explored, no specific therapy is currently available. Antioxidant drugs, such as N-acetylcysteine, have shown mixed results for treatment of PHLF and their use is controversial.^{196,197} Bio-artificial devices, mesenchymal stem cells as well as new molecular targets are currently being explored in patients at risk of or already suffering from PHLF. Portal pressure modulators like terlipressin and somatostatin have demonstrated mixed results as well. Surgical modulation of portal pressure by splenectomy or splenic artery ligation have also been implicated to decrease PHLF.¹⁹⁸ MARS and plasma exchange therapies have also been evaluated with inconsistent results.^{199–201} Liver transplantation is an effective treatment for fulminant PHLF but is seldomly possible given patient factors like oncological diagnosis or infections etc.^{202–204} Preoperative or early postoperative identification of high-risk patients remains key to enabling timely initiation of supportive treatments.

Discussion

Key message

This consensus paper builds upon and complements the E-AHPBA–ESSO–ESSR Innsbruck Consensus Guidelines on preoperative liver function assessment published in BJS in 2023,¹⁵ providing a continuum of guidance from preoperative

evaluation to postoperative care. Developed by a multidisciplinary expert panel, it aims to reduce practice variability and improve patient outcomes by synthesizing the best available evidence with the opinions and clinical knowledge of experts in the field. It provides a structured framework for peri- and postoperative management in patients undergoing liver resection, addressing six critical domains: thromboprophylaxis, antibiotics, prehabilitation, bile leak, bleeding, and liver failure. A total of 32 statements were generated.

Even though the review of the literature revealed that most of the available evidence informing these recommendations is limited in both quantity and quality, the expert panel has identified key practices that, if implemented, are believed to support improved outcomes in peri- and postoperative care following liver surgery.

Limitations

This consensus recommendation is subject to several limitations. First, the heterogeneity of definitions, particularly regarding perioperative complications but also regarding liver surgery procedures and indications. Then, the variability in the application of interventions such as prehabilitation, thromboprophylaxis, and bile leak management, limit generalizability. Several recommendations also rely on advanced diagnostics and therapeutics not universally available, challenging implementation across diverse healthcare settings. Many of the recommendations are based on low-to moderate-quality evidence, often derived from retrospective analyses, institutional protocols, or expert opinion, due to the lack of high-quality RCTs in key areas. As with many expert consensus processes, the strength of the recommendations reflects both the current evidence base and clinical judgment.

It is further necessary to consider the demographic and geographical composition, as well as the clinical specialty, of the involved experts. The vast majority of the faculty were drawn from high-income countries and were engaged in high-volume hospitals with specialised hepatobiliary surgery units. It is therefore evident that the clinical experience and available resources of the aforementioned units may not be directly transferable to low- or medium-volume surgical units, or to healthcare settings characterised by resource constraints. Consequently, the implementation of any recommendation outlined in this article is contingent upon the specific requirements, experience, and available resources of the respective institutions or countries.

The consensus threshold of 70 % allows the potential inclusion of statements with limited. Further limitations arise from the methodology applied. Incorporating a modified Delphi approach and expert opinions carries some risk of bias in the interpretation of the literature review, especially since many available publications were indeed published by members of the consensus group. Involvement of a neutral validation committee and full transparency through declaration of conflict of interest for each co-author addressed this potential limitation.

Finally, long-term outcome data are sparse, particularly in relation to late postoperative complications, underscoring the need for prospective, multicentre studies to validate and refine these recommendations.

Implementation considerations

Clinicians are encouraged to adapt these consensus recommendations according to institutional resources, local expertise, and patient populations. The recommendations are designed to align with other principles, such as Enhanced Recovery After Surgery (ERAS)⁶ and EASL–EASD–EASO Clinical Practice Guidelines.¹⁵² They are also implementable within both academic and community practice environments. Given the level of available evidence and the current rate of innovation in perioperative

management of surgical patients, including those undergoing liver surgery, the authors estimate that these recommendations will remain relevant for approximately five years.

Future research topics

To strengthen future versions of these recommendations and resolve ongoing uncertainties, several evidence gaps on perioperative management of patients undergoing liver resections have been identified throughout this consensus development process. These clinically relevant areas represent future research topics that could be addressed in a common, multicentric effort with the aim to improve clinical outcomes and standardize the perioperative management of patients undergoing complex liver resections. Equally, some topics were identified where additional research

Table 1 Priority areas for future research

Area	Rationale and unmet needs
Thromboprophylaxis following portal vein reconstruction (PVR) and liver resection	There is currently no strong evidence on specific anticoagulation protocols after PVR and hepatectomy available. Trials should define patients at risk for portal vein thrombosis (PVT) and compare effectiveness and costs with regard to duration of thromboprophylaxis and pharmacologic agents (e.g., LMWH vs. UFH).
PVT classification	The application of the 30-day threshold (according to transplant surgery) is pragmatic but arbitrary. Studies should focus on a possible pathophysiological distinction between early and late PVT. Definitions are needed that include time point, extent as well as duration of existence of PVT in order to optimize intervention and surveillance strategies.
Optimal duration of perioperative antibiotic therapy	There is uncertainty about sufficiency of short-course vs. prolonged antibiotic regimens. Future studies should identify preoperative, intraoperative, and postoperative risk factors to develop predictive models and to investigate potential cost-saving effects of different treatment regimens.
Prehabilitation	Standardized metrics for nutritional risk or optimal prehabilitation components are lacking. Trials should focus on assessment tools in order to improve patient selection, outcome, and economic challenges.
Bile leak risk modeling	Complex hepatectomies with biliary resections experience high morbidity. Studies aimed at risk modeling could guide selective drainage and imaging postoperatively.
Anastomotic stenting	There is no evidence supporting the use of trans-anastomotic stents even though it is still common practice in high-risk anastomoses. There is a need to develop predictive models based on preoperative, intraoperative, and postoperative risk factors and to identify potential cost-saving effects when applying stents in a certain cohort of patients.
Biomarker-based prediction of post-hepatectomy liver failure (PHLF)	Several markers are currently available and in clinical use. Comparative studies are needed to define the best suitable biomarker for clinical practice.

Table 2 Areas where future research May Be futile

Area	Rationale
Routine use of intraoperative drains	Multiple randomized controlled trials (RCTs) have shown that there is no benefit of drains in standard hepatectomies. Enhanced recovery protocols strongly advocate against routine use of drains as they contradict early mobilization.
Minimally-invasive surgery (MIS) and blood loss	Recent RCTs provide high level of evidence that MIS does not result in significantly lower blood loss compared to open surgery.

may yield limited clinical benefit, and that the authors define as “futile” since enough good evidence has been achieved and no future efforts should be invested in those areas (Tables 1 and 2).

Final statement

The present consensus paper has been developed for the purpose of supporting clinicians in the delivery of safe, evidence-informed, and standardized care across the perioperative course of liver resection. The text addresses fundamental clinical dilemmas, with the objective of providing clear guidance, thus making it a practical reference for surgical and multidisciplinary teams. It is imperative that ongoing research collaboration, implementation monitoring, and periodic updates are maintained to ensure the continued relevance of the program, and to enhance the outcomes for patients undergoing liver resection.

Acknowledgements

We thank Gerhard Lobenscheg from the patient support group “L(i)eberleben” in Tyrol, Austria, for reviewing the final draft of the manuscript as a patient representative.

Funding sources

This study was supported by an educational grant of BJS.

Data availability statement

Not applicable.

Conflicts of interest

AAS is CEO of the Start Up Capreolos that develops digital solutions for prehabilitation in major surgery. PS receives consulting fees from Tamirna and is co-owner of the patent of HepatomiR is getting an inventor fee by the Medical University of Vienna. All other authors have no conflicts of interest to report.

References

- van Keulen AM, Büttner S, Erdmann JI, Hagendoorn J, Hoogwater FJH, IJzermans JNM *et al.* (2023) Major complications and mortality after resection of intrahepatic cholangiocarcinoma: a systematic review and meta-analysis. *Surgery* 173:973–982.
- van Tuil T, Dhaif AA, Te Riele WW, van Ramshorst B, van Santvoort HC. (2019) Systematic review and meta-analysis of liver resection for colorectal metastases in elderly patients. *Dig Surg* 36:111–123.
- Kong J, Li G, Chai J, Yu G, Liu Y, Liu J. (2021) Impact of postoperative complications on long-term survival after resection of hepatocellular carcinoma: a systematic review and meta-analysis. *Ann Surg Oncol* 28:8221–8233.
- Chok KS, Ng KK, Poon RT, Lo CM, Fan ST. (2009) Impact of post-operative complications on long-term outcome of curative resection for hepatocellular carcinoma. *Br J Surg* 96:81–87.
- Lassen K, Nymo LS, Olsen F, Brudvik KW, Fretland Å, Søreide K. (2019) Contemporary practice and short-term outcomes after liver resections in a complete national cohort. *Langenbecks Arch Surg* 404:11–19.
- Joliat GR, Kobayashi K, Hasegawa K, Thomson JE, Padbury R, Scott M *et al.* (2023) Guidelines for perioperative care for liver surgery: enhanced recovery after surgery (ERAS) society recommendations 2022. *World J Surg* 47:11–34.
- Baltatzis M, Low R, Stathakis P, Sheen AJ, Siriwardena AK, Jamdar S. (2017) Efficacy and safety of pharmacological venous thromboembolism prophylaxis following liver resection: a systematic review and meta-analysis. *HPB* 19:289–296.
- Tanner B, Lu S, Zervoudakis G, Woodwyk A, Munene G. (2018) Coagulation profile following liver resection: does liver cirrhosis affect thromboelastography? *Am J Surg* 215:406–409.
- Shinkawa H, Tanaka S, Takemura S, Amano R, Kimura K, Nishioka T *et al.* (2019) Giving short-term prophylactic antibiotics in patients undergoing open and laparoscopic hepatic resection. *Ann Gastroenterol Surg* 3:506–514.
- O’Leary L, Lal P, Gilbert T, Quinn M, Díaz-Nieto R, Jones R *et al.* (2024) HPB SO02 - cutting surgical site infection rates in perihilar cholangiocarcinoma with informed changes to perioperative antibiotic prophylaxis: a quality improvement project in a UK hepatopancreatobiliary tertiary centre. *Br J Surg* 111(Supplement 9).
- Burchard PR, Dave YA, Loria AP, Parikh NB, Pineda-Solis K, Ruffolo LI *et al.* (2022) Early postoperative ERAS compliance predicts decreased length of stay and complications following liver resection. *HPB* 24: 1425–1432.
- Spetzler VN, Schepers M, Pinnschmidt HO, Fischer L, Nashan B, Li J. (2019) The incidence and severity of post-hepatectomy bile leaks is affected by surgical indications, preoperative chemotherapy, and surgical procedures. *Hepatobiliary Surg Nutr* 8:101–110.
- Rahbari NN, Garden OJ, Padbury R, Maddern G, Koch M, Hugh TJ *et al.* (2011) Post-hepatectomy haemorrhage: a definition and grading by the international study group of liver surgery (ISGLS). *HPB* 13:528–535.
- Lim C, Dokmak S, Farges O, Aussilhou B, Sauvanet A, Belghiti J. (2014) Reoperation for post-hepatectomy hemorrhage: increased risk of mortality. *Langenbecks Arch Surg* 399:735–740.
- Primavesi F, Maglione M, Cipriani F, Denecke T, Oberkofler CE, Starlinger P *et al.* (2023) Stättner S. E-AHPBA-ESSO-ESSR innsbruck consensus guidelines for preoperative liver function assessment before hepatectomy. *Br J Surg* 110:1331–1347.
- Abu Hilal M, Aldrighetti L, Dagher I, Edwin B, Troisi RI, Alikhanov R *et al.* (2018) The Southampton consensus guidelines for laparoscopic liver surgery: from indication to implementation. *Ann Surg* 268:11–18.
- SIGN 50: A guideline developer’s handbook. Scottish Intercollegiate Guidelines Network (SIGN). www.sign.ac.uk/guidelines/fulltext/50/index.html [Accessed: 1.March.2021].
- Dalkey NCHO. (1963) An experimental application of the Delphi method to the use of experts. *Manag Sci* 9:458–467.
- Nasa P, Jain R, Juneja D. (2021) Delphi methodology in healthcare research: how to decide its appropriateness. *World J Methodol* 11: 116–129.
- AGREE II-GRS Instrument Approval of guidelines research and evaluation (AGREE). <http://www.agreetrust.org/resource-centre/agree-iigrs-instrument/>. [Accessed 1 March 2021].
- Tzeng CW, Katz MH, Fleming JB, Pisters PW, Lee JE, Abdalla EK *et al.* (2012) Risk of venous thromboembolism outweighs post-hepatectomy bleeding complications: analysis of 5651 national surgical quality improvement program patients. *HPB* 14:506–513.
- Aloia TA, Geerts WH, Clary BM, Day RW, Hemming AW, D’Albuquerque LC *et al.* (2016) Venous thromboembolism prophylaxis in liver surgery. *J Gastrointest Surg* 20:221–229.

23. Edwards MA, Hussain MWA, Spaulding AC, Brennan E, Colibaseanu D, Stauffer J. (2023) Venous thromboembolism and bleeding after hepatectomy: role and impact of risk adjusted prophylaxis. *J Thromb Thrombolysis* 56:375–387.
24. Weiss MJ, Kim Y, Ejaz A, Spolverato G, Haut ER, Hirose K *et al.* (2014) Venous thromboembolic prophylaxis after a hepatic resection: patterns of care among liver surgeons. *HPB* 16:892–898.
25. Ainoa E, Uutela A, Nordin A, Mäkisalo H, Sallinen V. (2021) Pre- vs. postoperative initiation of thromboprophylaxis in liver surgery. *HPB* 23:1016–1024.
26. Hull RD, Pineo GF, MacIsaac S. (2001) Low-molecular-weight heparin prophylaxis: preoperative versus postoperative initiation in patients undergoing elective hip surgery. *Thromb Res* 101:V155–V162.
27. Lavikainen LI, Guyatt GH, Sallinen VJ, Karanicolas PJ, Couban RJ, Singh T *et al.* (2024) Systematic reviews and meta-analyses of the procedure-specific risks of thrombosis and bleeding in general abdominal, colorectal, upper gastrointestinal, and hepatopancreatobiliary surgery. *Ann Surg* 279:213–225.
28. Barbar S, Noventa F, Rossetto V, Ferrari A, Brandolin B, Perlati M *et al.* (2010) A risk assessment model for the identification of hospitalized medical patients at risk for venous thromboembolism: the padua prediction score. *J Thromb Haemostasis* 8:2450–2457.
29. Miyazaki M, Shimizu H, Ohtuka M, Kato A, Yoshitomi H, Furukawa K *et al.* (2017) Portal vein thrombosis after reconstruction in 270 consecutive patients with portal vein resections in hepatopancreatobiliary (HPB) surgery. *Am J Surg* 214:74–79.
30. Molina V, Sampson J, Ferrer J, Díaz A, Ayuso JR, Sánchez-Cabús S *et al.* (2017) Surgical treatment of perihilar cholangiocarcinoma: early results of en bloc portal vein resection. *Langenbecks Arch Surg* 402:95–104.
31. Ejaz A, Spolverato G, Kim Y, Lucas DL, Lau B, Weiss M *et al.* (2014) Defining incidence and risk factors of venous thromboembolism after hepatectomy. *J Gastrointest Surg* 18:1116–1124.
32. Onda S, Furukawa K, Shirai Y, Hamura R, Horiuchi T, Yasuda J *et al.* (2020) New classification-oriented treatment strategy for portal vein thrombosis after hepatectomy. *Ann Gastroenterol Surg* 4:701–709.
33. Zhou YM, Chen ZY, Li XD, Xu DH, Su X, Li B. (2016) Preoperative antibiotic prophylaxis does not reduce the risk of postoperative infectious complications in patients undergoing elective hepatectomy. *Dig Dis Sci* 61:1707–1713.
34. Hirokawa F, Hayashi M, Miyamoto Y, Asakuma M, Shimizu T, Komeda K *et al.* (2013) Evaluation of postoperative antibiotic prophylaxis after liver resection: a randomized controlled trial. *Am J Surg* 206:8–15.
35. Liew YX, Chian DM, Koh YX, Chlebicki MP. (2021) Reducing prophylactic antibiotics use in laparoscopic cholecystectomy and liver resections. *Eur J Gastroenterol Hepatol* 33:533–534.
36. Steccanella F, Amoretti P, Barbieri MR, Bellomo F, Puziello A. (2022) Antibiotic prophylaxis for hepato-biliopancreatic Surgery-A systematic review. *Antibiotics (Basel)* 11.
37. Matsumura M, Saiura A, Inoue Y, Ishizawa T, Mise Y, Takahashi Y. (2016) High rate of organ/space surgical site infection after hepatectomy with preexisting bilioenteric anastomosis. *World J Surg* 40:937–945.
38. Liakina V. (2023) Antibiotic resistance in patients with liver cirrhosis: prevalence and current approach to tackle. *World J Clin Cases* 11: 7530–7542.
39. Dirchwolf M, Marciano S, Martínez J, Ruf AE. (2018) Unresolved issues in the prophylaxis of bacterial infections in patients with cirrhosis. *World J Hepatol* 10:892–897.
40. Sugawara G, Yokoyama Y, Ebata T, Mizuno T, Yagi T, Ando M *et al.* (2018) Duration of antimicrobial prophylaxis in patients undergoing major hepatectomy with extrahepatic bile duct resection: a randomized controlled trial. *Ann Surg* 267:142–148.
41. Ishioka K, Hokuto D, Nomi T, Yasuda S, Yoshikawa T, Matsuo Y *et al.* (2018) Significance of bacterial culturing of prophylactic drainage fluid in the early postoperative period after liver resection for predicting the development of surgical site infections. *Surg Today* 48:625–631.
42. Yang T, Tu PA, Zhang H, Lu JH, Shen YN, Yuan SX *et al.* (2014) Risk factors of surgical site infection after hepatic resection. *Infect Control Hosp Epidemiol* 35:317–320.
43. Okamura K, Tanaka K, Miura T, Nakanishi Y, Noji T, Nakamura T *et al.* (2017) Randomized controlled trial of perioperative antimicrobial therapy based on the results of preoperative bile cultures in patients undergoing biliary reconstruction. *J Hepatobiliary Pancreat Sci* 24: 382–393.
44. Suc B, Panis Y, Belghiti J, Fékété F. (1992) 'Natural history' of hepatectomy. *Br J Surg* 79:39–42.
45. Stack CM, Gold HS, Wright SB, Baldini LM, Snyder GM. (2018) Perioperative antimicrobial prophylaxis and prevention of hepatobiliary surgical site infections. *Infect Control Hosp Epidemiol* 39: 1037–1041.
46. Chacon E, Eman P, Dugan A, Davenport D, Marti F, Ancheta A *et al.* (2019) Effect of operative duration on infectious complications and mortality following hepatectomy. *HPB* 21:1727–1733.
47. Nakayama H, Takayama T, Okubo T, Higaki T, Midorikawa Y, Moriguchi M *et al.* (2014) Subcutaneous drainage to prevent wound infection in liver resection: a randomized controlled trial. *J Hepatobiliary Pancreat Sci* 21:509–517.
48. Shen J, Ni Z, Qian Y, Wang B, Zheng S. (2020) Clinical prediction score for superficial surgical site infections: real-life data from a retrospective single-centre analysis of 812 hepatectomies. *Int Wound J* 17:16–20.
49. Dong ZM, Chidi AP, Goswami J, Han K, Simmons RL, Rosengart MR *et al.* (2015) Prior inpatient admission increases the risk of postoperative infection in hepatobiliary and pancreatic surgery. *HPB* 17: 1105–1112.
50. Isik O, Kaya E, Sarkut P, Dundar HZ. (2015) Factors affecting surgical site infection rates in hepatobiliary surgery. *Surg Infect* 16:281–286.
51. Kokudo T, Uldry E, Demartines N, Halkic N. (2015) Risk factors for incisional and organ space surgical site infections after liver resection are different. *World J Surg* 39:1185–1192.
52. Takahashi Y, Takesue Y, Fujiwara M, Tatsumi S, Ichiki K, Fujimoto J *et al.* (2018) Risk factors for surgical site infection after major hepatobiliary and pancreatic surgery. *J Infect Chemother* 24: 739–743.
53. Tang H, Lu W, Yang Z, Jiang K, Chen Y, Lu S *et al.* (2017) Risk factors and long-term outcome for postoperative intra-abdominal infection after hepatectomy for hepatocellular carcinoma. *Medicine (Baltim)* 96e6795.
54. Morikawa T, Ishida M, Iseki M, Aoki S, Hata T, Kawaguchi K *et al.* (2021) Liver resections in patients with prior bilioenteric anastomosis are predisposed to develop organ/space surgical site infections and biliary leakage: results from a propensity score matching analysis. *Surg Today* 51:526–536.
55. Sadamori H, Yagi T, Shinoura S, Umeda Y, Yoshida R, Satoh D *et al.* (2013) Risk factors for organ/space surgical site infection after

- hepatectomy for hepatocellular carcinoma in 359 recent cases. *J Hepatobiliary Pancreat Sci* 20:186–196.
56. Yasuda S, Hokuto D, Kamitani N, Matsuo Y, Doi S, Nakagawa K *et al.* (2023) Pre- and postoperative C-reactive protein as a risk factor of organ/space surgical site infection after hepatectomy. *Langenbecks Arch Surg* 408:13.
 57. Yusa T, Yamashita YI, Nitta H, Nakao Y, Itoyama R, Kitano Y *et al.* (2022) Efficacy of ring drape and unused sterile instruments for incisional SSI after hepatectomy. *Anticancer Res* 42:1007–1012.
 58. Gupta S, Sinha PK, Patil NS, Mohapatra N, Sindwani G, Garg N *et al.* (2022) Randomized control trial on perioperative antibiotic prophylaxis in live liver donors: are three doses enough? *J Hepatobiliary Pancreat Sci* 29:1124–1132.
 59. Murtha-Lemekhova A, Fuchs J, Teroerde M, Chiriac U, Klotz R, Hornuss D *et al.* (2022) Routine postoperative antibiotic prophylaxis offers no benefit after Hepatectomy-A systematic review and meta-analysis. *Antibiotics (Basel)* 11.
 60. Sakoda M, Iino S, Mataka Y, Kawasaki Y, Kurahara H, Maemura K *et al.* (2017) Influence of a shorter duration of post-operative antibiotic prophylaxis on infectious complications in patients undergoing elective liver resection. *Surg Infect* 18:149–156.
 61. Takayama T, Aramaki O, Shibata T, Oka M, Itamoto T, Shimada M *et al.* (2019) Antimicrobial prophylaxis for 1 day versus 3 days in liver cancer surgery: a randomized controlled non-inferiority trial. *Surg Today* 49:859–869.
 62. Bednarsch J, Czigany Z, Heij LR, Luedde T, van Dam R, Lang SA *et al.* (2021) Bacterial bile duct colonization in perihilar cholangiocarcinoma and its clinical significance. *Sci Rep* 11:2926.
 63. Chen Z, Jiang H, Wang Y, Liang R, Xu L, Lai J *et al.* (2021) Three-day postoperative antibiotics reduces post-hepatectomy infection rate in hepatitis B virus-related hepatocellular carcinoma. *J Gastroenterol Hepatol* 36:2531–2539.
 64. Gurusamy KS, Naik P, Davidson BR. (2011) Methods of decreasing infection to improve outcomes after liver resections. *Cochrane Database Syst Rev*, CD006933.
 65. Dewulf M, Verrips M, Coolsen MME, Olde Damink SWM, Den Dulk M, Bongers BC *et al.* (2021) The effect of prehabilitation on postoperative complications and postoperative hospital stay in hepatopancreato-biliary surgery a systematic review. *HPB* 23:1299–1310.
 66. Wang B, Shelat VG, Chow JLL, Huey PCW, Low JK, Woon WWL *et al.* (2020) Prehabilitation program improves outcomes of patients undergoing elective liver resection. *J Surg Res* 251:119–125.
 67. Cambriel A, Choisy B, Hedou J, Bonnet MP, Fellous S, Lefevre JH *et al.* (2023) Impact of preoperative uni- or multimodal prehabilitation on postoperative morbidity: meta-analysis. *BJS Open* 7.
 68. Dagorno C, Sommacale D, Laurent A, Attias A, Mongardon N, Levesque E *et al.* (2022) Prehabilitation in hepato-pancreato-biliary surgery: a systematic review and meta-analysis. A necessary step forward evidence-based sample size calculation for future trials. *J Vis Surg* 159:362–372.
 69. Deprato A, Verhoeff K, Purich K, Kung JY, Bigam DL, Dajani KZ. (2022) Surgical outcomes and quality of life following exercise-based prehabilitation for hepato-pancreato-biliary surgery: a systematic review and meta-analysis. *Hepatobiliary Pancreat Dis Int* 21:207–217.
 70. Nakajima H, Yokoyama Y, Inoue T, Nagaya M, Mizuno Y, Kadono I *et al.* (2019) Clinical benefit of preoperative exercise and nutritional therapy for patients undergoing hepato-pancreato-biliary surgeries for malignancy. *Ann Surg Oncol* 26:264–272.
 71. Cambriel A, Choisy B, Hedou J, Bonnet MP, Fellous S, Lefevre JH *et al.* (2023) Impact of preoperative uni- or multimodal prehabilitation on postoperative morbidity: meta-analysis. *BJS open* 7.
 72. Dunne DF, Jack S, Jones RP, Jones L, Lythgoe DT, Malik HZ *et al.* (2016) Randomized clinical trial of prehabilitation before planned liver resection. *Br J Surg* 103:504–512.
 73. Dunne DF, Jones RP, Lythgoe DT, Pilkington FJ, Palmer DH, Malik HZ *et al.* (2014) Cardiopulmonary exercise testing before liver surgery. *J Surg Oncol* 110:439–444.
 74. Kaibori M, Ishizaki M, Matsui K, Nakatake R, Yoshiuchi S, Kimura Y *et al.* (2013) Perioperative exercise for chronic liver injury patients with hepatocellular carcinoma undergoing hepatectomy. *Am J Surg* 206: 202–209.
 75. Kasvis P, Vigano A, Bui T, Carli F, Kilgour R. (2024) Cancer symptom burden negatively affects health-related quality of life in patients undergoing prehabilitation prior to liver resection: results from a 12-week randomized controlled trial. *Appl Physiol Nutr Metabol* 49:64–76.
 76. Sliwinski S, Werneburg E, Faqar-Uz-Zaman SF, Detemle C, Dreilich J, Mohr L *et al.* (2023) A toolbox for a structured risk-based prehabilitation program in major surgical oncology. *Front Surg* 10:1186971.
 77. Zhang H, Zhang L, Wang Y, Zou J, Zhang T. (2023) Sarcopenia predicts prognosis of patients undergoing liver resection for hepatocellular carcinoma: a systematic review and meta-analysis. *Nutr Cancer* 75:776–784.
 78. Kong Q, Gao Q, Li W, Chen Z. (2024) The impact of imaging-diagnosed sarcopenia on long-term prognosis after curative resection for hepatocellular carcinoma: a systematic review and meta-analysis. *Acad Radiol* 31:1272–1283.
 79. March C, Omari J, Thormann M, Pech M, Wienke A, Surov A. (2022) Prevalence and role of low skeletal muscle mass (LSMM) in hepatocellular carcinoma. A systematic review and meta-analysis. *Clin Nutr ESPEN* 49:103–113.
 80. O'Connell RM, O'Neill M, Mg OR, Cb OS, O'Sullivan AW. (2022) Sarcopaenia, obesity, sarcopaenic obesity and outcomes following hepatic resection for colorectal liver metastases: a systematic review and meta-analysis. *HPB* 24:1844–1853.
 81. Thormann M, Omari J, Pech M, Damm R, Croner R, Perrakis A *et al.* (2022) Low skeletal muscle mass and post-operative complications after surgery for liver malignancies: a meta-analysis. *Langenbecks Arch Surg* 407:1369–1379.
 82. Wagner D, Wienerroither V, Scherrer M, Thalhammer M, Faschinger F, Lederer A *et al.* (2023) Value of sarcopenia in the resection of colorectal liver metastases-a systematic review and meta-analysis. *Front Oncol* 13:1241561.
 83. Xu L, Jing Y, Zhao C, Zhang Q, Zhao X, Yang J *et al.* (2020) Preoperative computed tomography-assessed skeletal muscle index is a novel prognostic factor in patients with hepatocellular carcinoma following hepatectomy: a meta-analysis. *J Gastrointest Oncol* 11: 1040–1053.
 84. O'Connell RM, O'Neill M, Ó Ríordáin MG, Ó Súilleabháin CB, O'Sullivan AW. (2022) Sarcopaenia, obesity, sarcopaenic obesity and outcomes following hepatic resection for colorectal liver metastases: a systematic review and meta-analysis. *HPB* 24:1844–1853.

85. Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T *et al.* (2019) Writing group for the european working group on sarcopenia in older people 2 (EWGSOP2) atEGfE. Sarcopenia: revised european consensus on definition and diagnosis. *Age Ageing* 48:601.
86. Cornet M, Lim C, Salloum C, Lazzati A, Compagnon P, Pascal G *et al.* (2015) Prognostic value of sarcopenia in liver surgery. *J Vis Surg* 152: 297–304.
87. Clifford K, Woodfield JC, Tait W, Campbell HA, Baldi JC. (2023) Association of preoperative high-intensity interval training with cardiorespiratory fitness and postoperative outcomes among adults undergoing major surgery: a systematic review and meta-analysis. *JAMA Netw Open* 6e2320527.
88. Snowden CP, Prentis JM, Anderson HL, Roberts DR, Randles D, Renton M *et al.* (2010) Submaximal cardiopulmonary exercise testing predicts complications and hospital length of stay in patients undergoing major elective surgery. *Ann Surg* 251:535–541.
89. Moran J, Wilson F, Guinan E, McCormick P, Hussey J, Moriarty J. (2016) Role of cardiopulmonary exercise testing as a risk-assessment method in patients undergoing intra-abdominal surgery: a systematic review. *Br J Anaesth* 116:177–191.
90. Snowden CP, Prentis J, Jacques B, Anderson H, Manas D, Jones D *et al.* (2013) Cardiorespiratory fitness predicts mortality and hospital length of stay after major elective surgery in older people. *Ann Surg* 257:999–1004.
91. Steffens D, Ismail H, Denehy L, Beckenkamp PR, Solomon M, Koh C *et al.* (2021) Preoperative cardiopulmonary exercise test associated with postoperative outcomes in patients undergoing cancer surgery: a systematic review and meta-analyses. *Ann Surg Oncol* 28: 7120–7146.
92. Bongers BC. (2023) Steep ramp test protocol for preoperative risk assessment and short-term high-intensity interval training to evaluate, improve, and monitor cardiorespiratory fitness in surgical oncology. *J Surg Oncol* 127:891–895.
93. Tew GA, Ayyash R, Durrand J, Danjoux GR. (2018) Clinical guideline and recommendations on pre-operative exercise training in patients awaiting major non-cardiac surgery. *Anaesthesia* 73:750–768.
94. Butson G, Edbrooke L, Ismail H, Denehy L. (2024) CPPOC CIPaPoC. Exercise capacity prior to major cancer surgery: a cross-sectional observational study of the validity of the 6-minute walk and 30-second sit-to-stand tests. *Asia Pac J Clin Oncol* 20:497–506.
95. Dutton J, Zardab M, De Braal VJF, Hariharan D, MacDonald N, Hallworth S *et al.* (2021) The accuracy of pre-operative (P)-POSSUM scoring and cardiopulmonary exercise testing in predicting morbidity and mortality after pancreatic and liver surgery: a systematic review. *Ann Med Surg* 62:1–9.
96. Kumar R, Garcea G. (2018) Cardiopulmonary exercise testing in hepato-biliary & pancreas cancer surgery - a systematic review: are we any further than walking up a flight of stairs? *Int J Surg* 52: 201–207.
97. Trul-Kreuz IA, Akkerman M, Kieboom EAM, Nieuwenhuis MK, Houdijk H, Bongers BC. (2024) Clinimetric properties of the steep ramp test to assess cardiorespiratory fitness, its underlying physiological responses, and its current applications: a scoping review. *Arch Phys Med Rehabil* 105:2198–2213.
98. Struthers R, Erasmus P, Holmes K, Warman P, Collingwood A, Sneyd JR. (2008) Assessing fitness for surgery: a comparison of questionnaire, incremental shuttle walk, and cardiopulmonary exercise testing in general surgical patients. *Br J Anaesth* 101:774–780.
99. Gao B, Luo J, Liu Y, Zhong F, Yang X, Gan Y *et al.* (2020) Clinical efficacy of perioperative immunonutrition containing Omega-3-Fatty acids in patients undergoing hepatectomy: a systematic review and meta-analysis of randomized controlled trials. *Ann Nutr Metab* 76: 375–386.
100. Hachiya H, Aoki T, Iso Y, Shimizu T, Tago K, Park KH *et al.* (2020) Effects of branched-chain amino acids on postoperative tumor recurrence in patients undergoing curative resection for hepatocellular carcinoma: a randomized clinical trial. *J Hepatobiliary Pancreat Sci* 27:819–829.
101. Zhang B, Wei G, Li R, Wang Y, Yu J, Wang R *et al.* (2017) n-3 fatty acid-based parenteral nutrition improves postoperative recovery for cirrhotic patients with liver cancer: a randomized controlled clinical trial. *Clin Nutr* 36:1239–1244.
102. Wong CS, Praseedom R, Liau SS. (2020) Perioperative immunonutrition in hepatectomy: a systematic review and meta-analysis. *Ann Hepatobiliary Pancreat Surg* 24:396–414.
103. Shimada M, Morine Y, Nagano H, Hatano E, Kaiho T, Miyazaki M *et al.* (2015) Effect of TU-100, a traditional Japanese medicine, administered after hepatic resection in patients with liver cancer: a multi-center, phase III trial (JFMC40-1001). *Int J Clin Oncol* 20:95–104.
104. Nishi M, Shimada M, Uchiyama H, Ikegami T, Arakawa Y, Hanaoka J *et al.* (2012) The beneficial effects of Kampo medicine dai-ken-chu-to after hepatic resection: a prospective randomized control study. *Hepatogastroenterology* 59:2290–2294.
105. You XM, Mo XS, Ma L, Zhong JH, Qin HG, Lu Z *et al.* (2015) Randomized clinical trial comparing efficacy of simo decoction and acupuncture or chewing gum alone on postoperative ileus in patients with hepatocellular carcinoma after hepatectomy. *Medicine (Baltim)* 94e1968.
106. Hendry PO, van Dam RM, Bukkems SF, McKeown DW, Parks RW, Preston T *et al.* (2010) Randomized clinical trial of laxatives and oral nutritional supplements within an enhanced recovery after surgery protocol following liver resection. *Br J Surg* 97:1198–1206.
107. Jang SY, Ju EY, Kim DE, Kim JH, Kim YH, Son M *et al.* (2012) First flatus time and xerostomia associated with gum-chewing after liver resection. *J Clin Nurs* 21:2188–2192.
108. Lassen K, Kjaeve J, Fetveit T, Tranø G, Sigurdsson HK, Horn A *et al.* (2008) Allowing normal food at will after major upper gastrointestinal surgery does not increase morbidity: a randomized multicenter trial. *Ann Surg* 247:721–729.
109. Weimann A, Braga M, Carli F, Higashiguchi T, Hübner M, Klek S *et al.* (2017) ESPEN guideline: clinical nutrition in surgery. *Clin Nutr* 36: 623–650.
110. Gao X, Zhang Y, Qi X, Xiao Y, Gao T, Jin G *et al.* (2024) Early enteral nutrition versus early supplemental parenteral nutrition in patients undergoing major abdominal surgery: a secondary analysis of 2 randomized clinical trials. *Am J Clin Nutr* 119:1036–1043.
111. Gao LB, Tian H, Wang XG, Yu XF, Guan Y, Chen ML *et al.* (2015) Early enteral and parenteral nutritional support after hepatectomy in patients with hepatic carcinoma: a systematic review and meta-analysis. *OncoTargets Ther* 8:623–631.
112. Zhao XF, Wu N, Zhao GQ, Liu JF, Dai YF. (2016) Enteral nutrition versus parenteral nutrition after major abdominal surgery in patients

- with gastrointestinal cancer: a systematic review and meta-analysis. *J Invest Med* 64:1061–1074.
113. Ni CY, Wang ZH, Huang ZP, Zhou H, Fu LJ, Cai H *et al.* (2018) Early enforced mobilization after liver resection: a prospective randomized controlled trial. *Int J Surg* 54(Pt A):254–258.
 114. de Almeida EPM, de Almeida JP, Landoni G, Galas F, Fukushima JT, Fominskiy E *et al.* (2017) Early mobilization programme improves functional capacity after major abdominal cancer surgery: a randomized controlled trial. *Br J Anaesth* 119:900–907.
 115. Tang JH, Wang B, Chow JLL, Joseph PM, Chan JY, Abdul Rahman N *et al.* (2021) Improving postoperative mobilisation rates in patients undergoing elective major hepatopancreatobiliary surgery. *Postgrad Med J* 97:239–247.
 116. Chan KS, Wang B, Tan YP, Chow JLL, Ong EL, Junnarkar SP *et al.* (2021) Single-institution, postoperative mobilization clinical practice improvement program following hepatopancreatobiliary surgery during the COVID-19 pandemic: prospective cohort study. *JMIR Perioper Med* 4e30473.
 117. Hogenbirk RNM, Wijma AG, van der Plas WY, Hentzen J, de Bock GH, van der Schans CP *et al.* (2024) Actual physical activity after major abdominal cancer surgery: far from optimal. *Eur J Surg Oncol* 50107949.
 118. Barkley R, Khalil M, Shen P, Levine EA, Votanopoulos K, Clark CJ. (2020) Feasibility of low-cost accelerometers in measuring functional recovery after major oncologic surgery. *J Surg Oncol* 121:279–285.
 119. Vaska AI, Abbas S. (2019) The role of bile leak testing in liver resection: a systematic review and meta-analysis. *HPB* 21:148–156.
 120. Wang HQ, Yang J, Yang JY, Yan LN. (2013) Bile leakage test in liver resection: a systematic review and meta-analysis. *World J Gastroenterol* 19:8420–8426.
 121. Linke R, Ulrich F, Bechstein WO, Schnitzbauer AA. (2015) The White-test helps to reduce biliary leakage in liver resection: a systematic review and meta-analysis. *Ann Hepatol* 14:161–167.
 122. Liu M, Liu Z, Jin H, Jiao Y. (2015) The efficacy of fat emulsion and normal saline for bile leakage tests during hepatic resection: a randomized trial. *J Int Med Res* 43:378–384.
 123. Liu Z, Jin H, Li Y, Gu Y, Zhai C. (2012) Randomized controlled trial of the intraoperative bile leakage test in preventing bile leakage after hepatic resection. *Dig Surg* 29:510–515.
 124. Cristaudi A, Tarantino I, Scheiwiller A, Wiencierz A, Majno-Hurst P, Schmied BM *et al.* (2021) Diagnostic role of the 'white test' with lipidic solution in the early intraoperative identification of open bile ducts for the prevention of bile leakage after liver resection: study protocol for a randomised controlled multicentric superiority trial (BiLe-Trial). *BMJ Open* 11e049449.
 125. Hanaki T, Goto K, Morimoto M, Murakami Y, Matsunaga T, Yamamoto M *et al.* (2022) Surgical administration of indocyanine green in hepatectomy for improved bile leakage detection. *Anticancer Res* 42:4787–4793.
 126. Hanaki T, Goto K, Tokuyasu N, Endo Y, Sunada H, Noma H *et al.* (2023) Efficacy of indocyanine green systemic administration for bile leak detection after hepatectomy: a protocol for a prospective single-arm clinical trial with a historical control group. *BMJ Open* 13e068223.
 127. Hajibandeh S, Raza SS, Bartlett D, Dasari BVM, Sutcliffe RP. (2023) Abdominal drainage is contraindicated after uncomplicated hepatectomy: results of a meta-analysis of randomized controlled trials. *Surgery* 173:401–411.
 128. Gurusamy KS, Samraj K, Davidson BR. (2007) Routine abdominal drainage for uncomplicated liver resection. *Cochrane Database Syst Rev* CD006232.
 129. Sun HC, Qin LX, Lu L, Wang L, Ye QH, Ren N *et al.* (2006) Randomized clinical trial of the effects of abdominal drainage after elective hepatectomy using the crushing clamp method. *Br J Surg* 93:422–426.
 130. Liu CL, Fan ST, Lo CM, Wong Y, Ng IO, Lam CM *et al.* (2004) Abdominal drainage after hepatic resection is contraindicated in patients with chronic liver diseases. *Ann Surg* 239:194–201.
 131. Dezfouli SA, Ünal UK, Ghamarnejad O, Khajeh E, Ali-Hasan-Al-Saegh S, Ramouz A *et al.* (2021) Systematic review and meta-analysis of the efficacy of prophylactic abdominal drainage in major liver resections. *Sci Rep* 11:3095.
 132. Brooke-Smith M, Figueras J, Ullah S, Rees M, Vauthey JN, Hugh TJ *et al.* (2015) Prospective evaluation of the international study group for liver surgery definition of bile leak after a liver resection and the role of routine operative drainage: an international multicentre study. *HPB* 17:46–51.
 133. Gavrilidis P, Hidalgo E, de'Angelis N, Lodge P, Azoulay D. (2017) Re-appraisal of prophylactic drainage in uncomplicated liver resections: a systematic review and meta-analysis. *HPB* 19:16–20.
 134. Anweier N, Apaer S, Zeng Q, Wu J, Gu S, Li T *et al.* (2021) Is routine abdominal drainage necessary for patients undergoing elective hepatectomy? A protocol for systematic review and meta-analysis. *Medicine (Baltim)* 100e24689.
 135. Braunwarth E, Ratti F, Aldrighetti L, Al-Saffar HA, D Souza MA, Stureson C *et al.* (2023) Incidence and risk factors for anastomotic bile leakage in hepatic resection with bilioenteric reconstruction - a international multicenter study. *HPB* 25:54–62.
 136. Kokas B, Ulmann L, Rozman P, Farkas N, Szijártó A, Szűcs Á. (2024) Postoperative bile leak after hepato-pancreato-biliary surgery in malignant biliary obstruction: rates, treatments, and outcomes in a high-volume tertiary referral center. *BMC Surg* 24:410.
 137. Fukuoka H, Toyama H, Mizumoto T, Ishida J, Asari S, Goto T *et al.* (2023) Risk factors for recurrence of cholangitis after pancreaticoduodenectomy and comparison of stents in hepaticojejunostomy. *World J Surg* 47:2499–2506.
 138. Olthoff PB, Coelen RJ, Wiggers JK, Besselink MG, Busch OR, van Gulik TM. (2016) External biliary drainage following major liver resection for perihilar cholangiocarcinoma: impact on development of liver failure and biliary leakage. *HPB* 18:348–353.
 139. Kuang Y, Ji R, Yuan T, Liu M. (2024) Modified technique of hepatojejunostomy for biliary tract reconstruction after resection of tumors affecting the perihilar region: a case series. *BMC Surg* 24:102.
 140. Alegre Castellanos A, Molina Granados JF, Escibano Fernandez J, Gallardo Muñoz I, Triviño Tarradas FeA. (2012) Early phase detection of bile leak after hepatobiliary surgery: value of Gd-EOB-DTPA-enhanced MR cholangiography. *Abdom Imaging* 37:795–802.
 141. Kul M, Erden A, Düşünceli Atman E. (2017) Diagnostic value of Gd-EOB-DTPA-enhanced MR cholangiography in non-invasive detection of postoperative bile leakage. *Br J Radiol* 9020160847.

142. Argirò R, Sensi B, Siragusa L, Bellini L, Conte LE, Riccetti C *et al.* (2023) Liver-specific contrast-enhanced magnetic resonance Cholangio-Pancreatography (Ce-MRCP) in non-invasive diagnosis of iatrogenic biliary leakage. *Diagnostics* 13.
143. Sharma P, Kumar R, Das KJ, Singh H, Pal S, Parshad R *et al.* (2012) Detection and localization of post-operative and post-traumatic bile leak: hybrid SPECT-CT with 99mTc-Mebrofenin. *Abdom Imaging* 37: 803–811.
144. Adler DG, Papachristou GI, Taylor LJ, McVay T, Birch M, Francis G *et al.* (2017) Clinical outcomes in patients with bile leaks treated via ERCP with regard to the timing of ERCP: a large multicenter study. *Gastrointest Endosc* 85:766–772.
145. Dumonceau JM, Tringali A, Papanikolaou IS, Blero D, Mangiavillano B, Schmidt A *et al.* (2018) Endoscopic biliary stenting: indications, choice of stents, and results: European society of gastrointestinal endoscopy (ESGE) clinical guideline - updated October 2017. *Endoscopy* 50:910–930.
146. Arellano RS, Reid NJ, Kapoor B, Lorenz J. (2022) The role of interventional radiology in the management of refractory bile leaks. *Abdom Radiol (NY)* 47:1881–1890.
147. Farhat S, Bourrier A, Gaudric M, Dousset B, Scatton O, Chaussade S *et al.* (2011) Endoscopic treatment of biliary fistulas after complex liver resection. *Ann Surg* 253:88–93.
148. Fukuda K, Nakai Y, Mizuno S, Sato T, Noguchi K, Kanai S *et al.* (2022) Endoscopic bridge-and-seal of bile leaks using a fully covered self-expandable metallic stent above the papilla. *J Clin Med* 11.
149. Lorenzo D, Bromberg L, Arvanitakis M, Delhay M, Fernandez Y, Viesca M *et al.* (2022) Endoscopic internal drainage of complex bilomas and biliary leaks by transmural or transpapillary/transfistulary access. *Gastrointest Endosc* 95:131–139.e136.
150. Nasr JY, Hashash JG, Orons P, Marsh W, Slivka A. (2013) Rendezvous procedure for the treatment of bile leaks and injury following segmental hepatectomy. *Dig Liver Dis* 45:433–436.
151. Murtha-Lemekhova A, Fuchs J, Feiler S, Schulz E, Teroerde M, Kalkum E *et al.* (2022) Is metabolic syndrome a risk factor in hepatectomy? A meta-analysis with subgroup analysis for histologically confirmed hepatic manifestations. *BMC Med* 20:47.
152. (EASL) EAftSoTL, (EASD) EAftSoD, (EASO) EAftSoO. (2024) EASL-EASD-EASO clinical practice guidelines on the management of metabolic dysfunction-associated steatotic liver disease (MASLD). *J Hepatol* 81):492–542.
153. Barth RJ, Mills JB, Suriawinata AA, Putra J, Tosteson TD, Axelrod D *et al.* (2019) Short-term preoperative diet decreases bleeding after partial hepatectomy: results from a multi-institutional randomized controlled trial. *Ann Surg* 269:48–52.
154. Reeves JG, Suriawinata AA, Ng DP, Holubar SD, Mills JB, Barth RJ. (2013) Short-term preoperative diet modification reduces steatosis and blood loss in patients undergoing liver resection. *Surgery* 154: 1031–1037.
155. Park L, Gilbert R, Baker L, Shorr R, Workneh A, Turcotte S *et al.* (2020) The safety and efficacy of hypovolemic phlebotomy on blood loss and transfusion in liver surgery: a systematic review and meta-analysis. *HPB* 22:340–350.
156. Martel G, Lenet T, Wherrett C, Carrier FM, Monette L, Workneh A *et al.* (2023) Phlebotomy resulting in controlled hypovolemia to prevent blood loss in major hepatic resections (PRICE-2): study protocol for a phase 3 randomized controlled trial. *Trials* 24:38.
157. Ye H, Wu H, Li B, Zuo P, Chen C. (2023) Application of cardiovascular interventions to decrease blood loss during hepatectomy: a systematic review and meta-analysis. *BMC Anesthesiol* 23:89.
158. Pathak S, Hakeem A, Pike T, Toogood GJ, Simpson M, Prasad KR *et al.* (2015) Anaesthetic and pharmacological techniques to decrease blood loss in liver surgery: a systematic review. *ANZ J Surg* 85: 923–930.
159. Karanicolas PJ, Lin Y, McCluskey SA, Tarshis J, Thorpe KE, Wei A *et al.* (2024) Team HC. Tranexamic acid in patients undergoing liver resection: the HeLiX randomized clinical trial. *JAMA* 332:1080–1089.
160. Wang HQ, Yang JY, Yan LN. (2011) Hemihepatic versus total hepatic inflow occlusion during hepatectomy: a systematic review and meta-analysis. *World J Gastroenterol* 17:3158–3164.
161. Guangtong D, Hui L, Fulong Z, Lang C, Hongru L, Dongwen W *et al.* (2019) Pringle maneuver versus hemihepatic blood flow occlusion during hepatectomy: a systematic review and meta-analysis. *Int J Clin Exp Med* 12:11–25.
162. Fagenson AM, Gleeson EM, Nabi F, Lau KN, Pitt HA. (2021) When does a Pringle Maneuver cause harm? *HPB* 23:587–594.
163. Man K, Fan ST, Ng IO, Lo CM, Liu CL, Yu WC *et al.* (1999) Tolerance of the liver to intermittent pringle maneuver in hepatectomy for liver tumors. *Arch Surg* 134:533–539.
164. Kamarajah SK, Wilson CH, Bundred JR, Lin A, Sen G, Hammond JS *et al.* (2020) A systematic review and network meta-analysis of parenchymal transection techniques during hepatectomy: an appraisal of current randomised controlled trials. *HPB* 22:204–214.
165. Brustia R, Granger B, Scatton O. (2016) An update on topical haemostatic agents in liver surgery: systematic review and meta analysis. *J Hepatobiliary Pancreat Sci* 23:609–621.
166. Wells CI, Ratnayake CBB, Mentor K, Sen G, Hammond JS, French JJ *et al.* (2020) Haemostatic efficacy of topical agents during liver resection: a network meta-analysis of randomised trials. *World J Surg* 44:3461–3469.
167. Hendi M, Lv J, Cai XJ. (2021) Current status of laparoscopic hepatectomy for the treatment of hepatocellular carcinoma: a systematic literature review. *Medicine (Baltim)* 100e27826.
168. Ding Z, Fang H, Huang M, Yu T. (2023) Laparoscopic versus open in right posterior sectionectomy: a systematic review and meta-analysis. *Langenbecks Arch Surg* 408:25.
169. Nomi T, Hirokawa F, Kaibori M, Ueno M, Tanaka S, Hokuto D *et al.* (2020) Laparoscopic versus open liver resection for hepatocellular carcinoma in elderly patients: a multi-centre propensity score-based analysis. *Surg Endosc* 34:658–666.
170. Fichtinger RS, Aldrighetti LA, Abu Hilal M, Troisi RI, Sutcliffe RP, Besselink MG *et al.*, collaborative OIP. (2024) Laparoscopic versus open hemihepatectomy: the ORANGE II PLUS multicenter randomized controlled trial. *J Clin Oncol* 42:1799–1809.
171. Fretland Å, Dagenborg VJ, Bjørnelv GMW, Kazaryan AM, Kristiansen R, Fagerland MW *et al.* (2018) Laparoscopic versus open resection for colorectal liver metastases: the OSLO-COMET randomized controlled trial. *Ann Surg* 267:199–207.
172. van Keulen AM, Buettner S, Besselink MG, Busch OR, van Gulik TM, IJzermans JNM *et al.* (2021) Primary and secondary liver failure after major liver resection for perihilar cholangiocarcinoma. *Surgery* 170: 1024–1030.
173. Sultana A, Brooke-Smith M, Ullah S, Figueras J, Rees M, Vauthey JN *et al.* (2018) Prospective evaluation of the international study group for

- liver surgery definition of post hepatectomy liver failure after liver resection: an international multicentre study. *HPB* 20:462–469.
174. Mise Y, Vauthey JN, Zimmitti G, Parker NH, Conrad C, Aloia TA *et al.* (2015) Ninety-day postoperative mortality is a legitimate measure of hepatopancreatobiliary surgical quality. *Ann Surg* 262:1071–1078.
 175. Pereyra D, Mandorfer M, Santol J, Gregory L, Koeditz C, Ortmayr G *et al.* (2024) Von Willebrand Factor Antigen Improves Risk Stratification for Patients with a Diagnosis of Resectable Hepatocellular Carcinoma. *Ann Surg Oncol* 31:6526–6536.
 176. Fukushima K, Fukumoto T, Kuramitsu K, Kido M, Takebe A, Tanaka M *et al.* (2014) Assessment of ISGLS definition of posthepatectomy liver failure and its effect on outcome in patients with hepatocellular carcinoma. *J Gastrointest Surg* 18:729–736.
 177. Baumgartner R, Gilg S, Björnsson B, Hasselgren K, Ghorbani P, Sauter C *et al.* (2022) Impact of post-hepatectomy liver failure on morbidity and short- and long-term survival after major hepatectomy. *BJS Open* 6.
 178. Sparrelid E, Gilg S, van Gulik TM. (2020) Systematic review of MARS treatment in post-hepatectomy liver failure. *HPB* 22:950–960.
 179. Nakamura N, Hatano E, Iguchi K, Seo S, Taura K, Uemoto S. (2016) Posthepatectomy liver failure affects long-term function after resection for hepatocellular carcinoma. *World J Surg* 40:929–936.
 180. Wang DD, Dong MM, Xie YM, Xu FQ, Fu TW, Wu YC *et al.* (2024) Effect of early peri-operative arterial lactate concentration level ratios on post-hepatectomy liver failure. *Discov Oncol* 15:81.
 181. Connolly C, Stattner S, Niederwieser T, Primavesi F. (2020) Systematic review on peri-operative lactate measurements to predict outcomes in patients undergoing liver resection. *J Hepatobiliary Pancreat Sci* 27:359–370.
 182. Niederwieser T, Braunwarth E, Dasari BVM, Pufal K, Szatmary P, Hackl H *et al.* (2021) Early postoperative arterial lactate concentrations to stratify risk of post-hepatectomy liver failure. *Br J Surg* 108: 1360–1370.
 183. Wang H, Li L, Bo W, Liu A, Feng X, Hu Y *et al.* (2018) Immediate postoperative Fibrosis-4 predicts postoperative liver failure for patients with hepatocellular carcinoma undergoing curative surgery. *Dig Liver Dis* 50:61–67.
 184. Wang HQ, Yang J, Yang JY, Wang WT, Yan LN. (2014) Low immediate postoperative platelet count is associated with hepatic insufficiency after hepatectomy. *World J Gastroenterol* 20:11871–11877.
 185. Haegele S, Reiter S, Wanek D, Offensperger F, Pereyra D, Stremitzer S *et al.* (2016) Perioperative non-invasive indocyanine green-clearance testing to predict postoperative outcome after liver resection. *PLoS One* 11:e0165481.
 186. Thomas MN, Weninger E, Angele M, Bösch F, Pratschke S, Andrassy J *et al.* (2015) Intraoperative simulation of remnant liver function during anatomic liver resection with indocyanine green clearance (LiMON) measurements. *HPB* 17:471–476.
 187. Squires MH, Dann GC, Lad NL, Fisher SB, Martin BM, Kooby DA *et al.* (2014) Hypophosphataemia after major hepatectomy and the risk of post-operative hepatic insufficiency and mortality: an analysis of 719 patients. *HPB* 16:884–891.
 188. Hallet J, Karanicolas PJ, Zih FS, Cheng E, Wong J, Hanna S *et al.* (2016) Hypophosphatemia and recovery of post-hepatectomy liver insufficiency. *Hepatobiliary Surg Nutr* 5:217–224.
 189. Gyoeri GP, Pereyra D, Braunwarth E, Ammann M, Jonas P, Offensperger F *et al.* (2019) The 3-60 criteria challenge established predictors of postoperative mortality and enable timely therapeutic intervention after liver resection. *Hepatobiliary Surg Nutr* 8:111–124.
 190. Truant S, Baillet C, Fulbert M, Olivier A, Sergent G, Turpin A *et al.* (2020) Asymmetric kinetics of volume and function of the remnant liver after major hepatectomy as a key for postoperative outcome - a case-matched study. *HPB* 22:855–863.
 191. Yoshida M, Shiraishi S, Sakamoto F, Beppu T, Utsunomiya D, Okabe H *et al.* (2014) Assessment of hepatic functional regeneration after hepatectomy using (99m)Tc-GSA SPECT/CT fused imaging. *Ann Nucl Med* 28:780–788.
 192. Haegele S, Offensperger F, Pereyra D, Lahner E, Assinger A, Fleischmann E *et al.* (2015) Deficiency in thrombopoietin induction after liver surgery is associated with postoperative liver dysfunction. *PLoS One* 10:e0116985.
 193. Brunthaler L, Pereyra D, Brenner M, Santol J, Herrmann L, Schrottmaier WC *et al.* (2024) Intrahepatic neutrophil accumulation and extracellular trap formation are associated with posthepatectomy liver failure. *Hepatol Commun* 8.
 194. Xu ZY, Peng M, Fan MM, Zou QF, Li YR, Jiang D. (2024) Heparin is an effective treatment for preventing liver failure after hepatectomy. *World J Gastroenterol* 30:2881–2892.
 195. Kamali C, Brunnbauer P, Kamali K, Saqr AA, Arnold A, Harman Kamali G *et al.* (2024) Extracellular NAD. *Commun Biol* 7:991.
 196. Robinson SM, Saif R, Sen G, French JJ, Jaques BC, Charnley RM *et al.* (2013) N-acetylcysteine administration does not improve patient outcome after liver resection. *HPB* 15:457–462.
 197. Grendar J, Ouellet JF, McKay A, Sutherland FR, Bathe OF, Ball CG *et al.* (2016) Effect of N-acetylcysteine on liver recovery after resection: a randomized clinical trial. *J Surg Oncol* 114:446–450.
 198. Pei Y, Chai S, Zhang Y, Zhang Z, Chen X, Zhang W. (2019) Benefits of splenectomy and curative treatments for patients with hepatocellular carcinoma and portal hypertension: a retrospective study. *J Gastrointest Surg* 23:2151–2162.
 199. Monet C, De Jong A, Aarab Y, Piron L, Prades A, Carr J *et al.* (2022) Adverse events, short- and long-term outcomes of extra corporeal liver therapy in the intensive care unit: 16 years experience with MARS® in a single center. *Crit Care* 26:282.
 200. Gilg S, Sparrelid E, Engstrand J, Baumgartner R, Nowak G, Stål P *et al.* (2022) Molecular adsorbent recirculating system treatment in patients with post-hepatectomy liver failure: long-term results of a pilot study. *Scand J Surg* 111:48–55.
 201. Gilg S, Sparrelid E, Saraste L, Nowak G, Wahlin S, Strömberg C *et al.* (2018) The molecular adsorbent recirculating system in post-hepatectomy liver failure: results from a prospective phase I study. *Hepatol Commun* 2:445–454.
 202. Sparrelid E, Thorsen T, Sauter C, Jorns C, Stål P, Nordin A *et al.* (2022) Liver transplantation in patients with post-hepatectomy liver failure - a Northern European multicenter cohort study. *HPB* 24: 1138–1144.
 203. Thorsen T, Solheim JM, Labori KJ, Line PD, Aandahl EM. (2019) Liver transplantation as a lifesaving procedure for posthepatectomy liver failure and iatrogenic liver injuries. *Langenbecks Arch Surg* 404: 301–308.
 204. Azoulay D, Desterke C, Bhangu P, Serrablo A, De Martin E, Cauchy F *et al.* (2024) Rescue liver transplantation for posthepatectomy liver failure: a systematic review and survey of an international experience. *Transplantation* 108:947–957.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.hpb.2025.10.008>.

Topic 1: Perioperative thromboprophylaxis

Subtopic	Recommendation	Recommendation grade	Median level of evidence
Q1: When should perioperative thromboprophylaxis be commenced in liver surgery?	Pharmacological thromboprophylaxis is recommended for all patients undergoing liver surgery and can be initiated either pre- or postoperatively. Mechanical thromboprophylaxis (socks, pneumatic) should be considered routinely.	Conditional	2+
Q2: What is the optimal duration of perioperative thromboprophylaxis in liver surgery?	A 4-week period of administration of LMWH should be considered after major liver surgery. However, a prolonged (>4 weeks) prophylaxis may be indicated in high-risk patients based on individual risk stratification.	Conditional	2-
Q3: What anticoagulation should be used when performing hepatectomy with portal vein resection and reconstruction?	There is no strong evidence for specific perioperative prevention strategy of portal vein thrombosis in patients undergoing liver surgery with portal vein resection and reconstruction. The approach should be tailored according to the patient-specific factors and clinical judgement.	Conditional	2-
Q4: Is there a role for defining early and late portal vein thrombosis (PVT) after liver resection?	Defining early and late PVT after liver resection should be considered with the aim to standardize outcomes. A widely accepted definition specific to liver resection is lacking. A 30-day cut-off, as used in liver transplantation, is a pragmatic approach that may be adopted, with limitations related to partially different pathophysiological causes of PVT after liver resection compared to liver transplantation.	Conditional	4
Q5: Are there differences in the treatment of early and late portal vein thrombosis after liver resection?	There are no high-quality comparative studies to establish standardized treatment algorithms for early and late PVT after liver resection. Treatment options include anticoagulation, surgical intervention, and interventional radiology procedures. The decision on treatment strategy should be adjusted according to the time onset, thrombus extent and site, patient-specific factors, and institutional resources.	Conditional	4

Topic 2: Perioperative Antibiotics

Subtopic	Recommendation	Recommendation grade	Median level of evidence
Q1: Which antibiotics should be used as standard prophylaxis for liver surgery?	For liver surgery prophylaxis, antibiotic choice should align with local pathogen resistance patterns. First-generation cephalosporins such as cefazolin are preferred for livers with naive bile ducts, administered within 60 min before incision. Broader-spectrum agents, like ceftriaxone or Piperacillin-Tazobactam, are conditionally recommended for high-risk cases. Consultation with a microbiologist should be considered in MRSA-risk settings. In beta-lactamic allergic patients Fosfomycin or aminoglycosides should be selected.	Conditional	2++
Q2: Which patients should be defined as high-risk for developing clinically relevant SSI following hepatectomy?	Patients at higher risk of SSI can be identified by bacterial colonization resistant to commonly used prophylactic antibiotics, as well as by factors which make them generally more at risk for postoperative complications like underlying liver disease, preoperative biliary drainage, and synchronous resections of the gastrointestinal tract. In patients with preoperative biliary drainage cultures should be obtained at the time of the intervention. In case of bile duct resection and reconstruction intraoperative biliary cultures should be obtained during the surgical procedure.	Conditional	2+
Q3: Which patients are likely to benefit from extended antibiotic prophylaxis?	Although antibiotic prophylaxis is recommended, extended postoperative antibiotic prophylaxis for more than 24 h has not demonstrated any additional benefit in preventing postoperative infections after standard hepatectomies. Patients at high risk for developing clinically relevant SSI may benefit from extended antimicrobial duration (three to five days postoperatively).	Conditional	2+
Q4: What is the role of antibiotic treatment in cases of postoperative fever of unknown origin and no suspicion of infection or sepsis following liver resection?	Antibiotic therapy is not routinely recommended for postoperative fever of unknown origin if diagnostic workup excluded any signs of infection or sepsis. High-risk patients — such as those with preoperative biliary drainages, bilio-enteric anastomosis, immunosuppression, or comorbidities like diabetes mellitus — may benefit from targeted antibiotic therapy if early signs of infection are present.	Strong	1+

Topic 3: Prehabilitation, Nutrition, Mobilisation

Subtopic	Recommendation	Recommendation grade	Median level of evidence
Q1: What is the value of a prehabilitation program in patients undergoing hepatectomy?	Multimodal prehabilitation programs (exercise training, nutritional optimization, psychological interventions) should be considered for patients prior to hepatectomy for both benign and malignant liver disease following assessment to reduce postoperative morbidity and length of hospital stay.	Strong	1+
Q2: What is the influence of preoperative body composition for outcomes in patients undergoing hepatectomy?	Preoperative body composition influences outcomes in patients undergoing hepatectomy for various conditions of liver malignancies. Therefore, preoperative screening for low skeletal muscle mass (LSMM) should be done in every patient considered for hepatectomy for liver malignancies.	Strong	1+
Q3: What is the role of fitness screening and assessment in patients undergoing hepatectomy?	As aerobic fitness is an important modifiable risk factor and can be improved by a prehabilitation program, screening for this risk factor can be incorporated as a routine part of the preoperative assessment for patients undergoing liver surgery, particularly for those with borderline or reduced fitness, as it provides critical information for optimizing perioperative care and improving postoperative outcomes.	Conditional	1
Q4: Does perioperative immunonutrition offer a benefit in patients undergoing hepatectomy?	Immunonutrition can be administered as perioperative nutritional supplementation in patients undergoing hepatectomy as it has a positive effect on postoperative liver function and might foster the regenerative potential especially in vulnerable populations.	Conditional	1-
Q5: What is the role of prokinetic policies to improve time to first bowel movement/full standard diet?	Routine use of postoperative prokinetics or laxatives is recommended to stimulate bowel movement after hepatectomy, while prokinetics may shorten the time to first flatus or stool. In patients who have had a simultaneous liver and colonic resection, use of prokinetic treatment should be used with caution and based upon established colorectal protocols.	Conditional	1+
Q6: When and how should artificial nutrition be added to oral nutrition in the postoperative period?	Early oral nutrition should be preferred after hepatectomy. In patients requiring artificial nutrition support (malnourished patients, patients with complications causing several days of fasting), enteral nutrition through a nasofeeding tube should be preferred over total parenteral nutrition (TPN). Nutritional specialist assessment is recommended for patients with delayed oral intake.	Conditional	1-
Q7: How should (early) postoperative mobilization after hepatectomy be defined?	The ideal prescription on postoperative mobilization after hepatic surgery is not clear. Early mobilization after liver surgery should start immediately on postoperative day one with assisted active mobilization and sitting out of bed. Walking (with no precise target) should start at least on second postoperative day adapted to the patient's functional capacity and carefully supervised especially in elderly or frail patients.	Strong	1+

Topic 4: Post-hepatectomy bile leak including bilio-enteric anastomosis leaks

Subtopic	Recommendation	Recommendation grade	Median level of evidence
Q1: Is there a role for intraoperative bile leak test (i.e. white test, methylene blue, indocyanine green (ICG) ...) after hepatectomy without bilio-enteric anastomosis?	Intraoperative bile leak tests can be considered after liver resection (conditional), especially in major resections or complex procedures (strong), recording its use in the operation report. No specific test has shown clear superiority regarding identification of leaks, although the gauze (surface) test and the “white-test” might be preferred. Therefore, the choice of test should depend on centre expertise and availability.	Conditional (major/complex resections: Strong)	1-
Q2: Is there a role for drainage after hepatectomy without bilio-enteric reconstruction?	Routine abdominal drainage after an uncomplicated hepatectomy should be avoided.	Strong	1+
Q3: What is the value of biliary trans-anastomotic stents in complex bilio-enteric anastomoses in hepatectomy?	Trans-anastomotic stents can be considered in complex bilio-enteric anastomoses (e.g., extended resections, small bile ducts, friable tissue) but are not recommended for routine use. If used, they should be removed within 3 months to minimize complications. The choice of trans-anastomotic stent depends on centre availability and expertise.	Conditional	2-
Q4: Is there a role for routine imaging (computed tomography (CT), magnetic resonance (MR) cholangiography) in high-risk patients for bile leaks?	Routine use of advanced imaging modalities such as CT, MR, or single-photon emission computed tomography (SPECT) is not recommended for asymptomatic high-risk patients due to their cost and limited availability in most healthcare settings. For patients with suspected or detected bile leaks, the threshold for performing radiological imaging should be low. The choice of modality should be guided by the resources available at the treating centre.	Conditional	3
Q5: What is the role of endoscopy and interventional radiology in patients with clinically relevant bile leakage after hepatectomy?	In case of a clinically relevant bile leakage (>100 ml bile/24 h) when a drain is in place, an early (endoscopic retrograde cholangiopancreatography) ERCP and stent placement seems sensible as a strategy to resolve bile leakage quickly. When the patient has a bilioma with no drain in place, the first step should be percutaneous drainage; ERCP/stent should follow on the same conditions as above.	Conditional	2-

Topic 5: Post-hepatectomy haemorrhage (PHH)

Subtopic	Recommendation	Recommendation grade	Median level of evidence
Q1: What is the role of pre-operative strategies to reduce blood loss associated with liver resection (fat-reducing diet, nutritional changes, hypovolaemic phlebotomy)?	In patients who are overweight or obese, the implementation of a low-calorie, low-fat diet should be considered, provided that the surgical timeline permits adequate intervention. Pre-operative management of metabolic dysfunction-associated steatotic liver disease (MASLD) should align with the recommendations of the EASL–EASD–EASO clinical practice guidelines. In selected patients at high risk of significant intra-operative bleeding, pre-operative hypovolaemic phlebotomy may be considered as a strategy to reduce transfusion requirements.	Strong	2-
Q2: What is the role of anaesthetic management to reduce blood loss in patients undergoing hepatectomy?	Intraoperative low central venous pressure (CVP) should be maintained until completion of parenchymal transection to reduce blood loss and transfusion requirements. No specific anaesthetic agent is recommended. Tranexamic acid is not advised in liver resection for cancer as it does not decrease blood loss during liver resection. The role of acute normovolaemic haemodilution remains unclear.	Moderate to strong	2++
Q3: What is the role for intra-operative liver inflow occlusion in reducing intra-operative blood loss?	Inflow occlusion techniques such as the Pringle maneuver (PM) and hemihepatic occlusion (HHO) can reduce intraoperative blood loss and should be considered when major bleeding is anticipated. While intermittent Pringle occlusion is safe for up to 120 min, prolonged clamping may increase the risk of ischemia-reperfusion injury, particularly in patients with underlying liver dysfunction.	Moderate	2+
Q4: Is there an influence of parenchymal transection technique on blood loss?	Meta-analyses of RCTs suggest that bipolar energy devices may reduce intraoperative blood loss during parenchymal transection. However, due to the wide range of available techniques, the limited number of direct comparisons, and methodological variability across studies, no specific recommendation can currently be made regarding the optimal transection technique.	Moderate	1+
Q5: What is the value of haemostatic agents to reduce blood loss in patients undergoing hepatectomy?	No specific recommendations can be made as to the use of topical haemostats. The time to haemostasis is shorter when using topical haemostatic agents (fibrin sealant) but there is no significant difference in peri operative blood transfusion. There is no evidence of one individual topical sealant being superior to another.	Moderate	1+
Q6: Does minimally-invasive surgery influence blood loss in patients undergoing hepatectomy?	Minimally-invasive surgery (MIS) does not reduce blood loss compared to open liver surgery for minor nor for major hepatectomy. MIS liver surgery may be associated with less blood loss in some conditions (i.e. HCC) based on non-randomized observations.	Strong (minor/major hepatectomy), conditional (HCC)	1-

Topic 6: Post-hepatectomy liver failure (PHLF)

Subtopic	Recommendation	Recommendation grade	Median level of evidence
Q1: Is there a clinical relevance to subclassify PHLF by the main underlying pathomechanism?	PHLF involves complex and heterogenous pathophysiological mechanisms and classification as primary and secondary PHLF can be considered, because it impacts on preventive and therapeutic interventions and prognosis.	Conditional	2+
Q2: Should patients suffering from PHLF be monitored beyond 90 days after hepatectomy?	Primary and secondary PHLF usually occur within 90 days after surgery. Based on indirect evidence, PHLF monitoring beyond 90 days can be considered for patients who have recovered from PHLF independent of oncological follow-up.	Conditional	4
Q3: Which postoperative markers should be considered to monitor postoperative liver function recovery in patients at high risk of PHLF?	The following parameters should be used to monitor liver function in patients at high risk of PHLF: Bilirubin, INR, albumin, lactate, ammonia levels (in patients with signs of hepatic encephalopathy), AST, alanine aminotransferase (ALT), glucose and temperature. Furthermore, infection monitoring by complete blood count (CBC) and CRP should be performed, as well as kidney function monitoring using creatinine.	Strong	2+
Q4: Is there a value for early postoperative markers to predict PHLF?	Several biomarkers (e.g. lactate, phosphate, platelets) have been reported to allow for early postoperative identification of PHLF and their measurement can be considered in high-risk patients after hepatic resection to initiate supportive treatment early. There are no comparative studies, and future research is needed to identify the optimal biomarker.	Conditional	2+
Q5: Which treatments are available to support hepatic regeneration in PHLF?	Currently, no specific therapeutic interventions are available or approved to promote liver regeneration and avoid or treat progression of PHLF.	Strong	2+