

Updated clinical guideline for human papillomavirus vaccine: the Korean Society of Gynecologic Oncology guidelines.

Title: Updated clinical guideline for human papillomavirus vaccine: the Korean Society of Gynecologic Oncology guidelines.

Publication year: 2021

Journal: Journal of gynecologic oncology

Authors: Department of Obstetrics and Gynecology, Korea University Guro Hospital, Korea University College of Medicine, Seoul, Korea.

Society: American College of Cardiology

Origin country: United States

Target countries: United States

DOI: 10.3802/jgo.2021.32.e94

PMID: 34708596

Source: <https://pubmed.ncbi.nlm.nih.gov/34708596/>

pmc J Gynecol Oncol J Gynecol Oncol 883 jgo 101483150 JGO Journal of Gynecologic Oncology 2005-0380 2005-0399 Korean Society of Gynecologic Oncology and Colposcopy PMC8550930 PMC8550930.1 8550930 8550930 34708596 10.3802/jgo.2021.32.e94 1 Practice Guideline Updated clinical guideline for human papillomavirus vaccine: the Korean Society of Gynecologic Oncology guidelines <https://orcid.org/0000-0002-0724-9971> Cho Hyun-Woong 1 * <https://orcid.org/0000-0002-5783-4968> Min Kyung-Jin 2 * <https://orcid.org/0000-0002-9121-3954> Kwon Sang-Hoon 3 <https://orcid.org/0000-0001-9254-6024> Kim Kidong 4 <https://orcid.org/0000-0002-1645-7473> Kim Sunghoon 5 <https://orcid.org/0000-0003-3820-3412> Seong Seok Ju 6 <https://orcid.org/0000-0002-6103-2466> Song Yong Jung 7 <https://orcid.org/0000-0001-9005-7796> Lee Keun Ho 8 <https://orcid.org/0000-0002-5088-1905> Lee Shin-Wha 9 <https://orcid.org/0000-0002-6945-0398> Lee Jeong-Won 10 <https://orcid.org/0000-0002-0013-8304> Ju Woong 11 <https://orcid.org/0000-0002-7347-1052> Kim Young Tae 5 <https://orcid.org/0000-0003-3101-6403> Lee Jae Kwan 1 1 Department of Obstetrics and Gynecology, Korea University Guro Hospital, Korea University College of Medicine, Seoul, Korea . 2 Department of Obstetrics and Gynecology, Korea University Ansan Hospital, Korea University College of Medicine, Ansan, Korea . 3 Department of Obstetrics and Gynecology, Keimyung University School of Medicine, Daegu, Korea . 4 Department of Obstetrics and Gynecology, Seoul National University Bundang Hospital, Seongnam, Korea . 5 Department of Obstetrics and Gynecology, Yonsei University College of Medicine, Seoul, Korea . 6 Department of Obstetrics and Gynecology, CHA Gangnam Medical Center, CHA University, Seoul, Korea . 7 Department of Obstetrics and Gynecology, Pusan National University School of Medicine, Yangsan, Korea . 8 Department of Obstetrics and Gynecology, Seoul St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Seoul, Korea . 9 Department of Obstetrics and Gynecology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea . 10 Department of Obstetrics and Gynecology, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea . 11 Department of Obstetrics and Gynecology, Ewha Womans University Mokdong Hospital, Ewha Womans University School of Medicine, Seoul, Korea . Correspondence to Jae Kwan Lee. Department of Obstetrics and Gynecology, Korea University Guro Hospital, Korea University College of Medicine, 148 Gurodong-ro, Guro-gu, Seoul 08308, Korea. jkleee38@korea.ac.kr * Hyun-Woong Cho and Kyung-Jin Min contributed equally to this work.

11 2021 17 9 2021 32 6 392658 e94 02 7 2021 02 8 2021 07 9 2021 01 11 2021 09 11 2021 09 11 2021 Copyright © 2021. Asian Society of Gynecologic Oncology, Korean Society of Gynecologic Oncology, and Japan Society of Gynecologic Oncology 2021 Asian Society of Gynecologic Oncology, Korean Society of Gynecologic Oncology, and Japan Society of Gynecologic Oncology <https://creativecommons.org/licenses/by-nc/4.0/> This is an Open Access article distributed under

the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. Since the human papillomavirus (HPV) vaccine guidelines were developed by the Korean Society of Gynecologic Oncology (KSGO) in 2011, 2016, and 2019, several recent studies on the efficacy and safety of HPV vaccines in middle-aged women and men have been reported. Furthermore, there has been an ongoing debate regarding the efficacy of the HPV vaccine in women with prior HPV infection or who have undergone conization for cervical intraepithelial neoplasia (CIN). We searched and reviewed studies on the efficacy and safety of the HPV vaccine in middle-aged women and men and the efficacy of the HPV vaccine in patients infected with HPV and those who underwent conization for CIN. The KSGO updated their guidelines based on the results of the studies included in this review.

1. Key question 1: what is the recommended age for the 3-dose schedule of the HPV vaccine?

- The optimal age for the 3-dose schedule of the quadrivalent HPV vaccine is 9–45 years in girls and women and 9–26 years in boys and men (IA).
- The optimal age for the three-dose schedule of the bivalent HPV vaccine is 9–45 years in girls and women (IA) and 9–25 years in boys and men (IIC).
- The optimal age for the three-dose schedule of the nonavalent HPV vaccine is 9–45 years in girls and women and 9–26 years in boys and men (girls and women aged 9–26 years, IA; women aged 27–45 years, IIB; and boys and men aged 9–26 years, IA).

2. Key question 2: is the HPV vaccine effective in women already infected with HPV?

- In HPV-infected women already, the administration of the quadrivalent HPV vaccine can lower the risk of infection with an uninfected HPV subtype (IIB).
- In HPV-infected women already, the administration of the bivalent HPV vaccine can lower the risk of infection with an uninfected HPV subtype (IIB).
- In HPV-infected women already, the administration of the nonavalent HPV vaccine can lower the risk of infection with an uninfected HPV subtype (IIB).

3. Key question 3: is the HPV vaccine effective in preventing the recurrence of HPV-related diseases in patients who underwent conization?

- In women who underwent conization, the administration of the quadrivalent HPV vaccine can reduce the risk of recurrence of HPV-related diseases by preventing infection with the corresponding viral subtype (IIB).
- In women who underwent conization, the administration of the bivalent HPV vaccine can reduce the risk of recurrence of HPV-related diseases by preventing infection with the corresponding viral subtype (IIB).
- In women who underwent conization, the administration of the nonavalent HPV vaccine can reduce the risk of recurrence of HPV-related diseases by preventing infection with the corresponding viral subtype (IVB).

High-risk human papillomavirus (HPV) infection is associated with the incidence of cervical, vaginal, vulvar, penile, anal, and oropharyngeal cancers and is responsible for 5%–10% of all cancers [1]. In particular, cervical cancer accounts for 80% of HPV-related cancers [2]. HPV is the most

widespread sexually transmitted infection, with a prevalence of approximately 11%, and persistent HPV infection is a major cause of cervical precancerous lesions and cancer [3 4]. Prophylactic vaccination against high-risk HPV can prevent precancerous lesions of cervical cancer and cervical cancer, as well as cancers that occur in men, such as oropharyngeal cancer, anal cancer, and penile cancer [5 6 7 8].

To date, three HPV vaccines have been approved; they can prevent infection with HPV types 16 and 18, which cause approximately 70% of cervical cancers and other HPV-related tumors [6 9 10 11 12 13 14 15 16]. The quadrivalent vaccine can further prevent infection with HPV types 6 and 11, which account for 90% of genital warts [17]. The most recently introduced 9-valent vaccine can protect against infection with 5 additional types of high-risk HPV (31, 33, 45, 52, and 58), in addition to these 4 types of HPV [18].

The Korean Society of Gynecologic Oncology (KSGO) provided and updated the clinical recommendations for HPV vaccines in 2011 and 2016 [19 20] and announced a recommendation for the 9-valent HPV vaccine in 2019 [8].

However, many studies have been recently reported on the effectiveness of HPV vaccines in middle-aged women and men. Additionally, the effectiveness of HPV vaccines in women already infected with HPV or who have undergone conization for precancerous cervical lesions is controversial. Therefore, updated recommendations for HPV vaccines are required.

The key questions and scope of guideline were derived through a meeting of the committee. The committee tried to develop a guideline by systematically searching for the latest literature related to vaccines and evaluating evidence.

According to the criteria for selection and exclusion criteria of all key questions, 2 reviewers per document independently selected and excluded the literature. The primary selection and exclusion were made by looking at the title and the abstract, and the literature selected by even 1 reviewer was searched for the original article. After that, the committee members selected and excluded based on the original article, and if there was no agreement between the members, the final selection and exclusion was decided through discussion. Articles were searched on Ovid-MEDLINE, Ovid-Embase, and Cochrane Library.

In the selected literature, randomized controlled clinical researches were evaluated using risk of bias (ROB) of Cochrane, and non-randomized observational studies were assessed by NEWCASTLE-OTTAWA QUALITY ASSESSMENT SCALE COHORT STUDIES. The quality of the literature is assessed independently by 2 members, and it was determined by an agreement between the 2 members if the quality of the literature is different.

Data extraction from the final selected literature was carried out independently by 2 working-level members according to the predetermined data extraction form. If data extracted were different between the 2 members, final settlement was made by discussion. A summary of evidence is attached to the Data S2 . Classification of evidence and recommendations are displayed in Tables 1 and 2 .

The process of developing clinical guidelines and an evidence table for each key question is addressed in the Supplementary Materials (Data S1 and S2).

In 2016, the KSGO recommended 9–26 years of age as the appropriate age for receiving the three doses of the 4-valent vaccine and 9–25 years of age as the appropriate age for receiving the three doses of the 2-valent vaccine in girls and women [21]. Additionally, in 2019, girls and women aged 9–26 years were recommended as appropriate for receiving the three doses of the 9-valent vaccine [8]. In this guideline, the effectiveness and safety of the three doses of the HPV vaccine in middle-aged women and men have been investigated.

To date, 2 randomized controlled trials (RCTs) for women over 20 years of age have been published, and 5 papers have been identified. [22 23 24 25 26]. In a multinational clinical trial published in 2009 (Protocol 019, NCT00090220), an interim analysis was performed in women aged 24–45 years over an average follow-up period of 2.2 years. The vaccine's efficacies against infection or disease associated with 4 types of HPV (6, 11, 16, and 18) and 2 types of HPV (16 and 18) were 90.5% (95% confidence interval [CI]=73.7, 97.5) and 83.1% (95% CI=50.6, 95.8), respectively [22]. After 4 years of follow-up, the efficacies against diseases related to the 4 types of HPV were 88.7% (95% CI=78.1, 94.8), and 66.9% (95% CI=4.3, 90.6) in women who were not infected with the 4 types of HPV and those who were vaccinated at least once, respectively [23]. The seropositivity rates for antibodies against HPV types 6, 11, 16, and 18 at 48 months after the first vaccination were 91.5%, 92.0%, 97.4%, and 47.9%, respectively, and no serious side effects were reported [22 23].

In the study V501-041 (NCT00834106), the vaccine's efficacy against cervical intraepithelial neoplasia (CIN) ≥ 2 related to HPV types 16 and 18 over a 78-month follow-up period in Chinese women aged 20–45 years was 100% (95% CI=32.3, 100) and that against CIN grade 1+ related to HPV types 6, 11, 16 and 18 was also 100% (95% CI=70.9, 100). Additionally, its efficacy against persistent infection with HPV types 16 and 18 was 97.5% (95% CI=85.1, 99.9), and the efficacy against abnormal cytology related to HPV types 6, 11, 16, and 18 was 94.0% (95% CI=81.5, 98.8) [26]. No serious adverse reactions were observed [25].

An RCT involving adolescent male aged 9–15 years and 2 RCTs involving male participants aged 16–26 years have been published. In V501-018 (NCT00092547), the seropositivity rate in adolescent boys aged 9–15 years at 10 years after vaccination was 89.96%, and the antibody titer was 16%–42% higher in boys aged 9–12 years than in boys aged 13–15 years [5]. In the Merck protocol 020 trial (NCT00090285) for boys and men aged 16–26 years, the vaccine's efficacies against anal epithelial neoplasia related to HPV types 6, 11, 16, and 18 after a follow-up period of 2.2 years were 77.5% (95% CI=39.6, 93.3) and 50.3% (95% CI=25.7, 67.2) in the per protocol and intention to treat populations, respectively. Additionally, the incidence of anal intraepithelial neoplasia 2 and 3 in men who received the vaccine decreased by 74.9% (95% CI=8.8, 95.4) and 54.2% (95% CI=18.0, 75.3) in the per protocol and intention to treat populations, respectively [27]. In phase II study for boys and men aged 16–26 years, the vaccine's efficacies against external genital warts were 90.4% (95% CI=69.2, 98.1) and 65.5% (95% CI=45.8, 78.6) in the per protocol and intention to treatment populations, respectively [6].

For the 4-valent vaccine, 2 doses were recommended from the age of 9 to 13 years, based on previous studies (level of evidence I, recommendation level B).

The above results determined the following recommendation for target of quadrivalent HPV vaccine:

- The optimal age for the 3-dose schedule of the 4-valent HPV vaccine is 9–45 years in girls and women and 9–26 years in boys and men (IA).

Three RCTs [28 29 30] and three observation studies [31 32 33] on middle-aged women were identified. A RCT (VIVIANE study, NCT00294047) in women over 25 years of age evaluated and compared the efficacy of the vaccine in the following age groups: 26–35, 36–45, and 46–55 years [30]. The efficacy of the 2-valent vaccine against 6-month-long infection or mild cervical epithelial dysplasia associated with HPV types 16 or 18 was 81.1% (97.7% CI=52.1, 94.0) in all age groups, 83.5% (97.7% CI=45.0, 96.8) in the 26–35 years age group, and 77.2% (97.7% CI=2.8, 96.9) in the 36–45 years age group. In an RCT (study HPV-010, NCT00423046) comparing the 2-valent and 4-valent vaccines in women aged 18–45 years, the seropositivity rate for HPV type 16 at 60 months after vaccination was 100% in the 2-valent vaccine group and 97.5–100% in the 4-valent vaccine group. The seropositivity rate for HPV type 18 was 99.0–100% in the 2-valent vaccine group and

72.3%–84.4% in the 4-valent vaccine group [29]. However, the antibody titer and the duration of antibody production against HPV type 16 or 18 were found to decrease in women aged 27–35 years and 36–45 years compared with that in women aged 18–26 years (HPV 16 and 18: women aged 18–26 years, 68.2 and 40.6 years, respectively; women aged 27–35 years, 57.3 and 9.5 years, respectively; and women aged 36–45 years, 31.0 and 1.9 years, respectively). No serious adverse effects were observed. In men, a phase I/II RCT was noted [7]. Seven months after vaccination in boys aged 10–18 years, seropositivity was 100%, and no serious side effects were observed. Additionally, for the 2-valent vaccine, the 2 doses were recommended at the age of 9–14 years based on previous research outcomes, including those from RCTs, and the 2016 KSGO recommendation (level of evidence I, recommendation level B).

According to the aforementioned results, the recommendation was as follows:

- The optimal age for the three-dose schedule of the 2-valent HPV vaccine is 9–45 years in girls and women (IA) and 9–25 years in boys and men (IIC).

A recently published phase III RCT (NCT03158220) that evaluated the efficacy and safety of the 9-valent vaccine was identified [34]. In this RCT, the immunogenicity and stability of the 9-valent vaccine were evaluated in women aged 27–45 years compared with those in women aged 16–26 years. At 7 months of vaccination, antibody titers against HPV types 6, 11, 16, 18, 31, 33, 45, 52, and 58 were not significantly different between women aged 27–45 years and women aged 16–26 years (0.60–0.67). The seroconversion rate was 99% or higher for all HPV types. No serious vaccine-related side effects have been reported to date. In boys and men, in the recommendations published in 2019 by the KSGO the 9-valent vaccine was recommended for boys and men aged 9–26 years (level of evidence I, recommendation level A) [8].

Based on this information, the recommendations was as follows:

- The optimal age for the three-dose schedule of the 9-valent HPV vaccine is 9–45 years in girls and women and 9–26 years in boys and men (girls and women aged 9–26 years, IA; women aged 27–45 years, IIIB; and boys and men aged 9–26 years, IA).

In a randomized pilot study, 10 women aged 27–45 years who tested positive for HPV 16 were divided into the vaccinated and non-vaccinated groups. In the vaccination group, antibody titers increased by 24–930-fold, and the number of memory B cells increased by 3–27-fold [35].

Based on the above results, the following was recommended:

- In HPV-infected women already, the administration of the quadrivalent HPV vaccine can lower the risk of infection with an uninfected HPV subtype (IIB).

A phase III RCT (PATRICIA study, NCT00122681) in girls and women aged 15–25 years was identified. According to this study, when the 2-valent HPV vaccine was administered to women infected with HPV type 16 or 18 at the beginning of the study, the efficacy against CIN grade 2+ related to an HPV type other than the one causing infection was 90% (95% CI=31.8, 99.8) [36].

Based on the above results, the recommendation was as follows:

- In HPV-infected women already, the administration of the bivalent HPV vaccine can lower the risk of infection with an uninfected HPV subtype (IIB).

A combined study of three RCTs (9-valent Vaccine Study Protocol V503-001, Quaternary Vaccine Study Protocol V501-013, and V501-015) was identified. In this study, the incidence of cervical lesion related to HPV 31, 33, 45, 52, and 58 was decreased among women who tested positive at baseline for HPV 6, 11, 16, or 18 but tested negative for HPV 31, 33, 45, 52, and 58 (all grades:

95.1%; high grades: 91.1%). Similarly, in women who tested positive at baseline for HPV 31, 33, 45, or 52 but tested negative for HPV 6, 11, 16, and 18, the incidence of cervical disease related to HPV 6, 11, 16, and 18 was also significantly reduced (all grades: 97.4%; high grade: 95.8%) [12].

Based on the above results, the recommendation was as follows:

- In HPV-infected women already, the administration of the nonavalent HPV vaccine can lower the risk of infection with an uninfected HPV subtype (IIB).

In an RCT of 312 patients who received treatment such as conization for CIN grade 1–3, the recurrence rate of CIN grade 1–3 was significantly decreased (58.7%) in the vaccinated group compared with that in the control group [37]. In a prospective case-control study of patients who underwent cone resection for CIN, the risk of high-grade cervical epithelial tumors in vaccinated patients was reduced by 81.2% (95% CI=34.3, 95.7) [38]. In a retrospective study of 20–45-year-old Korean women who underwent cone resection for high-grade CIN, the recurrence rate of high-grade CIN was decreased in the vaccinated group (vaccinated group, 2.5%; unvaccinated group, 7.2%), and the recurrence risk of high-grade CIN was significantly higher in the unvaccinated group than in the vaccinated group (hazard ratio=2.84; 95% CI=1.335, 6.042) [39]. In a single-center retrospective study of women who underwent conization for cervical dysplasia, the recurrence of cervical dysplasia was less common in the vaccinated group than in the unvaccinated group (7.1% vs. 16.5%, $p=0.02$), and vaccination after conization was a significant factor in preventing recurrence (odds ratio [OR]=0.2; 95% CI=0.1, 0.6) [40].

Based on the above, the following was recommended:

- In women who underwent conization, the administration of the quadrivalent HPV vaccine can reduce the risk of recurrence of HPV-related diseases by preventing infection with the corresponding viral subtype (IIB).

In an RCT (PATRICIA, NCT00122681) including girls and women aged 15–25 years, vaccine efficacy against the recurrence of CIN grade ≥ 2 was 88.2% (95% CI=14.8, 99.7) in women who underwent conization after vaccination [41]. However, in a analysis of RCT (Costa Rica HPV Vaccine Trial, NCT00128661), no vaccine efficacy was noted against cervical precancerous lesions in patients who underwent conization for cervical lesions after vaccination (efficacy: high-grade squamous intraepithelial lesion + –163.3% [95% CI= –742, 18]; CIN2+ –55.5% [95% CI=–834, 74]) [42]. In a meta-analysis that included the above 2 studies (2 studies on the 2-valent vaccine and three studies on the 4-valent vaccine), vaccination before and after conization could reduce the recurrence of CIN grade ≥ 2 (pretreatment vaccination: OR=0.4; 95% CI=0.21, 0.78; post-treatment vaccination: OR=0.28; 95% CI=0.14, 0.56) [43].

Based on the above results, the following recommendations were made:

- In women who underwent conization, the administration of the bivalent HPV vaccine can reduce the risk of recurrence of HPV-related diseases by preventing infection with the corresponding viral subtype (IIB).

In a retrospective study of women who were vaccinated after conization (2-valent: 19.6%; 4-valent: 4.6%; 9-valent: 64.1%), the incidence of persistent or recurrent high-grade CIN was significantly lower in the vaccinated group than in the unvaccinated group (3.3% vs. 10.7%, $p=0.015$), and the risk of persistent or recurrent high-grade CIN was significantly decreased in the vaccinated group (OR=0.2; 95% CI=0.1, 0.7) [44].

Based on the above results, the following was recommended:

· In women who underwent conization, the administration of the nonavalent HPV vaccine can reduce the risk of recurrence of HPV-related diseases by preventing infection with the corresponding viral subtype (IVB).

The presented recommendations are based on evidence but have several controversial aspects. Although this guideline recommends HPV vaccination in middle-aged women, the optimal age for vaccination is between 9 and 26 years in girls and women. In women aged 27–45 years, vaccination safety has been sufficiently proven, but the vaccine may be less effective because these women may have already been exposed to the virus [45]. In addition, because only one RCT has been performed for the 9-valent vaccine in middle-aged women, the evidence is insufficient for this vaccine. Therefore, doctors should fully discuss the benefits of vaccination with the patient prior to vaccination. Moreover, further research is required to identify middle-aged women who can benefit from vaccination and to study the cost-effectiveness of vaccination in this population.

HPV vaccines are administered via a 2-dose schedule for most persons who initiate vaccination at 9–14 years of age. Based on several RCTs, 2 doses of the 4-valent and 2-valent vaccines are recommended for boys and girls aged 9–13 years and 9–14 years, respectively [21 46 47].

This guideline recommends a 2-valent vaccine for boys and men aged 9–25 years. However, only one phase I/II randomized study has evaluated immunogenicity in boys aged 10–18 years [7]. Owing to insufficient evidence, further studies on 2-valent HPV vaccines in adolescent boys and young adult men are required.

These guidelines recommend that the administration of the HPV vaccine can lower the risk of infection with an HPV subtype other than the ones causing the infection in women already infected with HPV. This must be interpreted carefully, as it can be misunderstood as the vaccine leading to HPV clearance in women after infection. Clinical studies have demonstrated that vaccination does not eliminate HPV already present in women [12 35 36]. In addition, evidence to support the effectiveness of vaccination in patients who have already undergone conization is still inconclusive. In a analysis study of an RCT, no vaccine efficacy was observed against cervical precancerous lesions in patients who underwent conization for cervical lesions after vaccination [42]. Although recent meta-analysis shows a significant risk reduction of recurrence or persistence of CIN2+ after conization [43 48 49], well-designed RCTs are needed in future.

Regarding the safety of the HPV vaccine, no associations with multiple sclerosis, other disorders of the central nervous system, demyelinating disorders, Guillain-Barré syndrome, neurological diseases, venous thromboembolism, and autoimmune diseases have been revealed in observational studies and randomized clinical studies [50 51 52 53]. In addition, in a recently published registry-based study on 11–14-year-old Korean girls, 33 serious complications, including Hashimoto's thyroiditis, rheumatoid arthritis, and malignant headache, were demonstrated to not be associated with HPV vaccination [54].

The present recommendation requires further updates based on further research on men and middle-aged women or on controversial aspects.

The authors would like to thank KSGO staff for their support throughout the whole guideline development process.

Conflict of Interest: Kyung-Jin Min and Seok Ju Seong serve as editors or editorial advisor of the Journal of Gynecologic Oncology (JGO), but have no role in the decision to publish this article. No other conflict of interest relevant to this article was reported.

Author Contributions: Conceptualization: C.H.W., M.K.J., L.J.K., L.K.H., L.J.W.

Data curation: C.H.W., M.K.J., K.S.H., K.K., K.S., S.S.J., S.Y.J., L.K.H., L.S.W., L.J.W., J.W., K.Y.T., L.J.K.

Formal analysis: C.H.W., M.K.J., K.S.H., K.K., K.S., S.S.J., S.Y.J., L.K.H., L.S.W., L.J.W., J.W.

Investigation: C.H.W., M.K.J., K.S.H., K.K., K.S., S.S.J., S.Y.J., L.K.H., L.S.W., L.J.W., J.W.

Writing - original draft: C.H.W., M.K.J.

Writing - review & editing: C.H.W., M.K.J., K.S.H., K.K., K.S., S.S.J., S.Y.J., L.K.H., L.S.W., L.J.W., J.W., K.Y.T., L.J.K.

Lyon IARC Working Group on the Evaluation of Carcinogenic Risks to Humans 2009 255 314 2
Bray F Ferlay J Laversanne M Brewster DH Gombe Mbalawa C Kohler B Cancer incidence in five continents: inclusion criteria, highlights from Volume X and the global status of cancer registration Int J Cancer 2015 137 2060 2071 26135522 10.1002/ijc.29670 3 Schiffman M Castle PE Jeronimo J Rodriguez AC Wacholder S Human papillomavirus and cervical cancer Lancet 2007 370 890 907 17826171 10.1016/S0140-6736(07)61416-0 4 Forman D de Martel C Lacey CJ Soerjomataram I Lortet-Tieulent J Bruni L Global burden of human papillomavirus and related diseases Vaccine 2012 30 Suppl 5 F12 F23 23199955 10.1016/j.vaccine.2012.07.055 5 Ferris DG Samakoses R Block SL Lazcano-Ponce E Restrepo JA Mehlsen J 4-Valent human papillomavirus (4vHPV) vaccine in preadolescents and adolescents after 10 years Pediatrics 2017 140 e20163947 29167376 10.1542/peds.2016-3947 6 Giuliano AR Palefsky JM Goldstone S Moreira ED Jr Penny ME Aranda C Efficacy of quadrivalent HPV vaccine against HPV Infection and disease in males N Engl J Med 2011 364 401 411 21288094 10.1056/NEJMoa0909537 PMC3495065 7 Petäjä T Keränen H Karppa T Kawa A Lantela S Siitari-Mattila M Immunogenicity and safety of human papillomavirus (HPV)-16/18 AS04-adjuvanted vaccine in healthy boys aged 10–18 years J Adolesc Health 2009 44 33 40 19101456 10.1016/j.jadohealth.2008.10.002 8 Min KJ Kwon SH Kim K Kim S Kim HJ Seong SJ Clinical guideline for 9-valent HPV vaccine: Korean Society of Gynecologic Oncology guideline J Gynecol Oncol 2019 30 e31 30740959 10.3802/jgo.2019.30.e31 PMC6393630 9 Konno R Shin HR Kim YT Song YS Sasagawa T Inoue M Human papillomavirus infection and cervical cancer prevention in Japan and Korea Vaccine 2008 26 Suppl 12 M30 M42 18945412 10.1016/j.vaccine.2008.05.006 10 Ault KA Future II Study Group Effect of prophylactic human papillomavirus L1 virus-like-particle vaccine on risk of cervical intraepithelial neoplasia grade 2, grade 3, and adenocarcinoma in situ: a combined analysis of four randomised clinical trials Lancet 2007 369 1861 1868 17544766 10.1016/S0140-6736(07)60852-6 11 Garland SM Kjaer SK Muñoz N Block SL Brown DR DiNubile MJ Impact and effectiveness of the quadrivalent human papillomavirus vaccine: a systematic review of 10 years of real-world experience Clin Infect Dis 2016 63 519 527 27230391 10.1093/cid/ciw354 PMC4967609 12 Giuliano AR Joura EA Garland SM Huh WK Iversen OE Kjaer SK Nine-valent HPV vaccine efficacy against related diseases and definitive therapy: comparison with historic placebo population Gynecol Oncol 2019 154 110 117 30982556 10.1016/j.ygyno.2019.03.253 13 Herrero R Quint W Hildesheim A Gonzalez P Struijk L Katki HA Reduced prevalence of oral human papillomavirus (HPV) 4 years after bivalent HPV vaccination in a randomized clinical trial in Costa Rica PLoS One 2013 8 e68329 23873171 10.1371/journal.pone.0068329 PMC3714284 14 Olsson SE Kjaer SK Sigurdsson K Iversen OE Hernandez-Avila M Wheeler CM Evaluation of quadrivalent HPV 6/11/16/18 vaccine efficacy against cervical and anogenital disease in subjects with serological evidence of prior vaccine type HPV infection Hum Vaccin 2009 5 696 704 19855170 10.4161/hv.5.10.9515 15 Schlecht NF Masika M Diaz A Nucci-Sack A Salandy A Pickering S Risk of oral human papillomavirus infection among sexually active female adolescents receiving the quadrivalent vaccine JAMA Netw Open 2019 2 e1914031 31651968 10.1001/jamanetworkopen.2019.14031 PMC6822084 16 Wilkin TJ Chen H Cespedes MS Leon-Cruz JT Godfrey C Chiao EY A randomized, placebo-controlled trial of the quadrivalent human papillomavirus vaccine in human immunodeficiency virus-infected adults aged 27 years or older: AIDS Clinical Trials Group Protocol A5298 Clin Infect Dis 2018 67 1339 1346 29659751 10.1093/cid/ciy274 PMC6186857 17 Schiller JT Castellsagué X Garland SM A review of clinical trials of human papillomavirus prophylactic vaccines Vaccine 2012 30 Suppl 5

F123 F138 23199956 10.1016/j.vaccine.2012.04.108 PMC4636904 18 Vesikari T Brodzski N van Damme P Diez-Domingo J Icardi G Petersen LK A randomized, double-blind, phase III study of the immunogenicity and safety of a 9-valent human papillomavirus L1 virus-like particle vaccine (V503) versus Gardasil® in 9-15-year-old girls *Pediatr Infect Dis J* 2015 34 992 998 26090572 10.1097/INF.0000000000000773 19 Committee KGCP Clinical guideline for bi-valent HPV vaccine: Korean Society of Gynecologic Oncology guidelines Seoul Korean Society of Gynecologic Oncology 2011 20 Committee KGCP Clinical guideline for quadri-valent HPV vaccine: Korean Society of Gynecologic Oncology guidelines Seoul Korean Society of Gynecologic Oncology 2011 21 Min KJ Kwon SH Kim S Kim HJ Seong SJ Song YJ Preventive vaccination against cervical cancer: Korean Society of Gynecologic Oncology guideline *J Gynecol Oncol* 2016 27 e30 27029751 10.3802/jgo.2016.27.e30 PMC4823361 22 Muñoz N Manalastas R Jr Pitisuttithum P Tresukosol D Monsonego J Ault K Safety, immunogenicity, and efficacy of quadrivalent human papillomavirus (types 6, 11, 16, 18) recombinant vaccine in women aged 24–45 years: a randomised, double-blind trial *Lancet* 2009 373 1949 1957 19493565 10.1016/S0140-6736(09)60691-7 23 Castellsagué X Muñoz N Pitisuttithum P Ferris D Monsonego J Ault K End-of-study safety, immunogenicity, and efficacy of quadrivalent HPV (types 6, 11, 16, 18) recombinant vaccine in adult women 24–45 years of age *Br J Cancer* 2011 105 28 37 21629249 10.1038/bjc.2011.185 PMC3137403 24 Luna J Plata M Gonzalez M Correa A Maldonado I Nossa C Long-term follow-up observation of the safety, immunogenicity, and effectiveness of Gardasil™ in adult women *PLoS One* 2013 8 e83431 24391768 10.1371/journal.pone.0083431 PMC3877052 25 Chen W Zhao Y Xie X Liu J Li J Zhao C Safety of a quadrivalent human papillomavirus vaccine in a phase 3, randomized, double-blind, placebo-controlled clinical trial among Chinese women during 90 months of follow-up *Vaccine* 2019 37 889 897 30638797 10.1016/j.vaccine.2018.12.030 26 Wei L Xie X Liu J Zhao Y Chen W Zhao C Efficacy of quadrivalent human papillomavirus vaccine against persistent infection and genital disease in Chinese women: a randomized, placebo-controlled trial with 78-month follow-up *Vaccine* 2019 37 3617 3624 30122646 10.1016/j.vaccine.2018.08.009 27 Palefsky JM Giuliano AR Goldstone S Moreira ED Jr Aranda C Jessen H HPV vaccine against anal HPV infection and anal intraepithelial neoplasia *N Engl J Med* 2011 365 1576 1585 22029979 10.1056/NEJMoa1010971 28 Bhatla N Suri V Basu P Shastri S Datta SK Bi D Immunogenicity and safety of human papillomavirus-16/18 AS04-adjuvanted cervical cancer vaccine in healthy Indian women *J Obstet Gynaecol Res* 2010 36 123 132 20178538 10.1111/j.1447-0756.2009.01167.x 29 Einstein MH Takacs P Chatterjee A Sperling RS Chakhtoura N Blatter MM Comparison of long-term immunogenicity and safety of human papillomavirus (HPV)-16/18 AS04-adjuvanted vaccine and HPV-6/11/16/18 vaccine in healthy women aged 18–45 years: end-of-study analysis of a phase III randomized trial *Hum Vaccin Immunother* 2014 10 3435 3445 25483701 10.4161/hv.36121 PMC4514070 30 Skinner SR Szarewski A Romanowski B Garland SM Lazcano-Ponce E Salmerón J Efficacy, safety, and immunogenicity of the human papillomavirus 16/18 AS04-adjuvanted vaccine in women older than 25 years: 4-year interim follow-up of the phase 3, double-blind, randomised controlled VIVIANE study *Lancet* 2014 384 2213 2227 25189358 10.1016/S0140-6736(14)60920-X 31 Schwarz TF Spaczynski M Schneider A Wysocki J Galaj A Perona P Immunogenicity and tolerability of an HPV-16/18 AS04-adjuvanted prophylactic cervical cancer vaccine in women aged 15–55 years *Vaccine* 2009 27 581 587 19022320 10.1016/j.vaccine.2008.10.088 32 Schwarz T Spaczynski M Kaufmann A Wysocki J Galaj A Schulze K Persistence of immune responses to the HPV-16/18 AS04-adjuvanted vaccine in women aged 15–55 years and first-time modelling of antibody responses in mature women: results from an open-label 6-year follow-up study *BJOG* 2015 122 107 118 25208608 10.1111/1471-0528.13070 PMC4489326 33 Schwarz TF Spaczynski M Schneider A Wysocki J Galaj A Schulze K Persistence of immune response to HPV-16/18 AS04-adjuvanted cervical cancer vaccine in women aged 15–55 years *Hum Vaccin* 2011 7 958 965 21892005 10.4161/hv.7.9.15999 PMC3225769 34 Joura EA Ulied A Vandermeulen C Rua Figueroa M Seppa I Hernandez Aguado JJ Immunogenicity and safety of a nine-valent human papillomavirus vaccine in women 27–45 years of age compared to women 16-26 years of age: an open-label phase 3 study *Vaccine* 2021 39 2800 2809 33676783 10.1016/j.vaccine.2021.01.074 35 Scherer EM Smith RA Gallego DF Carter JJ Wipf GC Hoyos M A single human papillomavirus vaccine dose improves B cell memory in previously infected subjects *EBioMedicine* 2016 10 55 64 27423190 10.1016/j.ebiom.2016.06.042 PMC5006801 36 Szarewski A Poppe WA Skinner SR Wheeler CM Paavonen J Naud P Efficacy of the human papillomavirus (HPV)-16/18 AS04-adjuvanted vaccine in

women aged 15–25 years with and without serological evidence of previous exposure to HPV-16/18 *Int J Cancer* 2012 131 106 116 21858807 10.1002/ijc.26362 37 Karimi-Zarchi M Allahqoli L Nehmati A Kashi AM Taghipour-Zahir S Alkatout I Can the prophylactic quadrivalent HPV vaccine be used as a therapeutic agent in women with CIN? A randomized trial *BMC Public Health* 2020 20 274 32106837 10.1186/s12889-020-8371-z PMC7045378 38 Ghelardi A Parazzini F Martella F Pieralli A Bay P Tonetti A SPERANZA project: HPV vaccination after treatment for CIN2 *Gynecol Oncol* 2018 151 229 234 30197061 10.1016/j.ygyno.2018.08.033 39 Kang WD Choi HS Kim SM Is vaccination with quadrivalent HPV vaccine after loop electrosurgical excision procedure effective in preventing recurrence in patients with high-grade cervical intraepithelial neoplasia (CIN2-3)? *Gynecol Oncol* 2013 130 264 268 23623831 10.1016/j.ygyno.2013.04.050 40 Petrillo M Dessole M Tinacci E Saderi L Muresu N Capobianco G Efficacy of HPV vaccination in women receiving LEEP for cervical dysplasia: a single institution's experience *Vaccines (Basel)* 2020 8 45 10.3390/vaccines8010045 PMC7157656 31991753 41 Garland SM Paavonen J Jaisamrarn U Naud P Salmerón J Chow SN Prior human papillomavirus-16/18 AS04-adjuvanted vaccination prevents recurrent high grade cervical intraepithelial neoplasia after definitive surgical therapy: post-hoc analysis from a randomized controlled trial *Int J Cancer* 2016 139 2812 2826 27541373 10.1002/ijc.30391 PMC5412942 42 Hildesheim A Gonzalez P Kreimer AR Wacholder S Schussler J Rodriguez AC Impact of human papillomavirus (HPV) 16 and 18 vaccination on prevalent infections and rates of cervical lesions after excisional treatment *Am J Obstet Gynecol* 2016 215 212.e1 212.e15 26892991 10.1016/j.ajog.2016.02.021 PMC4967374 43 Bartels HC Postle J Rogers AC Brennan D Prophylactic human papillomavirus vaccination to prevent recurrence of cervical intraepithelial neoplasia: a meta-analysis *Int J Gynecol Cancer* 2020 30 777 782 32276936 10.1136/ijgc-2020-001197 44 Del Pino M Martí C Torras I Henere C Munmany M Marimon L HPV vaccination as adjuvant to conization in women with cervical intraepithelial neoplasia: a study under real-life conditions *Vaccines (Basel)* 2020 8 245 10.3390/vaccines8020245 PMC7349984 32456136 45 Meites E Szilagyi PG Chesson HW Unger ER Romero JR Markowitz LE Human papillomavirus vaccination for adults: updated recommendations of the advisory committee on immunization practices *MMWR Morb Mortal Wkly Rep* 2019 68 698 702 31415491 10.15585/mmwr.mm6832a3 PMC6818701 46 Meites E Kempe A Markowitz LE Use of a 2-dose schedule for human papillomavirus vaccination - updated recommendations of the advisory committee on immunization practices *MMWR Morb Mortal Wkly Rep* 2016 65 1405 1408 27977643 10.15585/mmwr.mm6549a5 47 World Health Organization Human papillomavirus vaccines: WHO position paper, May 2017-recommendations *Vaccine* 2017 35 5753 5755 28596091 10.1016/j.vaccine.2017.05.069 48 Jentschke M Kampers J Becker J Sibbertsen P Hillemanns P Prophylactic HPV vaccination after conization: a systematic review and meta-analysis *Vaccine* 2020 38 6402 6409 32762871 10.1016/j.vaccine.2020.07.055 49 Lichter K Krause D Xu J Tsai SHL Hage C Weston E Adjuvant human papillomavirus vaccine to reduce recurrent cervical dysplasia in unvaccinated women: a systematic review and meta-analysis *Obstet Gynecol* 2020 135 1070 1083 32282601 10.1097/AOG.0000000000003833 50 Scheller NM Svanström H Pasternak B Arnheim-Dahlström L Sundström K Fink K Quadrivalent HPV vaccination and risk of multiple sclerosis and other demyelinating diseases of the central nervous system *JAMA* 2015 313 54 61 25562266 10.1001/jama.2014.16946 51 Ojha RP Jackson BE Tota JE Offutt-Powell TN Singh KP Bae S Guillain-Barre syndrome following quadrivalent human papillomavirus vaccination among vaccine-eligible individuals in the United States *Hum Vaccin Immunother* 2014 10 232 237 24013368 10.4161/hv.26292 PMC4181024 52 Arnheim-Dahlström L Pasternak B Svanström H Sparén P Hviid A Autoimmune, neurological, and venous thromboembolic adverse events after immunisation of adolescent girls with quadrivalent human papillomavirus vaccine in Denmark and Sweden: cohort study *BMJ* 2013 347 f5906 24108159 10.1136/bmj.f5906 PMC3805482 53 Leung TF Liu AP Lim FS Thollot F Oh HM Lee BW Comparative immunogenicity and safety of human papillomavirus (HPV)-16/18 AS04-adjuvanted vaccine and HPV-6/11/16/18 vaccine administered according to 2- and 3-dose schedules in girls aged 9–14 years: results to month 12 from a randomized trial *Hum Vaccin Immunother* 2015 11 1689 1702 26062002 10.1080/21645515.2015.1050570 PMC4514190 54 Yoon D Lee JH Lee H Shin JY Association between human papillomavirus vaccination and serious adverse events in South Korean adolescent girls: nationwide cohort study *BMJ* 2021 372 m4931 33514507 10.1136/bmj.m4931 PMC8030229 SUPPLEMENTARY MATERIALS Data S1 Development process of human papillomavirus (HPV) vaccine guideline

